

University of Rajshahi

Rajshahi-6205

Bangladesh.

RUCL Institutional Repository

<http://rulrepository.ru.ac.bd>

---

Department of Mathematics

MPhil Thesis

---

2011

# On Supra Fuzzy Topological Spaces

Hoque, Md. Fazlul

University of Rajshahi

---

<http://rulrepository.ru.ac.bd/handle/123456789/883>

*Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.*

# On Supra Fuzzy Topological Spaces



*A Thesis Submitted to the Department of Mathematics, University of Rajshahi, Rajshahi – 6205, Bangladesh, for the Degree of Master of Philosophy in Mathematics*

By

**MD. FAZLUL HOQUE**

Under the Supervision of  
**Professor Dr. Dewan Muslim Ali**  
Department of Mathematics  
University of Rajshahi  
Rajshahi-6205, Bangladesh


**June, 2011**

---

**Dedicated  
To  
My Beloved  
Parents**

## STATEMENT OF ORIGINALITY

I declare that the contents in my M. Phill. Thesis entitled “**On Supra Fuzzy Topological Spaces**” is original and accurate to the best of my knowledge. I also declare that the materials contained in my research work have not been previously published or written any person for degree or diploma.

  
28.06.2011  
(Md. Fazlul Hoque)  
M. Phill. Research Fellow  
Rajshahi University  
June, 2001

**Dr. Dewan Muslim Ali**  
M.Sc. (Raj) Ph.D. (Benaras)  
Professor  
Department of Mathematics



University of Rajshahi  
Rajshahi-6205, Bangladesh  
Phone: (Dept.) 750041-9/4108  
(Resi): (0721) 750528  
Fax: 0088-0721-750064  
E-mail: profdma@ru.ac.bd

No .....

Date .....

## CERTIFICATE

I have the pleasure in certifying that the M. Phill. Thesis entitled “On Supra Fuzzy Topological Spaces” submitted by Md. Fazlul Hoque in fulfillment of the requirement for the degree of M. Phill. in Mathematics, University of Rajshahi, Rajshahi-6205, Bangladesh has been completed under my supervision. I believe that the research work is an original one and it has not been submitted elsewhere for any degree.

I wish him a bright future and every success in life.

Supervisor

*Muslim*  
28.6.11

(Dr. Dewan Muslim Ali)  
Professor  
Department of Mathematics  
University of Rajshahi  
Rajshahi - 6205, Bangladesh.

# ACKNOWLEDGEMENT

Firstly, I give my limitless thanks to the Almighty Allah for giving me strength, endurance and ability to complete this course of study.

I would like to convey an immense grateful to a number of individuals for their kind advices, directions, suggestions and co-operation, which have enabled me to complete my thesis successfully.

I am bound to express my deepest respect and thanks to my respected supervisor Professor Dr. Dewan Muslim Ali, Department of Mathematics, University of Rajshahi, Bangladesh for his day-to-day monitoring, supervision and maximum assistance for the completion of the present study successfully.

I wish to thank the Chairman, Department of Mathematics, Rajshahi University, for providing necessary laboratory and other facilities during the course of thesis work.

I am thankful to Md. Sahadat Hossain, Assistant Professor, Department of Mathematics, Rajshahi University, for providing me the source of information and materials about my study. I also wish to express my sincere admiration, appreciation and gratitude to all the teachers of the Department of Mathematics, University of Rajshahi for providing me with invaluable suggestions during the period of my research work.

A special thank will go to the shoulders of my colleagues and friends at Pabna Science and Technology University, Pabna, for their friendly support to overcome the hurdles during the M. Phill. program.

Finally, I am ever grateful to my parents for their moral and constant blessing, good advice and encouragement. I would like to thank all of my family members and well wishers for their inspiration.

I fine, I am alone responsible for the errors and shortcomings in this study if there be any, I am sorry for that.

(Md. Fazlul Hoque)  
Department of Mathematics  
June, 2011

# CONTENTS

	<b>Page no.</b>
<b>Introduction</b>	1 - 3
<b>Chapter 1 Preliminaries</b>	4 - 16
<b>Chapter 2 <math>T_0</math> Spaces in Supra Fuzzy Topology</b>	17 - 28
<b>Chapter 3 <math>T_1</math> Spaces in Supra Fuzzy Topology</b>	29 - 39
<b>Chapter 4 <math>T_2</math> Spaces in Supra Fuzzy Topology</b>	40 - 51
<b>Chapter 5 Regular Spaces in Supra Fuzzy Topology</b>	52 - 60
<b>Chapter 6 Normal Spaces in Supra Fuzzy Topology</b>	61 - 66
<b>Chapter 7 <math>R_0</math> and <math>R_1</math> Spaces in Supra Fuzzy Topology</b>	67 - 77
<b>Bibliography</b>	78 - 87

# Introduction

The fundamental concept of a fuzzy set was introduced by L. A. Zadeh[111] in 1965 to provide a foundation for the development of many areas of knowledge. Consequently, this provides a natural frame work for generalizing many algebraic and topological concepts in various directions such as fuzzy groups, fuzzy rings, fuzzy vector spaces, fuzzy supra topology, fuzzy infra topology, fuzzy bitopology etc. many other branches of mathematics have been developed all over the world during the last five decades. In 1968, Chang [19] introduced the concepts of a fuzzy topological space by using the fuzzy set. Wong [105], Lowen[60], Hutton[48], Katsaras[52], Ali[3], Pu and Liu[72], etc., discussed various aspects of fuzzy topological spaces. Ying [74] introduced fuzzifying topology and developed this in a new direction with the semantic methods of continuous valued logic. In the frame work of fuzzifying topology, Sinha[93] introduced and studied  $T_0$ -,  $T_1$ -,  $T_2$ (Hausdorff)-,  $T_3$ (regular)-,  $T_4$ (normal)-, separation axioms. A.S. Mashhour et al.[64] introduced and studied the concepts of the family of fuzzifying semi-open sets, fuzzifying neighbourhood structure of a point and fuzzifying semi-closure of a fuzzy set. A.S. Mashhour et al.[64] also introduced and studied the  $R_0$  and  $R_1$  separation axioms and studied their relations with the  $T_1$  and  $T_2$ - separation axioms respectively. Also in fuzzifying topology they introduced and studied semi- $T_0$ -, semi- $R_0$ -, semi- $T_1$ -, semi- $R_1$ -, semi- $T_2$ (semi Hausdorff)-, semi- $T_3$ (semi regular)-, semi- $T_4$ (semi normal)-, separation axioms. In 1983, A.S. Mashhour et al.[64] introduced supra topological spaces and studies  $s$ -continuous functions and  $s^*$ - continuous functions. In 1987, M.E. Abd EL-Monsef et al. [1] introduced the fuzzy supra topological spaces and studied fuzzy supra continuous functions and characterized a number of basic concepts.

The purpose of this thesis is to introduce some new definitions of separation axioms in supra fuzzy topological spaces using the ideas of Ali [8]. Some of their equivalent formulations along with various new characterizations and results concerning the existing ones are presented here. Our criterion for definitions has been preserving as



much as possible the relation between the corresponding separation properties for supra fuzzy topological spaces. Moreover, it will be seen that the definitions of these axioms are 'good extensions' in the sense of Lowen [60].

We aim to develop theories of supra fuzzy  $T_0$ , supra fuzzy  $T_1$ , supra fuzzy  $T_2$ (Hausdorff), supra fuzzy SFR(supra fuzzy regular), supra fuzzy SFN(supra fuzzy normal), supra fuzzy  $R_0$ - and supra fuzzy  $R_1$ - separation axioms analogous to its counterpart in ordinary topology. The materials of this thesis have been divided into seven chapters, a brief scenario of which we present as follows.

Chapter one incorporates some of the basic definitions and results of fuzzy set, fuzzy topology, mapping, supra fuzzy topology and its mapping. These results are ready references for the work in the subsequent chapter. Results are stated without proof and can be seen in the papers referred to.

Our work starts from second chapter. In second chapter, we have introduced and studied  $T_0$  properties in supra fuzzy topological spaces. Here we add four more definitions to this list and we have established relations among them. All these four definitions are 'good extensions' of the corresponding concept  $T_0$  in a topological space. We prove that all the definitions are hereditary, productive and projective. Also we have studied some other properties of these concepts.

In third chapter, we have introduced and studied  $T_1$  properties in supra fuzzy topological spaces. Here we add four more definitions to this list and we have established relations among them. All these four definitions are 'good extensions' of the corresponding concept  $T_1$  in a topological space. We prove that all these concepts are hereditary, productive and projective. Also we have studied some other properties of these concepts.

We have introduced and studied  $T_2$ (Hausdorff) properties for supra fuzzy topological spaces, in chapter four. We have introduced here four more definitions and we

established relations among them. All these properties are 'good extensions' of the corresponding concept  $T_2$ (Hausdorff) in a topological space. We have observed that all the definitions are hereditary, productive and projective. Also we have studied some other properties of these concepts.

In fifth chapter, we have discussed SFR (supra fuzzy regular) properties in supra fuzzy topological spaces. Here we have obtained four more definitions and we established relations among them. We see that all these properties are 'good extensions' of the corresponding concept  $T_3$  (regularity) in topological spaces. We have discussed that all the properties are hereditary, productive and projective. We have also studied several other properties of these concepts.

In chapter six, we have introduced and studied SFN (supra fuzzy normal) properties in supra fuzzy topological spaces. We have given here three more definitions and we have established relations among them. All these are 'good extensions' of the corresponding concept  $T_4$  (normality) in a topological space. Some other pleasant properties of these concepts have been studied here.

In seventh chapter, we aim to introduce and study  $R_0$  and  $R_1$  properties in supra fuzzy topological spaces. Four more definitions of each concept are given and obtained implications among them. All these are found to be 'good extensions' of the corresponding concepts  $R_0$  and  $R_1$  in a topological space. We have found that all the concepts are hereditary, productive and projective. Some other properties of these concepts are also found.

# CHAPTER-1

## Preliminaries

### 1.1 Introduction:

In this chapter, we collect several basic definitions and results of the Fuzzy sets, Grade of membership, complement of fuzzy sets, some laws of fuzzy sets, different mappings on fuzzy sets, Fuzzy topological spaces, Supra fuzzy topological spaces, distinction between Fuzzy sets and Supra fuzzy sets,  $T_0$ ,  $T_1$ ,  $T_2$  –spaces, Fuzzy product topological spaces and Supra fuzzy product topological spaces which are to be used as ready references for understanding the subsequent chapters. Most of the results are quoted from various research papers. We make use the following general notations in this thesis.

- $\Lambda$  : Index set.
- $I = [0, 1]$  : Closed unit interval.
- $I_1 = [0, 1)$  : Right open unit interval.
- $I_0 = (0, 1]$  : Left open unit interval.
- $u, v, w, \dots$  : Supra Fuzzy sets.
- $(X, t)$  : Fuzzy topological space.
- $(X, t^*)$  : Supra Fuzzy topological space.
- $(X, T^*)$  : General supra topological space.
- $(X, T)$  : General topological space.
- $\prod_{i \in \Lambda} X_i$  : Usual product of  $X_i$ .

$(X, t^*_1 \times t^*_2)$ : Product of supra fuzzy topologies  $t^*_1$  and  $t^*_2$  on the set  $X$ .

$I_\alpha(t^*) = \{ u^{-1}(\alpha, 1] : u \in t^* \}, \alpha \in I_1$ : General supra topology on  $X$ .

**Definition 1.2[111]:** For a set  $X$ , a function  $u: X \rightarrow [0,1]$  is called a fuzzy set in  $X$ . For every  $x \in X$ ,  $u(x)$  represents the grade of membership of  $x$  in the fuzzy set  $u$ . Some authors say that  $u$  is a fuzzy subset of  $X$ . Thus a usual subset of  $X$ , is a special type of a fuzzy set in which the range of the function is  $\{0, 1\}$ .

**Definition 1.3[111]:** Let  $X$  be a nonempty set and  $A$  be a subset of  $X$ . The function

$1_A: X \rightarrow [0, 1]$  defined by  $1_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$  is called the characteristic

function of  $A$ . We also write  $1_x$  for the characteristic function of  $\{x\}$ .

**Definition 1.4[111]:** The characteristic functions of subsets of a set  $X$  are referred to as the crisp sets in  $X$ .

**Example 1.5:** Suppose  $X$  is real number  $R$  and the fuzzy set of real numbers much greater than 5 in  $X$  that could be defined by the continuous function  $U: X \rightarrow [0,1]$  such that

$$u(x) = \begin{cases} 0 & \text{if } x \leq 5 \\ \frac{x-5}{50} & \text{if } 5 < x < 55. \\ 1 & \text{if } x \geq 55 \end{cases}$$

**Definition 1.6[75]:** A fuzzy subset is empty if and only if grade of membership is identically zero in  $X$ . It is denoted by  $0$ .

**Definition 1.7[75]:** A fuzzy subset is whole if and only if its grade of membership is identically one in  $X$ . It is denoted by  $1$ .

**Definition 1.8[111]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then  $u$  is said to be subset of  $v$ , i.e.,  $u \subseteq v$  if and only if  $u(x) \leq v(x)$  for every  $x \in X$ .

**Definition 1.9[111]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then  $u$  is said to be equal to  $v$ , i.e.,  $u = v$  if and only if  $u(x) = v(x)$  for every  $x \in X$ .

**Definition 1.10[111]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then  $u$  is said to be the complement of  $v$ , i.e.,  $u = v^c$  if and only if  $u(x) = 1 - v(x)$ , for every  $x \in X$ . Obviously,  $(v^c)^c = v$ .

**Definition 1.11[19]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then the union  $w$  of  $u$  and  $v$ , i.e.,  $w = u \cup v$  if and only if  $w(x) = (u \cup v)(x) = \max \{ u(x), v(x) \}$ , for every  $x \in X$ . The union  $w$  is a fuzzy subset of  $X$ .

In general, if  $\Lambda$  be an index set and  $A = \{u_i : i \in \Lambda\}$  be a family of fuzzy sets of  $X$  then the union  $\cup u_i$  is defined by  $(\cup u_i)(x) = \sup \{u_i(x) : i \in \Lambda\}$ ,  $x \in X$ .

**Definition 1.12[19]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then the intersection  $m$  of  $u$  and  $v$ , i.e.,  $m = u \cap v$  is a fuzzy subset of  $X$  if and only if  $m(x) = (u \cap v)(x) = \min \{u(x), v(x)\}$ ,  $\forall x \in X$ , and  $(\cap u_i)(x) = \inf \{u_i(x) : i \in \Lambda\}$ ,  $x \in X$ , where  $\{u_i, i \in \Lambda\}$ .

**Definition 1.13[3]:** Let  $u$  and  $v$  be two fuzzy subsets of a set  $X$ . Then the difference of  $u$  and  $v$  is defined by  $u - v = u \cap v^c$ .

**Definition 1.14[3]:** If  $\alpha \in I$  and  $u \in I^X$  define by  $u(x) = \alpha$ , for all  $x \in X$ , we refer to  $u$  as a constant fuzzy set and denote it by  $\alpha$  itself. In particular, we have the constant fuzzy sets 0 and 1.

**Example 1.15:** Let  $X = \{x, y, z\}$  and  $u, v \in I^X$  are defined by  $u(x) = .6$ ,  $u(y) = .7$ ,  $u(z) = .5$  and  $v(x) = .7$ ,  $v(y) = .5$ ,  $v(z) = .4$ . Then  $(u \cup v)(x) = \max \{u(x), v(x)\} = .7$ ,  $(u \cup v)(y) = .7$ ,  $(u \cup v)(z) = .5$ .

$$\begin{aligned}
&= \max \{u(y), v(y)\} = .7, (u \cup v)(y) = \max \{u(y), v(y)\} = .5, (u \cap v)(x) = \min \{u(x), v(x)\} \\
&= .6, (u \cap v)(y) = \min \{u(y), v(y)\} = .5, (u \cap v)(z) = \min \{u(z), v(z)\} = .4., u^c(x) \\
&= 1-u(x) = .4, u^c(y) = 1-u(y) = .3, u^c(z) = 1-u(z) = .5.
\end{aligned}$$

### Laws of the algebra of fuzzy sets 1.16:

As in ordinary set theory, idempotent laws, associative law, commutative law, distributive laws, identity law, demorgan's laws hold in the case of fuzzy sets also. But the complement laws are not necessarily true. For example, if  $X = \{a, b, c\}$  and  $u$  is a fuzzy subset of  $X$  where is defined by

$$u = \{(a, .2), (b, .7), (c, 1)\},$$

$$\text{then } u^c = \{(a, .8), (b, .3), (c, 0)\}$$

$$\text{so } u \cup u^c = \{(a, .8), (b, .7), (c, 1)\} \neq 1,$$

$$u \cap u^c = \{(a, .2), (b, .3), (c, 0)\} \neq 0.$$

Also in ordinary set theory  $U \cap V = \phi$  if and only if  $U \subset V^c$ . But in fuzzy subsets reverse is not necessary true. For example if

$$v = \{(a, .6), (b, .2), (c, 0)\} \text{ then } u \subset v^c,$$

$$u \cap v = \{(a, .2), (b, .2), (c, 0)\} \neq 0.$$

**Definition 1.17[19]:** Let  $f$  be a mapping from a set  $X$  into a set  $Y$  and  $u$  is a fuzzy subset of  $X$ . Then  $f$  and  $u$  induced a fuzzy subset  $v=f(u)$  of  $Y$  whose membership function is defined by

$$\begin{aligned}
v(y) &= (f(u))(y) = \sup \{u(x)\} \text{ if } f^{-1}[\{y\}] \neq \phi, x \in X \\
&= 0, \text{ otherwise.}
\end{aligned}$$

**Definition 1.18[19]:** Let  $f$  be a mapping from a set  $X$  into  $Y$  and  $v$  be a fuzzy subset of  $Y$ . Then the inverse of  $v$  written as  $u = f^{-1}(v)$  is a fuzzy subset of  $X$  and is defined by

$$u(x) = (f^{-1}(v))(x) = v(f(x)), \text{ for } x \in X.$$

**Example 1.19:** Suppose that  $X = \{x, y, z, w\}$  and  $Y = \{a, b, c\}$ . Define  $f : X \longrightarrow Y$  by  $f(x)=b, f(y)=c, f(z)=a, f(w)=a$ . Let  $u \in I^X$  be given by  $u(x) = .2, u(y) = .3, u(z) = .5$  and  $u(w) = .4$ . Then  $(f(u))(a) = \sup \{u(z), u(w)\} = .5$ . Similarly,  $(f(u))(b) = \sup u(x) = .2$ , and  $(f(u))(c) = \sup u(y) = .3$ .

On the other hand, if  $v$  is a fuzzy set in  $Y$  given by  $v(a) = .6, v(b) = .8, v(c) = .7$ . Then  $(f^{-1}(v))(x) = v(f(x))=v(b)=.8, (f^{-1}(v))(y) = v(f(y))=v(c)=.7, (f^{-1}(v))(z) = v(f(z))=v(a)=.6, (f^{-1}(v))(w) = v(f(w))=v(a)=.6$ .

We now mention some properties of fuzzy subsets induced by mappings.

Let  $f$  be a mapping from  $X$  into  $Y$ ,  $u$  be a fuzzy subset of  $X$  and  $v$  be a fuzzy subset of  $Y$ .

Then the following properties are true [19].

- (a)  $f^{-1}(v^c) = (f^{-1}(v))^c$  for any fuzzy subset  $v$  of  $Y$ .
- (b)  $f(u^c) = (f(u))^c$  for any fuzzy subset  $u$  of  $X$ .
- (c)  $v_1 \subset v_2 \Rightarrow f^{-1}(v_1) \subset f^{-1}(v_2)$ , where  $v_1$  and  $v_2$  are two fuzzy subsets of  $Y$ .
- (d)  $u_1 \subset u_2 \Rightarrow f(u_1) \subset f(u_2)$ , where  $u_1$  and  $u_2$  are two fuzzy subsets of  $X$ .
- (e)  $v \supset f(f^{-1}(v))$ , for any fuzzy subset  $v$  of  $Y$ .
- (f)  $u \subset f^{-1}(f(u))$ , for any fuzzy subset  $u$  of  $X$ .
- (g) Let  $f$  be a function from  $X$  into  $Y$  and  $g$  be a function from  $Y$  into  $Z$ . Then  $(g \circ f)^{-1}(w) = f^{-1}(g^{-1}(w))$ , for any fuzzy subset  $w$  in  $Z$ , where  $(g \circ f)$  is the composition of  $g$  and  $f$ .

**Definition 1.20[72] :** A fuzzy point in  $X$  is a special type of fuzzy set in  $X$  with membership function  $p(x) = r, p(y) = 0, \forall y \neq x$ , where  $0 < r < 1$ . This fuzzy point is said to have support  $x$  and value  $r$  and this point is denoted by  $x_r$  or  $r1_x$ .

**Definition 1.21[72]** : A fuzzy point  $p$  is said to belong a fuzzy set  $u$  in  $X$  ( $p \in u$ ) if and only if  $p(x) < u(x)$  and  $p(y) \leq u(y) \quad \forall y \neq x$ . Evidently, every fuzzy set  $u$  can be expressed as the union of all the fuzzy points which belong to  $u$ .

**Definition 1.22[72]** : Two fuzzy sets  $u$  and  $v$  in  $X$  are said to be intersected if and only if there exist a point  $x \in X$  such that  $(u \cap v)(x) \neq 0$ . In this case we say that  $u$  and  $v$  intersect at  $x$ .

**Definition 1.23[3]**: Let  $X$  be a non empty set and  $u$  be a fuzzy set in  $X$ . A  $\alpha$ - cut of  $u$  is defined by  $\alpha_u = \{x : u(x) \geq \alpha, \alpha \in I\}$ .

**Definition 1.24[3]**: Let  $X$  be a non empty set and  $u$  be a fuzzy set in  $X$ . A strong  $\alpha$ - cut of  $u$  is defined by  $\alpha_{+u} = \{x : u(x) > \alpha, \alpha \in I\}$ . We see that  $\alpha$ -cut and strong  $\alpha$ -cut are crisp subsets of  $X$ .

**Definition 1.25[3]**: Let  $X$  be a non empty set and  $u$  be a fuzzy set in  $X$ . The support of  $u$  in  $X$  is the crisp subset of  $X$  that contains all the elements of  $X$  that have non-zero membership grads in  $u$ , i.e.,  $\text{supp } u = \{x : u(x) > 0\}$ . The 1-cut is called the core of  $u$ .

**Definition 1.26[3]**: The height  $h(u)$  of a fuzzy set  $u$  is the largest membership grade obtained by any element in that set, i.e.,  $h(u) = \sup_{x \in X} u(x)$ .

**Definition 1.27[3]**: For a finite fuzzy set, the cardinality  $|\alpha|$  defined as  $|\alpha| = \sum_{x \in X} \alpha(x)$ .

$\|\alpha\| = \frac{|\alpha|}{|X|}$  is called the relative cardinality of  $\alpha$ .



**Definition 1.28[3]:** A fuzzy set  $u$  is called normal when  $h(u) = 1$ ; it is called subnormal when  $h(u) < 1$ . The height of  $u$  may also be viewed as the supremum of  $\alpha$  for which  $\alpha_u \neq \phi$ .

**Definition 1.29[19]:** Let  $X$  be a non empty set and  $t$  be the collection of fuzzy sets in  $X$ . Then  $t$  is called a fuzzy topology on  $X$  if it satisfies the following conditions:

- (i)  $1, 0 \in t$ ,
- (ii) If  $u_i \in t$  for each  $i \in \Lambda$ , then  $\cup_{i \in \Lambda} u_i \in t$ .
- (iii) If  $u_1, u_2 \in t$  then  $u_1 \cap u_2 \in t$ .

If  $t$  is a fuzzy topology on  $X$ , then the pair  $(X, t)$  is called a fuzzy topological space (fts, in short) and members of  $t$  are called  $t$ -open (or simply open) fuzzy sets. If  $u$  is open fuzzy set, then the fuzzy sets of the form  $1-u$  are called  $t$ -closed (or simply closed) fuzzy sets.

**Definition 1.30[60]:** Let  $X$  be a nonempty set and  $t$  be a collection of fuzzy sets in  $X$  such that

- (i)  $1, 0 \in t$ ,
- (ii) If  $u_i \in t$  for each  $i \in \Lambda$ , then  $\cup_{i \in \Lambda} u_i \in t$ .
- (iii) If  $u_1, u_2 \in t$  then  $u_1 \cap u_2 \in t$ .
- (iv) all constant fuzzy sets in  $X$  belong to  $t$ .

Then  $t$  is called a fuzzy topology on  $X$ .

**Example 1.31:** Let  $X = \{a, b, c, d\}$ ,  $t = \{0, 1, u, v\}$ ,

where  $1 = \{(a, 1), (b, 1), (c, 1), (d, 1)\}$

$0 = \{(a, 0), (b, 0), (c, 0), (d, 0)\}$

$u = \{(a, .2), (b, .5), (c, .7), (d, .9)\}$

$v = \{(a, .3), (b, .5), (c, .8), (d, .95)\}$

Then  $(X, t)$  is a fuzzy topological space.

**Definition 1.32[64]:** Let  $X$  be a nonempty set. A subfamily  $t^*$  of  $X$  is said to be a supra topology on  $X$  if and only if

(i)  $1, 0 \in t^*$ ,

(ii) If  $u_i \in t^*$  for each  $i \in \Lambda$ , then  $\cup_{i \in \Lambda} u_i \in t^*$ .

Then the pair  $(X, t^*)$  is called a supra fuzzy topological spaces. The elements of  $t^*$  are called supra open sets in  $(X, t^*)$  and complement of supra open set is called supra closed set.

**Example 1.33:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = .8$ ,  $u(y) = .6$  and  $v(x) = .6$ ,  $v(y) = .8$ . Then we have  $w(x) = (u \cup v)(x) = .8$ ,  $w(y) = (u \cup v)(y) = .8$  and  $k(x) = (u \cap v)(x) = .6$ ,  $k(y) = (u \cap v)(y) = .6$ . If we consider  $t^*$  on  $X$  generated by  $\{0, u, v, w, 1\}$ , then  $t^*$  is supra fuzzy topology on  $X$  but  $t^*$  is not fuzzy topology. Thus we see that every fuzzy topology is supra fuzzy topology but the converse is not always true.

**Definition 1.34[72]:** Let  $u$  be a fuzzy set in  $(X, \tau)$ . The interior of  $u$  is defined as the union of all  $\tau$  – open sets contained in  $u$ . It is denoted by  $u^\circ$ . Evidently  $u^\circ$  is the largest open fuzzy set contained in  $u$  and  $(u^\circ)^\circ = u^\circ$ .

**Definition 1.35[72]:** The intersection of all the  $\tau$  – closed set containing  $u$  is called the closure of  $u$  denoted by  $\bar{u}$ . Obviously  $\bar{u}$  is the smallest closed set containing  $u$  and  $\overline{\bar{u}} = \bar{u}$ .

**Definition 1.36[101] :** A fuzzy set  $n$  in a fuzzy topological space  $(X, \tau)$  is called a neighborhood of a point  $x \in X$ , if and only if there exist  $u \in \tau$  such that  $u \subseteq n$  and  $u(x) = n(x) > 0$ .

**Example 1.37:** Let us consider the example 1.15, and  $n = \{ (a, .5), (b, .6), (c, .8), (d, .9) \}$ . Now  $n$  is a neighborhood of  $d \in X$ . Since  $u \in \tau$  such that  $u \subseteq n$  and  $u(d) = n(d) > 0$ ,  $n$  is a neighborhood of  $d$ . Similarly,  $n_1 = \{ (a, .7), (b, .6), (c, .75), (d, .9) \}$  is a neighborhood of  $d$ . We denoted the family of all neighborhoods of  $x$  by  $N_x$ .

**Definition 1.38[72]:** A fuzzy set  $u$  in a fuzzy topological space  $(X, \tau)$  is called a neighborhood of a fuzzy point  $x_r$  if and only if there exist a fuzzy set  $u_1 \in \tau$  such that  $x_r \in u_1 \subseteq u$ . A neighborhood  $u$  is called an open neighborhood if  $u$  is open. The family consisting of all the neighborhoods of  $x_r$  is called the system of  $x_r$ .

**Definition 1.39[73]:** Let  $(X, \tau)$  and  $(Y, s)$  be two topological spaces. Let  $\tau^*$  and  $s^*$  are associated supra topologies with  $\tau$  and  $s$  respectively. The function  $f : (X, \tau^*) \longrightarrow (Y, s^*)$  is called supra fuzzy continuous if and only if for every  $v \in s$ ,  $f^{-1}(v) \in \tau^*$ , the function  $f$  is called supra fuzzy homeomeric if and only if  $f$  is bijective and both  $f$  and  $f^{-1}$  are supra fuzzy continuous.

**Definition 1.40[64]** : The function  $f : (X, t^*) \longrightarrow (Y, s^*)$  is called supra fuzzy open if and only if for each open supra fuzzy set  $u$  in  $(X, t^*)$ ,  $f(u)$  is open supra fuzzy set in  $(Y, s^*)$ .

**Definition 1.41[64]**: The function  $f : (X, t^*) \longrightarrow (Y, s^*)$  is called supra fuzzy closed if and only if for each closed supra fuzzy set  $u$  in  $(X, t^*)$ ,  $f(u)$  is closed supra fuzzy set in  $(Y, s^*)$ .

**Proposition 1.42 ([73] Theorem 1.1)**: Let  $f : (X, t^*) \longrightarrow (Y, s^*)$  be supra fuzzy continuous function, then the following properties hold :

- (i) For every  $s^*$  – closed  $v$ ,  $f^{-1}(v)$  is  $t^*$  – closed.
- (ii) For each supra fuzzy point  $p$  in  $X$  and each neighborhood  $u$  of  $f(p)$ , then there exist a neighborhood  $v$  of  $p$  such that  $f(v) \subseteq u$ .
- (iii) For any supra fuzzy set  $u$  in  $X$ ,  $f(u) \subseteq f(u)$ .
- (iv) For any supra fuzzy set  $v$  in  $Y$ ,  $(f^{-1}(v)) \subseteq f^{-1}(v)$ .

**Proposition 1.43( [62] Theorem 3.1)**: Let  $f : (X, t^*) \longrightarrow (Y, s^*)$  be supra fuzzy open function , then the following properties hold:

- (i)  $f(u^\circ) \subseteq (f(u))^\circ$ , for each supra fuzzy set  $u$  in  $X$ .
- (ii)  $(f^{-1}(v))^\circ \subseteq f^{-1}(v^\circ)$ , for each supra fuzzy set  $v$  in  $Y$ .

**Proposition 1.44([63] Theorem 1.5 )** : Let  $f : (X, t^*) \longrightarrow (Y, s^*)$  be a function. Then  $f$  is closed if and only if  $f(u) \subseteq f(u)$  for each supra fuzzy set  $u$  in  $X$ .

**Definition 1.45[72]**: Let  $(X, t^*)$  be a supra fuzzy topological space and  $A$  be an ordinary subset of  $X$ . The class  $t^*_A = \{u / A : u \in t^*\}$  determines a supra fuzzy topology on  $A$ . This topology is called the subspace supra fuzzy topology on  $A$ .

**Definition 1.46[106]:** Let  $(X, \tau^*)$  be a supra fuzzy topological space. A subfamily  $B$  of  $\tau^*$  is a base for  $\tau^*$  if and only if each member of  $\tau^*$  can be express as the union of some members of  $B$ .

**Definition 1.47[106]:** Let  $(X, \tau^*)$  be a supra fuzzy topological space. A subfamily  $S$  of  $\tau^*$  is a sub- base for  $\tau^*$  if and only if the family of finite intersection of members of  $S$  forms a base for  $\tau^*$ .

**Definition 1.48[111] :** If  $u_1$  and  $u_2$  are two supra fuzzy subsets of  $X$  and  $Y$  respectively, then the Cartesian product  $u_1 \times u_2$  of two supra fuzzy subsets  $u_1$  and  $u_2$  is a supra fuzzy subsets of  $X \times Y$  defined by  $(u_1 \times u_2)(x, y) = \min(u_1(x), u_2(y))$ , for each pair  $(x, y) \in X \times Y$ .

**Definition 1.49[105] :** Let  $\{X_i, i \in \Lambda\}$ , be any class of sets and let  $X$  denoted the Cartesian product of these sets, i.e.,  $X = \prod_{i \in \Lambda} X_i$ . Note that  $X$  consists of all points  $p = \langle a_i, i \in \Lambda \rangle$ , where  $a_i \in X_i$ . Recall that , for each  $j_0 \in \Lambda$  , we define the projection  $\pi_{j_0}$  from the product set  $X$  to the coordinate space  $X_{j_0}$ . ie  $\pi_{j_0} : X \longrightarrow X_{j_0}$  by  $\pi_{j_0}(\langle a_i : i \in \Lambda \rangle) = a_{j_0}$ ,

These projections are used to define the product supra topology.

**Definition 1.50[106]:** If  $(X_1, \tau_1^*)$  and  $(X_2, \tau_2^*)$  be two supra fuzzy topological space and  $X = X_1 \times X_2$  be the usual product and  $\tau^*$  be the coarsest supra fuzzy topology on  $X$ , then each projection  $\pi_i : X \longrightarrow X_i, i = 1, 2$  is supra fuzzy continuous. The pair  $(X, \tau^*)$  is called the product space of the supra fuzzy topological spaces  $(X_1, \tau_1^*)$  and  $(X_2, \tau_2^*)$ .

**Proposition 1.51( [13] Theorem 3.6 ) :** If  $u$  is a supra fuzzy subset of a supra fuzzy topological space  $(X, \tau_1^*)$  and  $v$  is a supra fuzzy subsets of a supra fuzzy topological space  $(Y, \tau_2^*)$ , then  $u \times v \subseteq u \times v$ .

**Definition 1.52[105]:** Let  $\{X_\alpha\}_{\alpha \in \Lambda}$  be a family of nonempty sets. Let  $X = \prod_{\alpha \in \Lambda} X_\alpha$  be the usual product of  $X_\alpha$ 's and let  $\pi_\alpha$  be the projection from  $X$  into  $X_\alpha$ . Further assume that each  $X_\alpha$  is an supra fuzzy topological spaces with supra fuzzy topology  $t_\alpha^*$ . Now the supra fuzzy topology generated by  $\{\pi_\alpha^{-1}(b_\alpha) : b_\alpha \in t_\alpha^*, \alpha \in \Lambda\}$  as a sub basis, is called the product supra fuzzy topology on  $X$ . Clearly if  $w$  is a basis element in the product, then there exist  $\alpha_1, \alpha_2, \dots, \alpha_n \in \Lambda$  such that  $w(x) = \min\{b_\alpha(x_\alpha) : \alpha = 1, 2, 3, \dots, n\}$ , where  $x = (x_\alpha)_{\alpha \in \Lambda} \in X$ .

**Definition 1.53[111]:** Let  $(X, T)$  be a topological spaces and  $T^*$  be associated supra topology with  $T$ . Then the function  $f: X \rightarrow R$  is semi continuous if and only if  $\{x \in X : f(x) > \alpha\}$  is open for all  $\alpha \in R$ .

**Definition 1.54[60]:** Let  $(X, T)$  be a topological spaces and  $T^*$  be associated supra topology with  $T$ . Then the lower semi continuous topology on  $X$  is  $\omega(T^*) = \{u \in I^X : u^{-1}(\alpha, 1] \in T^*\}$  for each  $\alpha \in I_1$ . It can be shown that  $\omega(T^*)$  is a supra fuzzy topology on  $X$ .

Let  $P$  be the property of a topological space  $(X, T^*)$  and  $FP$  be its supra fuzzy topological analogue. Then  $FP$  is called a 'good extension' of  $P$  " iff the statement  $(X, T^*)$  has  $P$  iff  $(X, \omega(T^*))$  has  $FP$  " holds good for every topological space  $(X, T^*)$ .

**Definition 1.55[3]:** A fuzzy topological space  $(X, t)$  is said to be fuzzy  $T_0$  if and only if

- (a) for all distinct elements  $x, y \in X$ , there exists  $u \in t$  such that  $u(x) = 1, u(y) = 0$  or  $u(x) = 0, u(y) = 1$ ;
- (b) for all distinct elements  $x, y \in X$ , there exists  $u \in t$  such that  $u(x) < u(y)$  or  $u(y) < u(x)$ ;
- (c) for all distinct elements  $x, y \in X, \bar{x}_1 \cap \bar{y}_1 < 1$ .

**Definition 1.56[3]:** A fuzzy topological space  $(X, t)$  is said to be fuzzy  $T_1$  if and only if

- (a) for all distinct elements  $x, y \in X$ , there exist  $u, v \in \tau$  such that  $u(x) = 1, u(y) = 0$  and  $v(x) = 0, v(y) = 1$ ;
- (b) for all distinct elements  $x, y \in X$ , there exist  $u, v \in \tau$  such that  $u(x) > 0, u(y) = 0$  and  $v(x) = 0, v(y) > 0$ ;
- (c) for all distinct elements  $x, y \in X$ , there exist  $u, v \in \tau$  such that  $u(x) > u(y)$  and  $v(y) > v(x)$ .

**Definition 1.57[3]:** A fuzzy topological space  $(X, \tau)$  is said to be fuzzy Hausdorff or  $T_2$  if and only if

- (a) for all distinct elements  $x, y \in X$ , there exist  $u, v \in \tau$  such that  $u(x) = 1 = u(y)$  and  $u \cap v = 0$ ;
- (b) for all pair of distinct fuzzy points  $x_r, y_s \in X$ , there exist  $u, v \in \tau$  such that  $x_r \in u, y_s \in v$  and  $u \cap v = 0$ ;
- (c) for all distinct elements  $x, y \in X$ , there exist  $u, v \in \tau$  such that  $u(x) > 0, v(y) > 0$  and  $u \cap v = 0$ .

**Definition 1.58[3]:** A fuzzy topological space  $(X, \tau)$  is said to be fuzzy regular if and only if for each  $x \in X$  and closed fuzzy set  $u$  with  $u(x) = 0$ , there exists open fuzzy sets  $v, w \in \tau$  such that  $v(x) = 1, u \subseteq w$  and  $v \subseteq 1 - w$ .

**Definition 1.59[3]:** A fuzzy topological space  $(X, \tau)$  is said to be fuzzy normal if and only if for each close fuzzy set  $m$  and open fuzzy set  $u$  with  $m \subseteq u$ , there exists a fuzzy set  $v$  such that  $m \subseteq v^\circ \subseteq \bar{v} \subseteq u$ .

# CHAPTER-2

## $T_0$ Spaces in Supra Fuzzy Topology

### 2.1 Introduction

Four concepts of  $T_0$  supra fuzzy topological spaces are introduced and studied in this chapter. We also establish some relationships among them and study some other properties of these spaces.

**Definition 2.2:** Let  $(X, t)$  be a fuzzy topological space and  $t^*$  be associated supra topology with  $t$  and  $\alpha \in I_1$ . Then

(a)  $(X, t^*)$  is an  $\alpha - T_0(i)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$  or there exists  $v \in t^*$  such that  $v(x) \leq \alpha, v(y) = 1$ .

(b)  $(X, t^*)$  is an  $\alpha - T_0(ii)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $u(x) = 0, u(y) > \alpha$  or there exists  $v \in t^*$  such that  $v(x) > \alpha, v(y) = 0$ .

(c)  $(X, t^*)$  is an  $\alpha - T_0(iii)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $0 \leq u(x) \leq \alpha < u(y) \leq 1$  or there exists  $v \in t^*$  such that  $0 \leq v(y) \leq \alpha < v(x) \leq 1$ .

(d)  $(X, t^*)$  is a  $T_0(iv)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $u(x) \neq u(y)$ .


**Lemma 2.3:** Suppose  $(X, t)$  is a topological space and  $t^*$  is associated supra topology with  $t$  and  $\alpha \in I_1$ . Then the following implications are true:

(a)  $(X, t^*)$  is  $\alpha - T_0(i)$  implies  $(X, t^*)$  is  $\alpha - T_0(iii)$  implies  $(X, t^*)$  is  $T_0(iv)$ .

(b)  $(X, t^*)$  is  $\alpha - T_0(ii)$  implies  $(X, t^*)$  is  $\alpha - T_0(iii)$  implies  $(X, t^*)$  is  $T_0(iv)$ .



Also, these can be shown in a diagram as follows:



**Proof:** Suppose  $(X, t^*)$  is  $\alpha - T_0 (i)$ . We have to prove that  $(X, t^*)$  is  $\alpha - T_0 (iii)$ . Let  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha - T_0 (i)$ , for  $\alpha \in I_1$ , then by definition, there exists  $u \in t^*$  such that  $u(x) = 1$ ,  $u(y) \leq \alpha$  which shows that  $0 \leq u(y) \leq \alpha < u(x) \leq 1$ . Hence by definition (c),  $(X, t^*)$  is  $\alpha - T_0 (iii)$ .

Suppose  $(X, t^*)$  is  $\alpha - T_0 (iii)$ . Then, for  $x, y \in X$  with  $x \neq y$ , there exist  $u \in t^*$  such that  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ , i.e.,  $u(x) \neq u(y)$ , hence by definition,  $(X, t^*)$  is  $\alpha - T_0 (iv)$ .

Let  $(X, t^*)$  is  $\alpha - T_0 (ii)$ . Then, for  $x, y \in X$  with  $x \neq y$ , there exists  $u \in t^*$  such that  $u(x) = 0$  and  $u(y) > \alpha$ , which implies  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ . Hence, by definition,  $(X, t^*)$  is  $\alpha - T_0 (iii)$  and hence  $(X, t^*)$  is  $\alpha - T_0 (iv)$ .

Therefore, the proof is complete.

The non-implications among  $\alpha - T_0 (i)$ ,  $\alpha - T_0 (ii)$ ,  $\alpha - T_0 (iii)$  and  $T_0 (iv)$  are shown in the following examples, i.e., The following examples show that

(a)  $T_0 (iv)$  does not imply  $\alpha - T_0 (iii)$ , so, not imply  $\alpha - T_0 (i)$  and  $\alpha - T_0 (ii)$ .

(b)  $\alpha - T_0 (iii)$  does not imply  $\alpha - T_0 (i)$  and  $\alpha - T_0 (ii)$ .

(c)  $\alpha - T_0 (i)$  does not imply  $\alpha - T_0 (ii)$ .

(d)  $\alpha - T_0$  (ii) does not imply  $\alpha - T_0$  (i).

**Example 2.4:** Let  $X = \{x, y\}$  and  $u \in I^X$  is defined by  $u(x) = 0.4$ ,  $u(y) = 0.7$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . Then for  $\alpha = 0.8$ , we can easily show that  $(X, t^*)$  is  $T_0$ (iv) but  $(X, t^*)$  is not  $\alpha - T_0$ (iii), so, not  $\alpha - T_0$  (i) and  $\alpha - T_0$  (ii).

**Example 2.5:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0.5$ ,  $u(y) = 0.9$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . For  $\alpha = 0.7$ , we have  $0 \leq u(x) \leq 0.7 < u(y) \leq 1$ . Thus according to the definition,  $(X, t^*)$  is  $\alpha - T_0$  (iii) but  $(X, t^*)$  is not  $\alpha - T_0$  (i). Also we can easily show that  $(X, t^*)$  is not  $\alpha - T_0$ (ii).

**Example 2.6:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 1$ ,  $u(y) = 0.5$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . For  $\alpha = 0.7$ , we have  $u(x) = 1$  and  $u(y) \leq \alpha$ . Thus according to the definition  $(X, t^*)$  is  $\alpha - T_0$ (i) but  $(X, t^*)$  is not  $\alpha - T_0$  (ii).

**Example 2.7:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0$ ,  $u(y) = 0.8$ . Let the supra fuzzy supra topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . For  $\alpha = 0.4$ , it can easily show that  $(X, t^*)$  is  $\alpha - T_0$ (ii) but  $(X, t^*)$  is not  $\alpha - T_0$ (i).

This completes the proof.

**Lemma 2.8:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $\alpha, \beta \in I^*$  with  $0 \leq \alpha \leq \beta < 1$ , then

(a)  $(X, t^*)$  is  $\alpha - T_0$  (i) implies  $(X, t^*)$  is  $\beta - T_0$  (i).

(b)  $(X, t^*)$  is  $\beta$ - $T_0$ (ii) implies  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii).

(c)  $(X, t^*)$  is  $0$ - $T_0$ (ii) if and only if  $(X, t^*)$  is  $0$ - $T_0$ (iii).

**Proof:** Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_0$ (i). We have to show that  $(X, t^*)$  is  $\beta$ - $T_0$ (i). Let any two distinct elements  $x, y \in X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_0$ (i), for  $\alpha \in I_1$ , there is  $u \in t^*$  such that  $u(x) = 1$ , and  $u(y) \leq \alpha$ . This implies that  $u(x) = 1$ , and  $u(y) \leq \beta$ , since  $0 \leq \alpha \leq \beta < 1$ . Hence by definition,  $(X, t^*)$  is  $\beta$ - $T_0$ (i).

Suppose that  $(X, t^*)$  is  $\beta$ - $T_0$ (ii). Then, for  $x, y \in X$  with  $x \neq y$ , there exist  $u \in t^*$  such that  $u(x) = 0$  and  $u(y) > \beta$ , which implies  $u(x) = 0$  and  $u(y) > \alpha$ , since  $0 \leq \alpha \leq \beta < 1$ . Hence we have  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii).

**Example 2.9:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 1$ ,  $u(y) = 0.6$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.5$  and  $\beta = 0.8$ ;  $(X, t^*)$  is  $\beta$ - $T_0$ (i) but  $(X, t^*)$  is not  $\alpha$ - $T_0$ (i).

**Example 2.10:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0$ ,  $u(y) = 0.65$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.45$  and  $\beta = 0.75$ ;  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii) but  $(X, t^*)$  is not  $\beta$ - $T_0$ (ii).

In the same way, we can prove that  $(X, t^*)$  is  $0$ - $T_0$ (ii) if and only if  $(X, t^*)$  is  $0$ - $T_0$ (iii).

**Theorem 2.11:** Let  $(X, T)$  be a topological space and  $T^*$  be associated supra topology with  $T$  and  $\alpha \in I_1$ . Suppose that the following statements:

- (1)  $(X, T^*)$  be a  $T_0$ -space.
- (2)  $(X, \omega(T^*))$  be an  $\alpha$ - $T_0$ (i) space.

(3)  $(X, \omega(T^*))$  be an  $\alpha$ - $T_0$ (ii) space.

(4)  $(X, \omega(T^*))$  be an  $\alpha$ - $T_0$ (iii) space.

(5)  $(X, \omega(T^*))$  be a  $T_0$ (iv) space.

Then the following implications are true:

(a)  $(1) \Rightarrow (2) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .

(b)  $(1) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .

**Proof:** Suppose  $(X, T^*)$  is a  $T_0$  – topological space. We have to prove that  $(X, \omega(T^*))$  is  $\alpha$  -  $T_0$ (i) space. Suppose  $x$  and  $y$  are two distinct elements in  $X$ . Since  $(X, T^*)$  is  $T_0$ , there is  $U \in T^*$  such that  $x \in U, y \notin U$ . By the definition of  $lsc$ , we have  $1_U \in \omega(T^*)$  and  $1_U(x) = 1, 1_U(y) = 0$ . Hence we have  $(X, \omega(T^*))$  is  $\alpha$  -  $T_0$ (i) space. Also we have  $(X, \omega(T^*))$  is  $\alpha$  -  $T_0$ (ii) space. Further, it is easy to show that  $(2) \Rightarrow (4), (3) \Rightarrow (4)$  and  $(4) \Rightarrow (5)$ . We therefore prove that  $(4) \Rightarrow (1)$ .

Suppose  $(X, \omega(T^*))$  be a  $T_0$ (iv) space. We have to prove that  $(X, T^*)$  is  $T_0$  – space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, \omega(T^*))$  is  $T_0$ (iv), there is  $u \in \omega(T^*)$  such that  $u(x) < u(y)$  or  $u(x) > u(y)$ . Suppose  $u(x) < u(y)$ . Then for  $r \in I_1$ , such that  $u(x) < r < u(y)$ . We observe that  $x \notin u^{-1}(r, 1], y \in u^{-1}(r, 1]$ , and by definition of  $lsc, u^{-1}(r, 1] \in T^*$ . Hence  $(X, T^*)$  is  $T_0$  – space.

Thus it is seen that  $\alpha$  -  $T_0(p)$  is a good extension of its topological counterpart ( $p = i, ii, iii, iv$ ).

**Theorem 2.12:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $\alpha \in I_1$  and  $I_\alpha(t^*) = \{u^{-1}(\alpha, 1] : u \in t^*\}$ , then

(a)  $(X, t^*)$  is an  $\alpha$ -T<sub>0</sub>(i) implies  $(X, I_\alpha(t^*))$  is T<sub>0</sub>.

(b)  $(X, t^*)$  is an  $\alpha$ -T<sub>0</sub>(ii) implies  $(X, I_\alpha(t^*))$  is T<sub>0</sub>.

(c)  $(X, t^*)$  is an  $\alpha$ -T<sub>0</sub>(iii) if and only if  $(X, I_\alpha(t^*))$  is T<sub>0</sub>.

**Proof:** (a) Let  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  be  $\alpha$ -T<sub>0</sub>(i). Suppose  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(i), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$ . Since  $u^{-1}(\alpha, 1] \in I_\alpha(t^*), y \notin u^{-1}(\alpha, 1]$  and  $x \in u^{-1}(\alpha, 1]$ , We have that  $(X, I_\alpha(t^*))$  is T<sub>0</sub>-space.

(b) Let  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  be  $\alpha$ -T<sub>0</sub>(ii). Suppose  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(ii), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 0, u(y) > \alpha$ . Since  $u^{-1}(\alpha, 1] \in I_\alpha(t^*), y \in u^{-1}(\alpha, 1]$  and  $x \notin u^{-1}(\alpha, 1]$  thus, we have that  $(X, I_\alpha(t^*))$  is T<sub>0</sub>-space.

(c) Finally, suppose that  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(iii). Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(iii), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ . Since  $u^{-1}(\alpha, 1] \in I_\alpha(t^*), x \notin u^{-1}(\alpha, 1]$  and  $y \in u^{-1}(\alpha, 1]$ , so these implies  $(X, I_\alpha(t^*))$  is T<sub>0</sub>-space.

Conversely, suppose  $(X, I_\alpha(t^*))$  be T<sub>0</sub>-space. We shall prove that  $(X, t^*)$  is a  $\alpha$ -T<sub>0</sub>(iii) space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, I_\alpha(t^*))$  is T<sub>0</sub>-space, there exists  $u^{-1}(\alpha, 1] \in I_\alpha(t^*)$  such that  $x \in u^{-1}(\alpha, 1]$  and  $y \notin u^{-1}(\alpha, 1]$ , where  $u \in t^*$ . Then we have  $u(x) > \alpha, u(y) \leq \alpha$ , i.e.,  $0 \leq u(y) \leq \alpha < u(x) \leq 1$ , and hence by definition,  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(iii) space.

**Example 2.13:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0.7, u(y) = 0$ . Suppose the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, 1, \text{Constants}\}$ . Then by definition, for

$\alpha = 0.5$ ,  $(X, t^*)$  is not  $\alpha$ - $T_0$ (i) and  $(X, t^*)$  is not  $\alpha$ - $T_0$ (ii). Now  $I_\alpha(t^*) = \{X, \phi, \{x\}\}$ . Then we see that  $I_\alpha(t^*)$  is a supra topology on  $X$  and  $(X, I_\alpha(t^*))$  is a  $T_0$ -space. This completes the proof.

**Theorem 2.14:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $A \subseteq X$ , and  $t^*_A = \{u/A : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is  $\alpha$ - $T_0$ (i) implies  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (i).
- (b)  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii) implies  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (ii).
- (c)  $(X, t^*)$  is  $\alpha$ - $T_0$ (iii) implies  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (iii).
- (d)  $(X, t^*)$  is  $T_0$ (iv) implies  $(A, t^*_A)$  is  $T_0$ (iv).

**Proof:** (a) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_0$ (i). Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_0$ (i), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$ . For  $A \subseteq X$ , we have  $u/A \in t^*_A$  and  $(u/A)(x) = 1$  and  $(u/A)(y) \leq \alpha$  as  $x, y \in A$ . Hence by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (i).

(b) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii). Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 0, u(y) > \alpha$ . For  $A \subseteq X$ , we have  $u/A \in t^*_A$  and  $(u/A)(x) = 0$  and  $(u/A)(y) > \alpha$  as  $x, y \in A$ . Hence by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (ii).

(c) Suppose that  $(X, t^*)$  is  $\alpha$ - $T_0$ (iii). Then for  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_0$ (iii), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ . For  $A \subseteq X$ , we have  $u/A \in t^*_A$  and  $0 \leq (u/A)(x) \leq \alpha < (u/A)(y) \leq 1$  as  $x, y \in A$ . Hence, by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_0$ (iii).

Similarly, we can prove (d).

**Theorem 2.15:** Suppose that  $(X_i, t^*_i)$ ,  $i \in \Lambda$  be supra fuzzy topological spaces and  $X = \prod_{i \in \Lambda} X_i$  and  $t^*$  be the product supra fuzzy topology on  $X$ , then

- (a)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_0(i)$  if and only if  $(X, t^*)$  is  $\alpha - T_0(i)$ .
- (b)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_0(ii)$  if and only if  $(X, t^*)$  is  $\alpha - T_0(ii)$ .
- (c)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_0(iii)$  if and only if  $(X, t^*)$  is  $\alpha - T_0(iii)$ .
- (d)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $T_0(iv)$  if and only if  $(X, t^*)$  is  $T_0(iv)$ .

**Proof:** (a) Suppose that  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_0(i)$ . Let  $x, y \in X$  with  $x \neq y$ , then  $x_i \neq y_i$ , for some  $i \in \Lambda$ . Since  $(X_i, t^*_i)$  is  $\alpha - T_0(i)$ , for  $\alpha \in I_1$ , there exists  $u_i \in t^*_i$ ,  $i \in \Lambda$  such that  $u_i(x_i) = 1$  and  $u_i(y_i) \leq \alpha$ . But we have  $\pi_i(x) = x_i$ , and  $\pi_i(y) = y_i$ . Then  $u_i(\pi_i(x)) = 1$  and  $u_i(\pi_i(y)) \leq \alpha$ , i.e.,  $(u_i \circ \pi_i)(x) = 1$  and  $(u_i \circ \pi_i)(y) \leq \alpha$ . It follows that there exists  $(u_i \circ \pi_i) \in t^*$  such that  $(u_i \circ \pi_i)(x) = 1$ ,  $(u_i \circ \pi_i)(y) \leq \alpha$ . Hence by definition,  $(X, t^*)$  is  $\alpha - T_0(i)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_0(i)$  space. We have to show that  $(X_i, t^*_i)$ ,  $i \in \Lambda$  is  $\alpha - T_0(i)$ . Let  $a_i$  be a fixed element in  $X_i$  and  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j$  for some  $i \neq j\}$ . Thus  $A_i$  is a subset of  $X$  and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_0(i)$ ,  $(A_i, t^*_{A_i})$  is also  $\alpha - T_0(i)$ . Now we have  $A_i$  is homeomorphic image of  $X_i$ . Thus, we have  $(X_i, t^*_i)$ ,  $i \in \Lambda$  is  $\alpha - T_0(i)$ .

(b) Suppose that  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_0(ii)$ . Let  $x, y \in X$  with  $x \neq y$ , then  $x_i \neq y_i$ , for some  $i \in \Lambda$ . Since  $(X_i, t^*_i)$  is  $\alpha - T_0(ii)$ , for  $\alpha \in I_1$ , there exists  $u_i \in t^*_i$ ,  $i \in \Lambda$  such that  $u_i(x_i) = 0$  and  $u_i(y_i) > \alpha$ . But we have  $\pi_i(x) = x_i$ , and  $\pi_i(y) = y_i$ . Then

$u_i(\pi_i(x))=0$  and  $u_i(\pi_i(y))>\alpha$ , i.e.,  $(u_i \circ \pi_i)(x)=0$  and  $(u_i \circ \pi_i)(y)>\alpha$ . It follows that, there exists  $(u_i \circ \pi_i) \in t^*$  such that  $(u_i \circ \pi_i)(x)=0, (u_i \circ \pi_i)(y)>\alpha$ . Hence, by definition,  $(X, t^*)$  is  $\alpha - T_0(ii)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_0(ii)$  space. We have to show that  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - T_0(ii)$ . Let  $a_i$  be a fixed element in  $X_i$  and  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Thus  $A_i$  is a subset of  $X$  and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_0(ii)$ ,  $(A_i, t^*_{A_i})$  is also  $\alpha - T_0(ii)$ . Then, we have  $A_i$  is homeomorphic image of  $X_i$ . Thus, we have  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - T_0(ii)$

(c) Suppose that  $\forall i \in \Lambda, (X_i, t^*_i)$  is  $\alpha - T_0(iii)$ . Let  $x, y \in X$  with  $x \neq y$ , then  $x_i \neq y_i$ , for some  $i \in \Lambda$ . Since  $(X_i, t^*_i)$  is  $\alpha - T_0(iii)$ , for  $\alpha \in I_1$ , there exists  $u_i \in t^*_i, i \in \Lambda$  such that  $0 \leq (u_i)(x_i) \leq \alpha < (u_i)(y_i) \leq 1$ . But we have  $\pi_i(x) = x_i$ , and  $\pi_i(y) = y_i$ . Then, we have  $0 \leq u_i(\pi_i(x)) \leq \alpha < u_i(\pi_i(y)) \leq 1$ , i.e.,  $0 \leq (u_i \circ \pi_i)(x) \leq \alpha < (u_i \circ \pi_i)(y) \leq 1$ . It follows that, there exists  $(u_i \circ \pi_i) \in t^*$  such that  $0 \leq (u_i \circ \pi_i)(x) \leq \alpha < (u_i \circ \pi_i)(y) \leq 1$ . Hence, by definition,  $(X, t^*)$  is  $\alpha - T_0(iii)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_0(iii)$  space. We have to show that  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - T_0(iii)$ . Let  $a_i$  be a fixed element in  $X_i$  and  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Thus  $A_i$  is a subset of  $X$  and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_0(iii)$ , so  $(A_i, t^*_{A_i})$  is also  $\alpha - T_0(iii)$ . Then, we have  $A_i$  is homeomorphic image of  $X_i$ . Thus, we have  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - T_0(iii)$ .

Similarly, (d) can be proved.



**Theorem 2.16:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be a one-one, onto and open map, then

- (a)  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(i) implies  $(Y, s^*)$  is  $\alpha$ -T<sub>0</sub>(i).
- (b)  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(ii) implies  $(Y, s^*)$  is  $\alpha$ -T<sub>0</sub>(ii).
- (c)  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(iii) implies  $(Y, s^*)$  is  $\alpha$ -T<sub>0</sub>(iii).
- (d)  $(X, t^*)$  is T<sub>0</sub>(iv) implies  $(Y, s^*)$  is T<sub>0</sub>(iv).

**Proof:** (a) Suppose  $(X, t^*)$  be  $\alpha$ -T<sub>0</sub>(i). We have to prove that  $(Y, s^*)$  is  $\alpha$ -T<sub>0</sub>(i). Let  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ . Since  $f$  is onto, there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1$ ,  $f(x_2) = y_2$  and hence  $x_1 \neq x_2$  as  $f$  is one-one. Again, since  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(i), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 1$ ,  $u(y) \leq \alpha$ .

$$\text{Now, } f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$$

$$= 1.$$

$$f(u)(y_2) = \{ \text{Sup } u(x_2) : f(x_2) = y_2 \}$$

$$\leq \alpha.$$

Since  $f$  is open,  $f(u) \in s^*$  as  $u \in t^*$ . We observe that there exists  $f(u) \in s^*$  such that  $f(u)(y_1) = 1$ ,  $f(u)(y_2) \leq \alpha$ . Hence by definition,  $(Y, s^*)$  is  $\alpha$ -T<sub>0</sub>(i).

(b) Suppose  $(X, t^*)$  be  $\alpha$ -T<sub>0</sub>(ii). Then for  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ , there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1$ ,  $f(x_2) = y_2$ , since  $f$  is onto, and hence  $x_1 \neq x_2$  as  $f$  is one-one. Again, since  $(X, t^*)$  is  $\alpha$ -T<sub>0</sub>(ii), for  $\alpha \in I_1$ , there exists  $u \in t^*$  such that  $u(x) = 0$ ,  $u(y) > \alpha$ .

Now,  $f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$

$$= 0.$$

$f(u)(y_2) = \{ \text{Sup } u(x_2) : f(x_2) = y_2 \}$

$$> \alpha.$$

Since  $f$  is open,  $f(u) \in s^*$  as  $u \in t^*$ . We observe that there exists  $f(u) \in s^*$  such that  $f(u)(y_1) = 0$ ,  $f(u)(y_2) > \alpha$ . Hence by definition,  $(Y, s^*)$  is  $\alpha - T_0$  (ii).

Similarly, (c) and (d) can be proved.

**Theorem 2.17:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f : X \longrightarrow Y$  be continuous and one-one map, then

- (a)  $(Y, s^*)$  is  $\alpha - T_0$  (i) implies  $(X, t^*)$  is  $\alpha - T_0$  (i).
- (b)  $(Y, s^*)$  is  $\alpha - T_0$  (ii) implies  $(X, t^*)$  is  $\alpha - T_0$  (ii).
- (c)  $(Y, s^*)$  is  $\alpha - T_0$  (iii) implies  $(X, t^*)$  is  $\alpha - T_0$  (iii).
- (d)  $(Y, s^*)$  is  $T_0$  (iv) implies  $(X, t^*)$  is  $T_0$  (iv).

**Proof:** (a) Suppose  $(Y, s^*)$  be  $\alpha - T_0$  (i). We have to prove that  $(X, t^*)$  is  $\alpha - T_0$  (i). Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , then  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also since  $(Y, s^*)$  is  $\alpha - T_0$  (i), for  $\alpha \in I_1$ , there exists  $u \in s^*$  such that  $u(f(x_1)) = 1$  and  $u(f(x_2)) \leq \alpha$ . This implies that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(u)(x_2) \leq \alpha$ , since  $u \in s^*$  and  $f$  is continuous, then  $f^{-1}(u) \in t^*$ . Thus there is a  $f^{-1}(u) \in t^*$  such that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(u)(x_2) \leq \alpha$ . Hence  $(X, t^*)$  is  $\alpha - T_0$  (i).

(b) Suppose  $(Y, s^*)$  be  $\alpha - T_0$  (ii). Then, for  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , we have,  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also since  $(Y, s^*)$  is  $\alpha - T_0$  (ii), for  $\alpha \in I_1$ , there exists  $u \in s^*$  such

that  $u(f(x_1)) = 0$  and  $u(f(x_2)) > \alpha$ . This implies that  $f^{-1}(u)(x_1) = 0$ ,  $f^{-1}(u)(x_2) > \alpha$ , since  $u \in s^*$  and  $f$  is continuous, then  $f^{-1}(u) \in t^*$ . Thus there is a  $f^{-1}(u) \in t^*$  such that  $f^{-1}(u)(x_1) = 0$ ,  $f^{-1}(u)(x_2) > \alpha$ . Hence  $(X, t^*)$  is  $\alpha$ - $T_0$ (ii).

(c) Suppose  $(Y, s^*)$  be  $\alpha$ - $T_0$ (iii). Then, for  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , we have,  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also, since  $(Y, s^*)$  is  $\alpha$ - $T_0$ (iii), for  $\alpha \in I_1$ , there exists  $u \in s^*$  such that  $0 \leq u(f(x_1)) \leq \alpha < u(f(x_2)) \leq 1$ . This implies that  $0 \leq f^{-1}(u)(x_1) \leq \alpha < f^{-1}(u)(x_2) \leq 1$ , since  $u \in s^*$  and  $f$  is continuous, then  $f^{-1}(u) \in t^*$ . Thus, there is a  $f^{-1}(u) \in t^*$  such that  $0 \leq f^{-1}(u)(x_1) \leq \alpha < f^{-1}(u)(x_2) \leq 1$ . Hence  $(X, t^*)$  is  $\alpha$ - $T_0$ (iii).

Similarly, (d) can be proved.

# CHAPTER 3

## $T_1$ Spaces in Supra Fuzzy Topology

### 3.1 Introduction

In this chapter, we introduce and study four concepts of  $T_1$  supra fuzzy topological spaces. We also establish some relationships among them and study some other properties of these spaces.


**Definition 3.2:** Let  $(X, t)$  be a fuzzy topological space and  $t^*$  be associated supra topology with  $t$  and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is an  $\alpha - T_1(i)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$  and there exists  $v \in t^*$  such that  $v(x) \leq \alpha, v(y) = 1$ .
- (b)  $(X, t^*)$  is an  $\alpha - T_1(ii)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $u(x) = 0, u(y) > \alpha$  and there exists  $v \in t^*$  such that  $v(x) > \alpha, v(y) = 0$ .
- (c)  $(X, t^*)$  is an  $\alpha - T_1(iii)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u \in t^*$  such that  $0 \leq u(y) \leq \alpha < u(x) \leq 1$  and there exists  $v \in t^*$  such that  $0 \leq v(x) \leq \alpha < v(y) \leq 1$ .
- (d)  $(X, t^*)$  is a  $T_1(iv)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u, v \in t^*$  such that  $u(x) < u(y)$  and  $v(x) > v(y)$ .

**Lemma 3.3:** Suppose  $(X, t)$  is a topological space and  $t^*$  is associated supra topology with  $t$  and  $\alpha \in I_1$ . Then the following implications are hold:

- (a)  $(X, t^*)$  is  $\alpha - T_1(i)$  implies  $(X, t^*)$  is  $\alpha - T_1(iii)$  implies  $(X, t^*)$  is  $T_1(iv)$ .
- (b)  $(X, t^*)$  is  $\alpha - T_1(ii)$  implies  $(X, t^*)$  is  $\alpha - T_1(iii)$  implies  $(X, t^*)$  is  $T_1(iv)$ .

One can show these in a diagram given below:



**Proof:** Suppose  $(X, t^*)$  is  $\alpha - T_1(i)$ . We have to prove that  $(X, t^*)$  is  $\alpha - T_1(iii)$ . Let  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha - T_1(i)$ , for  $\alpha \in I_1$ , then by definition, there exist  $u, v \in t^*$  such that  $u(x)=1, u(y) \leq \alpha$  and  $v(y)=1, v(x) \leq \alpha$  which shows that  $0 \leq u(y) \leq \alpha < u(x) \leq 1$  and  $0 \leq v(x) \leq \alpha < v(y) \leq 1$ . Hence by definition (c),  $(X, t^*)$  is  $\alpha - T_1(iii)$ .

Let  $(X, t^*)$  is  $\alpha - T_1(iii)$ . Then for any  $x$  and  $y$  be two distinct elements in  $X$ , there exist  $u, v \in t^*$  such that  $0 \leq u(y) \leq \alpha < u(x) \leq 1$ , i.e.,  $u(y) < u(x)$ . Hence, by definition,  $(X, t^*)$  is  $T_1(iv)$ .

Let  $(X, t^*)$  is  $\alpha - T_1(ii)$ . Then for any  $x$  and  $y$  be two distinct elements in  $X$ , there exist  $u, v \in t^*$  such that  $u(x) = 0, u(y) > \alpha$  and  $v(x) = 0, v(y) > \alpha$ , i.e.,  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ , and  $0 \leq v(y) \leq \alpha < v(x) \leq 1$ . Hence, by definition,  $(X, t^*)$  is  $T_1(iii)$  and  $(X, t^*)$  is  $T_1(iv)$ .

The non-implications among  $\alpha - T_1(i)$ ,  $\alpha - T_1(ii)$ ,  $\alpha - T_1(iii)$  and  $T_1(iv)$  are shown in the following examples: .

**Example 3.4:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0.52, u(y) = 0.95$  and  $v(x) = 0.95, v(y) = 0.52$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.81$ , we have  $0 \leq u(x) \leq 0.81 < u(y) \leq 1$  and  $0 \leq v(y) \leq 0.81 < v(x) \leq 1$ .

Thus according to the definition,  $(X, t^*)$  is  $\alpha - T_1(iii)$  but  $(X, t^*)$  is not  $\alpha - T_1(i)$ .

Also, we can easily show that  $(X, t^*)$  is not  $\alpha - T_1(ii)$ .

**Example 3.5:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 1, u(y) = 0.53$  and  $v(x) = 0.53, v(y) = 1$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.76$ , we have  $u(x) = 1, u(y) \leq \alpha$  and  $v(y) = 1, v(x) \leq \alpha$ . Thus according to the definition  $(X, t^*)$  is  $\alpha$ - $T_1$ (i) but  $(X, t^*)$  is not  $\alpha$ - $T_1$ (ii).

**Example 3.6:** Let  $X = \{x, y\}$  and  $u \in I^X$  be defined by  $u(x) = 0, u(y) = 0.83$  and  $v(x) = 0.83, v(y) = 0$ . Let the supra fuzzy supra topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.45$ , it can easily show that  $(X, t^*)$  is  $\alpha$ - $T_1$ (ii) but  $(X, t^*)$  is not  $\alpha$ - $T_1$ (i).

**Example 3.7:** Let  $X = \{x, y\}$  and  $u \in I^X$  is defined by  $u(x) = 0.35, u(y) = 0.72$  and  $v(x) = 0.72, v(y) = 0.35$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then for  $\alpha = 0.82$ , we can easily show that  $(X, t^*)$  is  $T_1$ (iv) but  $(X, t^*)$  is not  $\alpha$ - $T_1$ (iii) and hence not  $\alpha$ - $T_1$ (i) and  $\alpha$ - $T_1$ (ii).

Thus the proof is complete.

**Theorem 3.8:** Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $\alpha, \beta \in I^*$  with  $0 \leq \alpha \leq \beta < 1$ . Then

- (a)  $(X, t^*)$  is  $\alpha$ - $T_1$ (i) implies  $(X, t^*)$  is  $\beta$ - $T_1$ (i).
- (b)  $(X, t^*)$  is  $\beta$ - $T_1$ (ii) implies  $(X, t^*)$  is  $\alpha$ - $T_1$ (ii).
- (c)  $(X, t^*)$  is  $0$ - $T_1$ (iii) if and only if  $(X, t^*)$  is  $0$ - $T_1$ (iii).

**Proof:** (a) Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_1$ (i). We have to show that  $(X, t^*)$  is  $\beta$ - $T_1$ (i). Let any two distinct elements  $x, y \in X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_1$ (i), for  $\alpha \in I_1$ , there is  $u, v \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$  and  $v(y) = 1, v(x) \leq \alpha$ . This implies that  $u(x) = 1, u(y) \leq \beta$ , and  $v(x) \leq \beta, v(y) = 1$ , since  $0 \leq \alpha \leq \beta < 1$ . Hence by definition,  $(X, t^*)$  is  $\beta$ - $T_1$ (i).

(b) Suppose  $(X, t^*)$  is  $\beta - T_1(ii)$ . We have to show that  $(X, t^*)$  is  $\alpha - T_1(ii)$ . Then for any two distinct elements  $x, y \in X$ , there are  $u, v \in t^*$  such that  $u(x) = 0, u(y) > \beta$  and  $v(y) = 0, v(x) > \beta$ , for  $\beta \in I_1$ . This implies that  $u(x) = 0, u(y) > \alpha$  and  $v(y) = 0, v(x) > \alpha$ , since  $0 \leq \alpha \leq \beta < 1$ . Hence, by definition,  $(X, t^*)$  is  $\alpha - T_1(ii)$ .

The proof of (c) is trivial.

**Example 3.9:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 1, u(y) = 0.62$  and  $v(x) = 0.62, v(y) = 1$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.51$  and  $\beta = 0.84$ ; Thus  $(X, t^*)$  is  $\beta - T_1(i)$  but  $(X, t^*)$  is not  $\alpha - T_1(i)$ .

**Example 3.10:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 0, u(y) = 0.65$  and  $v(x) = 0.66, v(y) = 0$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.41$  and  $\beta = 0.84$ ;  $(X, t^*)$  is  $\alpha - T_1(ii)$  but  $(X, t^*)$  is not  $\beta - T_1(ii)$ .

**Theorem 3.11:** Let  $(X, T)$  be a topological space and  $T^*$  be associated supra topology with  $T$  and  $\alpha \in I_1$ . Suppose that the following statements:

- (1)  $(X, T^*)$  be a  $T_1$  - space.
- (2)  $(X, \omega(T^*))$  be an  $\alpha - T_1(i)$  space.
- (3)  $(X, \omega(T^*))$  be an  $\alpha - T_1(ii)$  space.
- (4)  $(X, \omega(T^*))$  be an  $\alpha - T_1(iii)$  space .
- (5)  $(X, \omega(T^*))$  be a  $T_1(iv)$  space .

Then the followings are true:

- (a)  $(1) \Rightarrow (2) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .
- (b)  $(1) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .

**Proof:** Suppose  $(X, T^*)$  is a  $T_1$  – topological space. We have to prove that  $(X, \omega(T^*))$  is  $\alpha - T_1(i)$  space. Suppose  $x$  and  $y$  are two distinct elements in  $X$ . Since  $(X, T^*)$  is  $T_1$  -spaces, there are  $U, V \in T^*$  such that  $x \in U, y \notin U$  and  $x \notin V, y \in V$ . By the definition of  $lsc$ , we have  $1_U, 1_V \in \omega(T^*)$  and  $1_U(x) = 1, 1_U(y) = 0$  and  $1_V(x) = 0, 1_V(y) = 1$ . Hence we have  $(X, \omega(T^*))$  is  $\alpha - T_1(i)$  -space. Also we have  $(X, \omega(T^*))$  is  $\alpha - T_1(ii)$  space. Further, it is easy to show that  $(2) \Rightarrow (4), (3) \Rightarrow (4)$  and  $(4) \Rightarrow (5)$ . We therefore prove that  $(5) \Rightarrow (1)$ .

Suppose  $(X, \omega(T^*))$  be a  $T_1(iv)$  - space. We have to prove that  $(X, T^*)$  is  $T_1$  – space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, \omega(T^*))$  is  $T_1(iv)$ , there are  $u, v \in \omega(T^*)$  such that  $u(x) < u(y)$  and  $v(x) > v(y)$ . Suppose that  $r, s \in I_1$  such that  $u(x) < r < u(y)$  and  $v(x) > s > v(y)$ . We observe that  $x \notin u^{-1}(r, 1], y \in u^{-1}(r, 1]$  and  $x \in v^{-1}(s, 1], y \notin v^{-1}(s, 1]$ , and by definition of  $lsc$ ,  $u^{-1}(r, 1]$  and  $v^{-1}(s, 1] \in T^*$ . Therefore  $(X, T^*)$  is  $T_1$  – space.

Thus it is seen that  $\alpha - T_1(p)$  is a good extension of its topological counter part ( $p = i, ii, iii, iv$ ).

**Theorem 3.12:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $\alpha \in I_1$  and  $I_\alpha(t^*) = \{u^{-1}(\alpha, 1] : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is an  $\alpha - T_1(i)$ -space implies  $(X, I_\alpha(t^*))$  is  $T_1$  -space.
- (b)  $(X, t^*)$  is an  $\alpha - T_1(ii)$ -space implies  $(X, I_\alpha(t^*))$  is  $T_1$  -space.
- (c)  $(X, t^*)$  is an  $\alpha - T_1(iii)$ -space if and only if  $(X, I_\alpha(t^*))$  is  $T_1$  -space.

**Proof:** (a) Let  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  be  $\alpha - T_1(i)$ . Suppose  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha - T_1(i)$ , for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$  and  $v(x) \leq \alpha, v(y) = 1$ . Since  $u^{-1}(\alpha, 1]$ ,



$v^{-1}(\alpha, 1] \in I_\alpha(t^*)$ ,  $y \notin u^{-1}(\alpha, 1]$ ,  $x \in u^{-1}(\alpha, 1]$  and  $x \notin v^{-1}(\alpha, 1]$ ,  $y \in v^{-1}(\alpha, 1]$ .

We have that  $(X, I_\alpha(t^*))$  is  $T_1$ -space.

(b) Suppose that  $(X, t^*)$  be  $\alpha - T_1(i)$ . Then for any  $x$  and  $y$  be two distinct elements in  $X$ , there exist  $u, v \in t^*$  such that  $u(x) = 0$ ,  $u(y) > \alpha$  and  $v(x) > \alpha$ ,  $v(y) = 0$ , for  $\alpha \in I_1$ . Since  $u^{-1}(\alpha, 1]$ ,  $v^{-1}(\alpha, 1] \in I_\alpha(t^*)$ ,  $y \in u^{-1}(\alpha, 1]$ ,  $x \notin u^{-1}(\alpha, 1]$  and  $x \in v^{-1}(\alpha, 1]$ ,  $y \notin v^{-1}(\alpha, 1]$ . We have that  $(X, I_\alpha(t^*))$  is  $T_1$ -space.

(c) Suppose that  $(X, t^*)$  is  $\alpha - T_1(iii)$ -space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, t^*)$  is  $\alpha - T_1(iii)$ , for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $0 \leq u(x) \leq \alpha < u(y) \leq 1$  and  $0 \leq v(y) \leq \alpha < v(x) \leq 1$ . Since  $u^{-1}(\alpha, 1]$ ,  $v^{-1}(\alpha, 1] \in I_\alpha(t^*)$ ,  $x \notin u^{-1}(\alpha, 1]$ ,  $y \in u^{-1}(\alpha, 1]$  and  $x \in v^{-1}(\alpha, 1]$ ,  $y \notin v^{-1}(\alpha, 1]$ , so these implies  $(X, I_\alpha(t^*))$  is  $T_1$ -space.

Conversely, suppose  $(X, I_\alpha(t^*))$  be  $T_1$ -space. We shall prove that  $(X, t^*)$  is a  $\alpha - T_1(iii)$  space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, I_\alpha(t^*))$  is  $T_1$ -space, there exist  $u^{-1}(\alpha, 1]$ ,  $v^{-1}(\alpha, 1] \in I_\alpha(t^*)$  such that  $x \in u^{-1}(\alpha, 1]$  and  $y \notin u^{-1}(\alpha, 1]$ ;  $x \notin v^{-1}(\alpha, 1]$  and  $y \in v^{-1}(\alpha, 1]$ , where  $u, v \in t^*$ . Then we have  $u(x) > \alpha$ ,  $u(y) \leq \alpha$  and  $v(x) \leq \alpha$ ,  $v(y) > \alpha$ , i.e.,  $0 \leq u(y) \leq \alpha < u(x) \leq 1$  and  $0 \leq v(x) \leq \alpha < v(y) \leq 1$ , and hence by definition,  $(X, t^*)$  is  $\alpha - T_1(iii)$ -space.

**Example 3.13:** Let  $X = \{x, y\}$  and  $u, v, w \in I^X$  be defined by,  $u(x) = 1$ ,  $u(y) = 0.63$ ,  $v(x) = 0.43$ ,  $v(y) = 0.83$ ,  $w(x) = 0.23$ ,  $w(y) = 0.29$ . Suppose the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, w, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.51$ ,  $(X, t^*)$  is not  $\alpha - T_1(i)$  and  $(X, t^*)$  is not  $\alpha - T_1(ii)$ . Now  $I_\alpha(t^*) = \{X, \phi, \{x\}, \{y\}\}$ . Then we see that  $I_\alpha(t^*)$  is a supra topology on  $X$  and  $(X, I_\alpha(t^*))$  is a  $T_1$ -space. This completes the proof.

**Theorem 3.14:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $A \subseteq X$ , and  $t^*_A = \{u/A : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is  $\alpha - T_1$  (i) implies  $(A, t^*_A)$  is  $\alpha - T_1$  (i).
- (b)  $(X, t^*)$  is  $\alpha - T_1$  (ii) implies  $(A, t^*_A)$  is  $\alpha - T_1$  (ii).
- (c)  $(X, t^*)$  is  $\alpha - T_1$  (iii) implies  $(A, t^*_A)$  is  $\alpha - T_1$  (iii).
- (d)  $(X, t^*)$  is  $T_1$  (iv) implies  $(A, t^*_A)$  is  $T_1$  (iv).

**Proof:** (a) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha - T_1$ (i) space. Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha - T_1$  (i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$  and  $v(x) \leq \alpha, v(y) = 1$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) = 1$  and  $(u/A)(y) \leq \alpha$  and  $(v/A)(x) \leq \alpha, (v/A)(y) = 1$ , as  $x, y \in A$ . Hence, by definition,  $(A, t^*_A)$  is  $\alpha - T_1$ (i).

(b) Suppose that  $(X, t^*)$  is  $\alpha - T_1$ (ii)-space. Then for  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$  and there exist  $u, v \in t^*$  such that  $u(x) = 0, u(y) > \alpha$  and  $v(x) > \alpha, v(y) = 0$ , for  $\alpha \in I_1$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) = 0$  and  $(u/A)(y) > \alpha$  and  $(v/A)(x) > \alpha, (v/A)(y) = 0$ , as  $x, y \in A$ . Hence, by definition,  $(A, t^*_A)$  is  $\alpha - T_1$ (ii).

(c) Suppose that  $(X, t^*)$  is  $\alpha - T_1$ (iii)-space. Then for  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$  and there exist  $u, v \in t^*$  such that  $0 \leq u(y) \leq \alpha < u(x) \leq 1$  and  $0 \leq v(x) \leq \alpha < v(y) \leq 1$ , for  $\alpha \in I_1$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $0 \leq (u/A)(y) \leq \alpha < (u/A)(x) \leq 1$  and  $0 \leq (v/A)(x) \leq \alpha < (v/A)(y) \leq 1$ , as  $x, y \in A$ . Hence, by definition,  $(A, t^*_A)$  is  $\alpha - T_1$ (iii).

Similarly, (d) can be proved.

**Theorem 3.15:** Suppose that  $(X_i, t^*_i)$ ,  $i \in \Lambda$  be supra fuzzy topological spaces and  $X = \prod_{i \in \Lambda} X_i$  and  $t^*$  be the product supra fuzzy topology on  $X$ , then

- (a)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_1(i)$  if and only if  $(X, t^*)$  is  $\alpha - T_1(i)$ .
- (b)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_1(ii)$  if and only if  $(X, t^*)$  is  $\alpha - T_1(ii)$ .
- (c)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_1(iii)$  if and only if  $(X, t^*)$  is  $\alpha - T_1(iii)$ .
- (d)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $T_1(iv)$  if and only if  $(X, t^*)$  is  $T_1(iv)$ .

**Proof:** (a) Suppose that  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_1(i)$ . Let  $x, y \in X$  with  $x \neq y$ , then  $x_i \neq y_i$ , for some  $i \in \Lambda$ . Since  $(X_i, t^*_i)$  is  $\alpha - T_1(i)$ , for  $\alpha \in I_1$ , there exists  $u_i \in t^*_i$ ,  $i \in \Lambda$  such that  $u_i(x_i) = 1$ ,  $u_i(y_i) \leq \alpha$  and  $v_i(y_i) = 1$ ,  $v_i(x_i) \leq \alpha$ . But we have  $\pi_i(x) = x_i$ , and  $\pi_i(y) = y_i$ . Then  $u_i(\pi_i(x)) = 1$ ,  $u_i(\pi_i(y)) \leq \alpha$  and  $v_i(\pi_i(y)) = 1$ ,  $v_i(\pi_i(x)) \leq \alpha$ , i.e.,  $(u_i \circ \pi_i)(x) = 1$ ,  $(u_i \circ \pi_i)(y) \leq \alpha$  and  $(v_i \circ \pi_i)(y) = 1$ ,  $(v_i \circ \pi_i)(x) \leq \alpha$ . It follows that there exist  $(u_i \circ \pi_i), (v_i \circ \pi_i) \in t^*$  such that  $(u_i \circ \pi_i)(x) = 1$ ,  $(u_i \circ \pi_i)(y) \leq \alpha$  and  $(v_i \circ \pi_i)(y) = 1$ ,  $(v_i \circ \pi_i)(x) \leq \alpha$ . Hence by definition,  $(X, t^*)$  is  $\alpha - T_1(i)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_1(i)$  space. We have to show that  $(X_i, t^*_i)$ ,  $i \in \Lambda$  is  $\alpha - T_1(i)$ . Let  $a_i$  be a fixed element in  $X_i$  and  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j$  for some  $i \neq j\}$ . Thus  $A_i$  is a subset of  $X$  and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_1(i)$ ,  $(A_i, t^*_{A_i})$  is also  $\alpha - T_1(i)$ . Now we have  $A_i$  is homeomorphic image of  $X_i$ . Thus, we have  $(X_i, t^*_i)$ ,  $i \in \Lambda$  is  $\alpha - T_1(i)$ .

(b) Suppose that  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha - T_1(ii)$ . Then, for  $x, y \in X$  with  $x \neq y$ , we have  $x_i \neq y_i$ , for some  $i \in \Lambda$ , there exists  $u_i \in t^*_i$ ,  $i \in \Lambda$  such that  $u_i(x_i) = 0$ ,  $u_i(y_i) > \alpha$  and  $v_i(y_i) = 0$ ,  $v_i(x_i) > \alpha$ , for  $\alpha \in I_1$ . But, we have  $\pi_i(x) = x_i$ , and  $\pi_i(y) = y_i$ . Then

$u_i(\pi_i(x)) = 0$ ,  $u_i(\pi_i(y)) > \alpha$  and  $v_i(\pi_i(y)) = 0$ ,  $v_i(\pi_i(x)) > \alpha$ , i.e.,  $(u_i \circ \pi_i)(x) = 0$ ,  $(u_i \circ \pi_i)(y) > \alpha$  and  $(v_i \circ \pi_i)(y) = 0$ ,  $(v_i \circ \pi_i)(x) > \alpha$ . It follows that there exist  $(u_i \circ \pi_i), (v_i \circ \pi_i) \in t^*$  such that  $(u_i \circ \pi_i)(x) = 0$ ,  $(u_i \circ \pi_i)(y) > \alpha$  and  $(v_i \circ \pi_i)(y) = 0$ ,  $(v_i \circ \pi_i)(x) > \alpha$ . Hence, by definition,  $(X, t^*)$  is  $\alpha - T_1$  (ii).

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_1$  (ii) space. Let  $a_i$  be a fixed element in  $X_i$  and  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Thus  $A_i$  is a subset of  $X$  and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_1$  (ii),  $(A_i, t^*_{A_i})$  is also  $\alpha - T_1$  (i). Thus we have  $A_i$  is homeomorphic image of  $X_i$ . Hence  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - T_1$  (i).

Similarly, (c) and (d) can be proved.

**Theorem 3.16:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be a one-one, onto and open map, then

- (a)  $(X, t^*)$  is  $\alpha - T_1$  (i) implies  $(Y, s^*)$  is  $\alpha - T_1$  (i).
- (b)  $(X, t^*)$  is  $\alpha - T_1$  (ii) implies  $(Y, s^*)$  is  $\alpha - T_1$  (ii).
- (c)  $(X, t^*)$  is  $\alpha - T_1$  (iii) implies  $(Y, s^*)$  is  $\alpha - T_1$  (iii).
- (d)  $(X, t^*)$  is  $T_1$  (iv) implies  $(Y, s^*)$  is  $T_1$  (iv).

**Proof:** (a) Let  $(X, t^*)$  be  $\alpha - T_1$  (i). We have to prove that  $(Y, s^*)$  is  $\alpha - T_1$  (i). Let  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ . Since  $f$  is onto, there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1$ ,  $f(x_2) = y_2$  and also  $x_1 \neq x_2$  as  $f$  is one-one. Again since  $(X, t^*)$  is  $\alpha - T_1$  (i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x_1) = 1$ ,  $u(x_2) \leq \alpha$  and  $v(x_1) \leq \alpha$ ,  $v(x_2) = 1$ .

$$\text{Now, } f(u)(y_1) = \{\text{Sup } u(x_1) : f(x_1) = y_1\}$$

$$= 1.$$

$$f(u)(y_2) = \{ \text{Sup } u(x_2) : f(x_2) = y_2 \}$$

$$\leq \alpha,$$

$$\text{and } f(v)(y_1) = \{ \text{Sup } v(x_1) : f(x_1) = y_1 \}$$

$$\leq \alpha.$$

$$f(v)(y_2) = \{ \text{Sup } v(x_2) : f(x_2) = y_2 \}$$

$$= 1.$$

Since  $f$  is open then  $f(u) \in s^*$  as  $u \in t^*$ . We observe that there exist  $f(u), f(v) \in s^*$  such that  $f(u)(y_1) = 1$ ,  $f(u)(y_2) \leq \alpha$  and  $f(v)(y_1) \leq \alpha$ ,  $f(v)(y_2) = 1$ . Hence by definition,  $(Y, s^*)$  is  $\alpha - T_1(i)$

(b) Let  $(X, t^*)$  be  $\alpha - T_1(ii)$ . Then for  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ , there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1$ ,  $f(x_2) = y_2$ , since  $f$  is onto, and also  $x_1 \neq x_2$  as  $f$  is one-one. Again, since  $(X, t^*)$  is  $\alpha - T_1(i)$ , for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x_1) = 0$ ,  $u(x_2) > \alpha$  and  $v(x_1) > \alpha$ ,  $v(x_2) = 0$ .

$$\text{Now, } f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$$

$$= 0.$$

$$f(u)(y_2) = \{ \text{Sup } u(x_2) : f(x_2) = y_2 \}$$

$$> \alpha,$$

$$\text{and } f(v)(y_1) = \{ \text{Sup } v(x_1) : f(x_1) = y_1 \}$$

$$> \alpha.$$

$$f(v)(y_2) = \{ \text{Sup } v(x_2) : f(x_2) = y_2 \}$$

$$= 0.$$

Since  $f$  is open, then  $f(u) \in s^*$  as  $u \in t^*$ . We observe that there exist  $f(u), f(v) \in s^*$  such

that  $f(u)(y_1) = 0$ ,  $f(u)(y_2) > \alpha$  and  $f(v)(y_1) > \alpha$ ,  $f(v)(y_2) = 0$ . Hence by definition,  $(Y, s^*)$  is  $\alpha - T_1(ii)$ .

Similarly, (c) and (d) can be proved.

**Theorem 3.17:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f : X \longrightarrow Y$  be continuous and one-one map, then

- (a)  $(Y, s^*)$  is  $\alpha - T_1(i)$  implies  $(X, t^*)$  is  $\alpha - T_1(i)$ .
- (b)  $(Y, s^*)$  is  $\alpha - T_1(ii)$  implies  $(X, t^*)$  is  $\alpha - T_1(ii)$ .
- (c)  $(Y, s^*)$  is  $\alpha - T_1(iii)$  implies  $(X, t^*)$  is  $\alpha - T_1(iii)$ .
- (d)  $(Y, s^*)$  is  $T_1(iv)$  implies  $(X, t^*)$  is  $T_1(iv)$ .

**Proof:** (a) Let  $(Y, s^*)$  be  $\alpha - T_1(i)$ . We have to prove that  $(X, t^*)$  is  $\alpha - T_1(i)$ . Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , then  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also since  $(Y, s^*)$  is  $\alpha - T_1(i)$ , for  $\alpha \in I_1$ , there exist  $u, v \in s^*$  such that  $u(f(x_1)) = 1$ ,  $u(f(x_2)) \leq \alpha$  and  $v(f(x_1)) \leq \alpha$ ,  $v(f(x_2)) = 1$ . This implies that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(u)(x_2) \leq \alpha$  and  $f^{-1}(v)(x_1) \leq \alpha$ ,  $f^{-1}(v)(x_2) = 1$ , since  $u, v \in s^*$  and  $f$  is continuous, then  $f^{-1}(u), f^{-1}(v) \in t^*$ . Thus there are  $f^{-1}(u), f^{-1}(v) \in t^*$  such that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(u)(x_2) \leq \alpha$  and  $f^{-1}(v)(x_1) \leq \alpha$ ,  $f^{-1}(v)(x_2) = 1$ . Hence  $(X, t^*)$  is  $\alpha - T_1(i)$ .

(b) Let  $(Y, s^*)$  be  $\alpha - T_1(ii)$ . Then for  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , we have  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also we have, for  $\alpha \in I_1$ , there exist  $u, v \in s^*$  such that  $u(f(x_1)) = 0$ ,  $u(f(x_2)) > \alpha$  and  $v(f(x_1)) > \alpha$ ,  $v(f(x_2)) = 0$ . This implies that  $f^{-1}(u)(x_1) = 0$ ,  $f^{-1}(u)(x_2) > \alpha$  and  $f^{-1}(v)(x_1) > \alpha$ ,  $f^{-1}(v)(x_2) = 0$ , since  $u, v \in s^*$  and  $f$  is continuous, then  $f^{-1}(u), f^{-1}(v) \in t^*$ . Thus there are  $f^{-1}(u), f^{-1}(v) \in t^*$  such that  $f^{-1}(u)(x_1) = 0$ ,  $f^{-1}(u)(x_2) > \alpha$  and  $f^{-1}(v)(x_1) > \alpha$ ,  $f^{-1}(v)(x_2) = 0$ . Hence  $(X, t^*)$  is  $\alpha - T_1(ii)$ .

Similarly, (c) and (d) can be proved.

# CHAPTER 4

## $T_2$ Spaces in Supra Fuzzy Topology

### 4.1 Introduction

Four concepts of  $T_2$  supra fuzzy topological spaces are introduced and studied in this chapter. We also establish some relationships among them and study some other properties of these spaces.


**Definition 4.2:** Let  $(X, t)$  be a fuzzy topological space and  $t^*$  be associated supra topology with  $t$  and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is an  $T_2(i)$  space if and only if for all distinct elements  $x, y \in X$ , there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v = 0$ .
- (b)  $(X, t^*)$  is an  $\alpha - T_2(ii)$  space if and only if for all distinct elements  $x, y \in X$ , there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ .
- (c)  $(X, t^*)$  is an  $\alpha - T_2(iii)$  space if and only if for all distinct elements  $x, y \in X$ , there exists  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v = 0$ .
- (d)  $(X, t^*)$  is an  $\alpha - T_2(iv)$  space if and only if for all distinct elements  $x, y \in X$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ .

**Lemma 4.3:** Suppose  $(X, t)$  is a topological space and  $t^*$  is associated supra topology with  $t$  and  $\alpha \in I_1$ . Then the following implications are hold:

- (a)  $(X, t^*)$  is  $T_2(i)$  implies  $(X, t^*)$  is  $\alpha - T_2(ii)$  implies  $(X, t^*)$  is  $\alpha - T_2(iv)$ .
- (b)  $(X, t^*)$  is  $\alpha - T_2(i)$  implies  $(X, t^*)$  is  $\alpha - T_2(iii)$  implies  $(X, t^*)$  is  $\alpha - T_2(iv)$ .

Also, these can be shown in a diagram as follows:



**Proof:** (a) Suppose  $(X, t^*)$  is  $T_2 (i)$ . We have to prove that  $(X, t^*)$  is  $\alpha - T_2 (ii)$ . Let  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $T_2 (i)$ , for  $\alpha \in I_1$ , then by definition, there exists  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v = 0$ , which shows that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ . Hence by definition (b),  $(X, t^*)$  is  $\alpha - T_2(ii)$ . Also, we see that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . Hence  $(X, t^*)$  is  $\alpha - T_2(iv)$ .

(b) Suppose  $(X, t^*)$  is  $T_2 (i)$ . Then for any  $x$  and  $y$  be two distinct elements in  $X$ , there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v = 0$ , for  $\alpha \in I_1$  which shows that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v = 0$ . Hence by definition (c),  $(X, t^*)$  is  $\alpha - T_2(iii)$  and hence  $(X, t^*)$  is  $\alpha - T_2(iv)$ .

The following examples show the non-implications among  $T_2 (i), \alpha - T_2 (ii), \alpha - T_2 (iii)$  spaces, i.e.,

- (1)  $\alpha - T_2 (iv)$  does not imply  $\alpha - T_2 (ii)$ , so not imply  $T_2 (i)$ .
- (2)  $\alpha - T_2 (iv)$  does not imply  $\alpha - T_2 (iii)$ , (3)  $\alpha - T_2 (ii)$  does not imply  $\alpha - T_2 (iii)$ ,
- (4)  $\alpha - T_2 (iii)$  does not imply  $\alpha - T_2 (ii)$ , (5)  $\alpha - T_2 (iii)$  does not imply  $T_2 (i)$ ,
- (6)  $\alpha - T_2 (ii)$  does not imply  $T_2 (i)$ .

**Example 4.4:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 0.95, u(y) = 0.43$  and  $v(x) = 0.31, v(y) = 0.95$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.75$ , we have  $u(x) = 0.95 > \alpha, v(y) = 0.95 > \alpha$  and  $u \cap v \leq \alpha$ . Thus



according to the definition,  $(X, t^*)$  is  $\alpha - T_2$ (iv) but  $(X, t^*)$  is not  $\alpha - T_2$ (ii) and  $(X, t^*)$  is not  $\alpha - T_2$ (iii). So,  $(X, t^*)$  is not  $T_2$ (i).

**Example 4.5:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 1, u(y) = 0.43$  and  $v(x) = 0.43, v(y) = 1$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.53$ , we have,  $u(x) = 1, v(y) = 1$ , and  $u \cap v \leq \alpha$ . Thus according to the definition,  $(X, t^*)$  is  $\alpha - T_2$ (ii) but  $(X, t^*)$  is not  $\alpha - T_2$ (iii) and  $(X, t^*)$  is not  $T_2$ (i).

**Example 4.6:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 0.85, u(y) = 0$  and  $v(x) = 0, v(y) = 0.85$ . Let the supra fuzzy supra topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . For  $\alpha = 0.53$ , it can easily show that  $(X, t^*)$  is  $\alpha - T_2$ (iii) but  $(X, t^*)$  is not  $\alpha - T_1$ (ii) and  $(X, t^*)$  is not  $T_2$ (i).

**Theorem 4.7:** Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $\alpha, \beta \in t^*$  with  $0 \leq \alpha \leq \beta < 1$ , then

- (a)  $(X, t^*)$  is  $\alpha - T_2$  (ii) implies  $(X, t^*)$  is  $\beta - T_2$  (ii).
- (b)  $(X, t^*)$  is  $\beta - T_2$  (iii) implies  $(X, t^*)$  is  $\alpha - T_2$ (iii).
- (c)  $(X, t^*)$  is  $0 - T_2$ (iii) if and only if  $(X, t^*)$  is  $0 - T_2$ (iv).

**Proof:** (a) Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha - T_2$ (ii).

We have to show that  $(X, t^*)$  is  $\beta - T_2$ (ii). Let any two distinct elements  $x, y \in X$ . Since

$(X, t^*)$  is  $\alpha - T_2$ (ii), for  $\alpha \in I_1$ , there is  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ .

This implies that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \beta$  as  $0 \leq \alpha \leq \beta < 1$ . Hence by definition,

$(X, t^*)$  is  $\beta - T_2$  (ii).

(b) Suppose that  $(X, t^*)$  is  $\beta - T_2$ (iii). We have to show that  $(X, t^*)$  is  $\alpha - T_2$ (iii). Then, any

two distinct elements  $x, y \in X$ , there is  $u, v \in t^*$  such that  $u(x) > \beta, v(y) > \beta$  and

$u \cap v = 0$ , for  $\alpha \in I_1$ . This implies that  $u(x) > \alpha$ ,  $v(y) > \alpha$  and  $u \cap v = 0$  as  $0 \leq \alpha \leq \beta < 1$ .

Hence by definition,  $(X, t^*)$  is  $\alpha - T_2$  (iii).

(c) The proof is trivial.

**Example 4.8:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 1$ ,  $u(y) = 0.62$  and  $v(x) = 0.82$ ,  $v(y) = 1$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.31$  and  $\beta = 0.84$ ;  $(X, t^*)$  is  $\beta - T_2$  (ii) but  $(X, t^*)$  is not  $\alpha - T_2$  (ii).

**Example 4.9:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  be defined by  $u(x) = 0$ ,  $u(y) = 0.72$  and  $v(x) = 0.88$ ,  $v(y) = 0$ . Let the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then, by definition, for  $\alpha = 0.33$  and  $\beta = 0.85$ ;  $(X, t^*)$  is  $\alpha - T_2$  (iii) but  $(X, t^*)$  is not  $\beta - T_2$  (iii).

**Theorem 4.10:** Let  $(X, T)$  be a topological space and  $T^*$  be associated supra topology with  $T$  and  $\alpha \in I_1$ . Suppose that the following statements:

- (1)  $(X, T^*)$  be a  $T_2$  - space.
- (2)  $(X, \omega(T^*))$  be a  $T_2$  (i) space.
- (3)  $(X, \omega(T^*))$  be an  $\alpha - T_2$  (ii) space.
- (4)  $(X, \omega(T^*))$  be an  $\alpha - T_2$  (iii) space.
- (5)  $(X, \omega(T^*))$  be an  $\alpha - T_2$  (iv) space.

Then the followings are true:

- (a)  $(1) \Rightarrow (2) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .
- (b)  $(1) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ .

**Proof:** Suppose  $(X, T^*)$  is a  $T_2$  - topological space. We have to prove that  $(X, \omega(T^*))$  is  $T_2$ (i) space. Suppose  $x$  and  $y$  are two distinct elements in  $X$ . Since  $(X, T^*)$  is  $T_2$  -spaces,

there are  $U, V \in T^*$  such that  $x \in U, y \in V$  and  $U \cap V = \phi$ . By the definition of  $lsc$ , we have  $1_U, 1_V \in \omega(T^*)$  and  $1_U(x) = 1, 1_V(y) = 1$  and  $1_U \cap 1_V = 0$ . If  $1_U \cap 1_V \neq 0$ , then there exists  $z \in X$  such that  $(1_U \cap 1_V)(z) \neq 0$  implies  $1_U(z) \neq 0, 1_V(z) \neq 0$  implies  $z \in U, z \in V$  implies  $z \in U \cap V \Rightarrow U \cap V \neq \phi$ , a contradiction. So that  $1_U \cap 1_V = 0$ , and consequently  $(X, \omega(T^*))$  is  $T_2(i)$ . Also, we see that  $(X, \omega(T^*))$  is  $\alpha - T_2(ii)$ .

Further, it is easy to show that  $(2) \Rightarrow (4), (3) \Rightarrow (4)$  and  $(4) \Rightarrow (5)$ . We therefore prove that  $(5) \Rightarrow (1)$ .

Now, suppose  $(X, \omega(T^*))$  be a  $T_2(iv)$  - space. We have to prove that  $(X, T^*)$  is  $T_2$  - space.

Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, \omega(T^*))$  is  $T_2(iv)$ , for  $\alpha \in I_1$ , there are  $u, v \in \omega(T^*)$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . We have that  $u^{-1}(\alpha, 1]$  and  $v^{-1}(\alpha, 1] \in T^*$ ,  $\alpha \in I_1$  and  $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ . Moreover  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ . For if  $z \in u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1]$ , then  $z \in u^{-1}(\alpha, 1], z \in v^{-1}(\alpha, 1]$  implies  $u(z) > \alpha$  and  $v(z) > \alpha$  implies  $(u \cap v)(z) > \alpha$ , a contradiction as  $(u \cap v)(z) \leq \alpha$ . Therefore  $(X, T^*)$  is  $T_2$  - space.

Thus it is seen that  $\alpha - T_2(p)$  is a good extension of its topological counter part ( $p = i, ii, iii, iv$ ).

**Theorem 4.11:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $\alpha \in I_1$  and  $I_\alpha(t^*) = \{u^{-1}(\alpha, 1] : u \in t^*\}$ . Then

- (a)  $(X, t^*)$  is an  $\alpha - T_2(ii)$ -space implies  $(X, I_\alpha(t^*))$  is  $T_2$  -space.
- (b)  $(X, t^*)$  is an  $\alpha - T_2(iii)$ -space implies  $(X, I_\alpha(t^*))$  is  $T_2$  -space.
- (c)  $(X, t^*)$  is an  $\alpha - T_2(iv)$ -space if and only if  $(X, I_\alpha(t^*))$  is  $T_2$  -space

**Proof:** (a) Let  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  be  $\alpha - T_2(ii)$ . Suppose  $x$  and  $y$  be any two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha - T_2(ii)$ , for  $\alpha \in I_1$ , there exist

$u, v \in t^*$  such that  $u(x) = 1, v(y) = 1$  and  $u \cap v \leq \alpha$ . But for every  $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t^*)$  and  $y \in v^{-1}(\alpha, 1], x \in u^{-1}(\alpha, 1]$  and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$  as  $u \cap v \leq \alpha$ . We have that  $(X, I_\alpha(t^*))$  is  $T_2$ -space.

(b) Let  $(X, t^*)$  be  $\alpha$ - $T_2$ (iii). Then for any  $x$  and  $y$  be two distinct elements in  $X$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v = 0$ , for  $\alpha \in I_1$ . But for every  $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t^*)$  and  $y \in v^{-1}(\alpha, 1], x \in u^{-1}(\alpha, 1]$  and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$  as  $u \cap v = 0$ . We have that  $(X, I_\alpha(t^*))$  is  $T_2$ -space.

(c) Suppose  $(X, t^*)$  is  $\alpha$ - $T_2$ (iv)-space. We have to prove that  $(X, I_\alpha(t^*))$  is  $T_2$ -space. Suppose  $x$  and  $y$  are two distinct elements in  $X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_2$ (iv)-space, for  $\alpha \in I_1$ , there are  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . But for every  $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t^*)$ . So we have  $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$  and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ , as  $u \cap v \leq \alpha$ . Hence it is clear that  $(X, I_\alpha(t^*))$  is  $T_2$ -space.

Conversely, suppose that  $(X, I_\alpha(t^*))$  is  $T_2$ -space. Let  $x, y \in X$  with  $x \neq y$ . Since  $(X, I_\alpha(t^*))$  is  $T_2$ -space, there are  $U, V \in I_\alpha(t^*)$ , such that  $x \in U, y \in V$  and  $U \cap V = \phi$ . Again since  $U, V \in I_\alpha(t^*)$ , so we get  $u, v \in t^*$  such that  $U = u^{-1}(\alpha, 1], V = v^{-1}(\alpha, 1]$ . This implies that  $u(x) > \alpha, v(y) > \alpha$  and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi \Rightarrow (u \cap v)^{-1}(\alpha, 1] = \phi$ , i.e.,  $u \cap v \leq \alpha$ . So, we see that  $(X, t^*)$  is  $\alpha$ - $T_2$ (iv).

This completes the proof.

**Example 4.12:** Suppose that  $X = \{x, y\}$  and  $u, v \in t^*$  defined by  $u(x) = 0.82, u(y) = 0.21, v(x) = 0.31$  and  $v(y) = 0.74$ . Suppose the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.53, (X, t^*)$  is not  $\alpha$ - $T_2$ (ii). Now

$I_\alpha(t^*) = \{X, \phi, \{x\}, \{y\}\}$ . Then we can easily show that  $I_\alpha(t^*)$  is a topology on  $X$  and  $(X, I_\alpha(t^*))$  is  $T_2$  space.

Similarly, we can easily prove that  $(X, t^*)$  is  $\alpha$ - $T_2$ (iii)-space implies  $(X, I_\alpha(t^*))$  is  $T_2$ -space.

**Theorem 4.13:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $A \subseteq X$ , and  $t^*_A = \{u/A : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is  $T_2$  (i) implies  $(A, t^*_A)$  is  $T_2$  (i).
- (b)  $(X, t^*)$  is  $\alpha$ - $T_2$  (ii) implies  $(A, t^*_A)$  is  $\alpha$ - $T_2$  (ii).
- (c)  $(X, t^*)$  is  $\alpha$ - $T_2$  (iii) implies  $(A, t^*_A)$  is  $\alpha$ - $T_2$  (iii).
- (d)  $(X, t^*)$  is  $\alpha$ - $T_2$  (iv) implies  $(A, t^*_A)$  is  $\alpha$ - $T_2$  (iv).

**Proof:** (a) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $T_2$ (i) space. Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $T_2$  (i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v = 0$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) = 1 = (v/A)(y)$  and  $u/A \cap v/A = 0$ . Hence by definition,  $(A, t^*_A)$  is  $T_2$ (i).

(b) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_2$ (ii)-space. Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_2$  (ii), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) = 1 = (v/A)(y)$  and  $u/A \cap v/A \leq \alpha$ . Hence by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_2$ (ii).

(c) Suppose that  $(X, t^*)$  is  $\alpha$ - $T_2$ (iii)-space. Then for  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . and for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$

and  $u \cap v = 0$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) > \alpha$ ,  $(v/A)(y) > \alpha$  and  $u/A \cap v/A = 0$ . Hence by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_2$ (iii).

(d) Suppose that  $(X, t^*)$  be a supra fuzzy topological space and  $(X, t^*)$  is  $\alpha$ - $T_2$ (iv)-space. Let  $x, y \in A$  with  $x \neq y$ , so that  $x, y \in X$ , as  $A \subseteq X$ . Since  $(X, t^*)$  is  $\alpha$ - $T_2$ (iii), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha$ ,  $v(y) > \alpha$  and  $u \cap v \leq \alpha$ . For  $A \subseteq X$ , we have  $u/A, v/A \in t^*_A$  and  $(u/A)(x) > \alpha$ ,  $(v/A)(y) > \alpha$  and  $u/A \cap v/A \leq \alpha$ . Hence by definition,  $(A, t^*_A)$  is  $\alpha$ - $T_2$ (iv).

**Theorem 4.14:** Suppose that  $(X_i, t^*_i)$ ,  $i \in \Lambda$  be supra fuzzy topological spaces and  $X = \prod_{i \in \Lambda} X_i$  and  $t^*$  be the product supra fuzzy topology on  $X$ , then

- (a)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $T_2$ (i) if and only if  $(X, t^*)$  is  $T_2$ (i).
- (b)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $T_2$ (ii) if and only if  $(X, t^*)$  is  $\alpha$ - $T_2$ (ii).
- (c)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $T_2$ (iii) if and only if  $(X, t^*)$  is  $\alpha$ - $T_2$ (iii).
- (d)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $T_2$ (iv) if and only if  $(X, t^*)$  is  $\alpha$ - $T_2$ (iv).

**Proof:** (b) Suppose  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  be an  $\alpha$ - $T_2$ (ii). Let  $x, y$  be two distinct points in  $X = \prod_{i \in \Lambda} X_i$ , then there exist an  $x_i \neq y_i$  in  $X_i$ . Thus, for  $\alpha \in I_1$ , there exist  $u_i, v_i \in t^*_i$  such that  $u_i(x_i) = 1 = v_i(y_i)$  and  $u_i \cap v_i \leq \alpha$ . But we have  $\pi_i(x) = x_i$ ,  $\pi_i(y) = y_i$ , then  $u_i(\pi_i(x)) = 1 = v_i(\pi_i(y))$  and  $(u_i \cap v_i) \circ \pi_i \leq \alpha$ . Hence  $(u_i \circ \pi_i)(x) = 1 = (v_i \circ \pi_i)(y)$  and  $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) \leq \alpha$ . Put  $u = u_i \circ \pi_i$ ,  $v = v_i \circ \pi_i$ , then  $u, v \in t^*$  with  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ . Hence by definition,  $(X, t^*)$  is  $\alpha$ - $T_2$ (ii).

Conversely, suppose that  $(X, t^*)$  is  $\alpha$ - $T_2$ (ii). For some  $i \in \Lambda$ , let  $a_i$  be a fixed element in  $X_i$ . Suppose that  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Then  $A_i$  is a subsets of  $X$  and therefore  $(A_i, t^*_{A_i})$  is a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha$ - $T_2$ (ii) space.

Then, we have also  $(A_i, t^*_{A_i})$  is  $\alpha - T_2(ii)$  space. Furthermore,  $A_i$  is homeomorphic image of  $X_i$ . Hence, by definition, we have  $(X_i, t^*_i)$  is  $\alpha - T_2(ii)$  space.

(c) Suppose  $\forall i \in \Lambda, (X_i, t^*_i)$  be an  $\alpha - T_2(iii)$ . Let  $x, y$  be two distinct points in  $X = \prod_{i \in \Lambda} X_i$ , then there exist an  $x_i \neq y_i$  in  $X_i$ . Since  $(X_i, t^*_i)$  is an  $\alpha - T_2(iii)$ , for  $\alpha \in I_1$ , there exist  $u_i, v_i \in t^*_i$  such that  $u_i(x_i) > \alpha, v_i(y_i) > \alpha$  and  $u_i \cap v_i = 0$ . But we have  $\pi_i(x) = x_i, \pi_i(y) = y_i$ , then  $u_i(\pi_i(x)) > \alpha, v_i(\pi_i(y)) > \alpha$  and  $(u_i \cap v_i) \circ \pi_i = 0$ . Hence  $(u_i \circ \pi_i)(x) > \alpha, (v_i \circ \pi_i)(y) > \alpha$  and  $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) = 0$ . Put  $u = u_i \circ \pi_i, v = v_i \circ \pi_i$ , then  $u, v \in t^*$  with  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v = 0$ . Hence, we have  $(X, t^*)$  is  $\alpha - T_2(iii)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_2(iii)$ . For some  $i \in \Lambda$ , let  $a_i$  be a fixed element in  $X_i$ . Suppose that  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Then  $A_i$  is a subsets of  $X$  and therefore  $(A_i, t^*_{A_i})$  is a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_2(iii)$  space. Then, we have also  $(A_i, t^*_{A_i})$  is  $\alpha - T_2(iii)$  space. Furthermore,  $A_i$  is homeomorphic image of  $X_i$ . Hence by definition, we have  $(X_i, t^*_i)$  is  $\alpha - T_2(iii)$  space.

(d) Suppose  $\forall i \in \Lambda, (X_i, t^*_i)$  be an  $\alpha - T_2(iv)$ . Let  $x, y$  be two distinct points in  $X = \prod_{i \in \Lambda} X_i$ , then there exist an  $x_i \neq y_i$  in  $X_i$ . Since  $(X_i, t^*_i)$  is an  $\alpha - T_2(iv)$ , for  $\alpha \in I_1, \exists u_i, v_i \in t^*_i$  such that  $u_i(x_i) > \alpha, v_i(y_i) > \alpha$  and  $u_i \cap v_i \leq \alpha$ . But we have  $\pi_i(x) = x_i, \pi_i(y) = y_i$ , then  $u_i(\pi_i(x)) > \alpha, v_i(\pi_i(y)) > \alpha$  and  $(u_i \cap v_i) \circ \pi_i \leq \alpha$ . Hence  $(u_i \circ \pi_i)(x) > \alpha, (v_i \circ \pi_i)(y) > \alpha$  and  $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) \leq \alpha$ . Put  $u = u_i \circ \pi_i, v = v_i \circ \pi_i$ , then  $u, v \in t^*$  with  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . Hence we have  $(X, t^*)$  is  $\alpha - T_2(iv)$ .

Conversely, suppose that  $(X, t^*)$  is  $\alpha - T_2$ (iv). For some  $i \in \Lambda$ , let  $a_i$  be a fixed element in  $X_i$ . Suppose that  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$ . Then  $A_i$  is a subsets of  $X$  and therefore  $(A_i, t^*_{A_i})$  is a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - T_2$ (iv) space. Then, we have also  $(A_i, t^*_{A_i})$  is  $\alpha - T_2$ (iv) space. Furthermore,  $A_i$  is homeomorphic image of  $X_i$ . Hence by definition, we have  $(X_i, t^*_{X_i})$  is  $\alpha - T_2$ (iv) space.

Similarly, (a) can be proved.

**Theorem 4.15:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be a one-one, onto and open map, then

(a)  $(X, t^*)$  is  $T_2$ (i) implies  $(Y, s^*)$  is  $T_2$ (i).

(b)  $(X, t^*)$  is  $\alpha - T_2$ (ii) implies  $(Y, s^*)$  is  $\alpha - T_2$ (ii).

(c)  $(X, t^*)$  is  $\alpha - T_2$ (iii) implies  $(Y, s^*)$  is  $\alpha - T_2$ (iii).

(d)  $(X, t^*)$  is  $\alpha - T_2$ (iv) implies  $(Y, s^*)$  is  $\alpha - T_2$ (iv).

**Proof:** (b) Let  $(X, t^*)$  be  $\alpha - T_2$ (ii). We have to prove that  $(Y, s^*)$  is  $\alpha - T_2$ (ii). Let  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ . Since  $f$  is onto, there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1, f(x_2) = y_2$  and also  $x_1 \neq x_2$  as  $f$  is one-one. Again since  $(X, t^*)$  is  $\alpha - T_2$ (i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x_1) = 1 = v(x_1)$  and  $u \cap v \leq \alpha$ .

$$\text{Now, } f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$$

$$= 1$$

$$f(v)(y_2) = \{ \text{Sup } v(x_2) : f(x_2) = y_2 \}$$

$$= 1$$

$$\text{and } f(u \cap v)(y_1) = \{ \text{Sup } (u \cap v)(x_1) : f(x_1) = y_1 \}$$

$$f(u \cap v)(y_2) = \{ \text{Sup } (u \cap v)(x_2) : f(x_2) = y_2 \}$$

Hence  $f(u \cap v) \leq \alpha \Rightarrow f(u) \cap f(v) \leq \alpha$ .



Since  $f$  is open then  $f(u), f(v) \in s^*$  as  $u, v \in t^*$ . We observe that there exist  $f(u), f(v) \in s^*$  such that  $f(u)(y_1) = 1, f(v)(y_2) = 1$  and  $f(u) \cap f(v) \leq \alpha$ . Hence by definition,  $(Y, s^*)$  is  $\alpha - T_2(ii)$ .

(c) Let  $(X, t^*)$  be  $\alpha - T_2(iii)$ . Then for  $y_1, y_2 \in Y$  with  $y_1 \neq y_2$ , there exist  $x_1, x_2 \in X$  with  $f(x_1) = y_1, f(x_2) = y_2$ , since  $f$  is onto, and also  $x_1 \neq x_2$  as  $f$  is one-one. Again, since  $(X, t^*)$  is  $\alpha - T_2(iii)$ , for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x_1) > \alpha, v(x_1) > \alpha$  and  $u \cap v = 0$ .

$$\text{Now, } f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$$

$$> \alpha$$

$$f(v)(y_2) = \{ \text{Sup } v(x_2) : f(x_2) = y_2 \}$$

$$> \alpha$$

$$\text{and } f(u \cap v)(y_1) = \{ \text{Sup } (u \cap v)(x_1) : f(x_1) = y_1 \}$$

$$f(u \cap v)(y_2) = \{ \text{Sup } (u \cap v)(x_2) : f(x_2) = y_2 \}$$

$$\text{Hence } f(u \cap v) = 0 \Rightarrow f(u) \cap f(v) = 0.$$

Since  $f$  is open then  $f(u), f(v) \in s^*$  as  $u, v \in t^*$ . We observe that there exist  $f(u), f(v) \in s^*$  such that  $f(u)(y_1) > \alpha, f(v)(y_2) > \alpha$  and  $f(u) \cap f(v) = 0$ . Hence by definition,  $(Y, s^*)$  is  $\alpha - T_2(iii)$ .

Similarly, (a) and (d) can be proved.

**Theorem 4.16:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f : X \longrightarrow Y$  be continuous and one-one map, then

(a)  $(Y, s^*)$  is  $T_2(i)$  implies  $(X, t^*)$  is  $T_2(i)$ .

(b)  $(Y, s^*)$  is  $\alpha - T_2(ii)$  implies  $(X, t^*)$  is  $\alpha - T_2(ii)$ .

(c)  $(Y, s^*)$  is  $\alpha - T_2(iii)$  implies  $(X, t^*)$  is  $\alpha - T_2(iii)$ .

(d)  $(Y, s^*)$  is  $\alpha - T_2(iv)$  implies  $(X, t^*)$  is  $\alpha - T_2(iv)$ .

**Proof: (b)** Let  $(Y, s^*)$  be  $\alpha - T_2(ii)$ . Let  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ , then  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also since  $(Y, s^*)$  is  $\alpha - T_2(ii)$ , for  $\alpha \in I_1$ , there exist  $u, v \in s^*$  such that  $u(f(x_1)) = 1 = v(f(x_2))$  and  $u \cap v \leq \alpha$ . This implies that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(v)(x_2) = 1$  and  $f^{-1}(u \cap v) \leq \alpha$ , i.e.,  $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$ , since  $u, v \in s$  and  $f$  is continuous then  $f^{-1}(u), f^{-1}(v) \in t^*$ . Now it is clear that there exist  $f^{-1}(u), f^{-1}(v) \in t^*$  such that  $f^{-1}(u)(x_1) = 1$ ,  $f^{-1}(v)(x_2) = 1$  and  $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$ . Hence  $(X, t^*)$  is  $\alpha - T_2(ii)$ .

(c) Let  $(Y, s^*)$  be  $\alpha - T_2(iii)$ . Then for  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ ,  $f(x_1) \neq f(x_2)$  in  $Y$ , since  $f$  is one-one. Also since  $(Y, s^*)$  is  $\alpha - T_2(iii)$ , for  $\alpha \in I_1$ , there exist  $u, v \in s^*$  such that  $u(f(x_1)) > \alpha$ ,  $v(f(x_2)) > \alpha$  and  $u \cap v = 0$ . This implies that  $f^{-1}(u)(x_1) > \alpha$ ,  $f^{-1}(v)(x_2) > \alpha$  and  $f^{-1}(u \cap v) = 0$ , i.e.,  $f^{-1}(u) \cap f^{-1}(v) = 0$ , since  $u, v \in s$  and  $f$  is continuous then  $f^{-1}(u), f^{-1}(v) \in t^*$ . Thus there exist  $f^{-1}(u), f^{-1}(v) \in t^*$  such that  $f^{-1}(u)(x_1) > \alpha$ ,  $f^{-1}(v)(x_2) > \alpha$  and  $f^{-1}(u) \cap f^{-1}(v) = 0$ . Hence  $(X, t^*)$  is  $\alpha - T_2(iii)$ .

Similarly, (a) and (d) can be proved.

# CHAPTER 5

## Regular Spaces in Supra Fuzzy Topology

### 5.1 Introduction

We introduce and study supra fuzzy regular spaces and we establish some relationships among them in this chapter. We also study some other properties of these concepts and obtain their several features.


**Definition 5.2:** Let  $(X, t)$  be a fuzzy topological space and  $t^*$  be associated supra topology with  $t$  and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is an  $\alpha$  - SFR (i) space if and only if for all  $w \in (t^*)^c$  with  $w(x) < 1$ ,  $\forall x \in X$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ .
- (b)  $(X, t^*)$  is an  $\alpha$  -SFR(ii) space if and only if for all  $w \in (t^*)^c$  with  $w(x) < 1, \forall x \in X$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ .
- (c)  $(X, t^*)$  is an  $\alpha$  -SFR(iii) space if and only if for all  $w \in (t^*)^c$  with  $w(x) = 0, \forall x \in X$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ .
- (d)  $(X, t^*)$  is an  $\alpha$  -SFR(iv) space if and only if for all  $w \in (t^*)^c$  with  $w(x) = 0, \forall x \in X$ , there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}, u \cap v \leq \alpha$ .

**Theorem 5.3:** Let  $(X, t)$  be a fuzzy topological space and  $t^*$  be associated supra topology with  $t$ . Then the following implications are true:

- (a)  $(X, t^*)$  is  $\alpha$  - SFR(i) implies  $(X, t^*)$  is  $\alpha$  - SFR(ii) implies  $(X, t^*)$  is  $\alpha$  - SFR(iv).
- (b)  $(X, t^*)$  is  $\alpha$  - SFR(i) implies  $(X, t^*)$  is  $\alpha$  - SFR(iii) implies  $(X, t^*)$  is  $\alpha$  - SFR(iv).

Also, these can be shown in a diagram as follows:



**Proof:** First, suppose that  $(X, t^*)$  is  $\alpha$  - SFR (i). We have to prove that  $(X, t^*)$  is  $\alpha$  - SFR(ii). Let  $w \in (t^*)^c$  with  $w(x) < 1, x \in X$ . Since  $(X, t^*)$  is  $\alpha$  -SFR(i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . Now, we see that  $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . Hence by definition  $(X, t^*)$  is  $\alpha$  -SFR (ii).

Suppose that  $(X, t^*)$  is  $\alpha$  - SFR (ii). Let  $w \in (t^*)^c$  with  $w(x) < 1, x \in X$ , then for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) > 1, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . Now, we see that  $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . Hence by definition  $(X, t^*)$  is  $\alpha$  -SFR (iv).

In the same way, we can prove that

$$(X, t^*) \text{ is } \alpha\text{-SFR(i)} \Rightarrow (X, t^*) \text{ is } \alpha\text{-SFR(iii)} .$$

$$(X, t^*) \text{ is } \alpha\text{-SFR(iii)} \Rightarrow (X, t^*) \text{ is } \alpha\text{-SFR(iv)} .$$

Now, we give some examples to show the non implication among  $\alpha$ - SFR (i),  $\alpha$  - SFR (ii),  $\alpha$ - SFR (iii) and  $\alpha$  - SFR (iv).

**Example 5.4:** Let  $X = \{ x, y \}$  and  $u, v \in I^X$  are defined by  $u(x) = 0.9, u(y) = 0, v(x) = 0.5$  and  $v(y) = 1$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{constants}\}$ . Then for  $w = 1 - u$  and  $\alpha = 0.7$ , we see that  $(X, t^*)$  is  $\alpha$  -SFR(ii) but  $(X, t^*)$  is not  $\alpha$ -SFR(i) .

**Example 5.5:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 0.2, u(y) = 0.3, v(x) = 0.3, v(y) = 0.2$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then for  $w = 1 - u$  and  $\alpha = 0.5$ , we see that  $(X, t^*)$  is  $\alpha$ -SFR(iii) and  $(X, t^*)$  is  $\alpha$ -SFR(iv), but  $(X, t^*)$  is not  $\alpha$ -SFR(ii) as they do not exist any  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ .

**Example 5.6:** Let  $X = \{x, y\}$  and  $u, v, w \in I^X$  are defined by  $u(x) = 0.9, u(y) = 0, v(x) = 0.5, v(y) = 1, w(x) = 1, w(y) = 0$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, w, 1, \text{Constants}\}$ . Then for  $\alpha = 0.6$  and  $p = 1 - w$ , it is seen that  $(X, t^*)$  is  $\alpha$ -SFR(iv) but  $(X, t^*)$  is not  $\alpha$ -SFR(iii).

This completes the proof.

**Theorem 5.7:** If  $\alpha, \beta \in t^*$  with  $0 \leq \alpha \leq \beta < 1$ , then

(a)  $(X, t^*)$  is  $\alpha$ -SFR(i) implies  $(X, t^*)$  is  $\beta$ -SFR(i).

(b)  $(X, t^*)$  is  $\alpha$ -SFR(iii) implies  $(X, t^*)$  is  $\beta$ -SFR(iii).

**Proof:** Suppose that  $(X, t^*)$  is  $\alpha$ -SFR (i). We have to prove that  $(X, t^*)$  is  $\beta$ -SFR (i). Let  $w \in (t^*)^c$  and  $x \in X$  with  $w(x) < 1$ . Since  $(X, t^*)$  is  $\alpha$ -SFR(i), for  $\alpha \in I_1$ , there exist  $u, v \in t^*$  such that  $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . Since  $\alpha \leq \beta$ , then  $u \cap v \leq \beta$ . So it is observed that  $(X, t^*)$  is  $\beta$ -SFR (i).

**Example 5.8:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 1, u(y) = 0, v(x) = 0.7, v(y) = 1$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{constants}\}$ . Then for  $w = 1 - u, \alpha = 0.75, \beta = 0.6$ . We see that  $(X, t^*)$  is  $\beta$ -SFR(i) but  $(X, t^*)$  is not  $\alpha$ -SFR(i).

In the same way, we can prove that

$(X, t^*)$  is  $\alpha$ -SFR(iii) implies  $(X, t^*)$  is  $\beta$ -SFR(iii).

**Theorem 5.9:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $I_\alpha(t^*) = \{ u^{-1}(\alpha, 1] : u \in t^* \}$ , then  $(X, t^*)$  is 0-SFR (i) implies  $(X, I_0(t^*))$  is Regular.

**Proof:** Suppose  $(X, t^*)$  be a 0-SFR (i). We have to prove that  $(X, I_0(t^*))$  is Regular. Let  $V$  be a closed set in  $I_0(t^*)$  and  $x \in X$  such that  $x \notin V$ . Then  $V^c \in I_0(t^*)$  and  $x \in V^c$ . So, by the definition of  $I_0(t^*)$ , there exists an  $u \in t^*$  such that  $V^c = u^{-1}(0, 1]$ , i.e.,  $u(x) > 0$ . Since  $u \in t^*$ , then  $u^c$  is closed supra fuzzy set in  $t^*$  and  $u^c(x) < 1$ . Since  $(X, t^*)$  is 0-SFR(i), there exist  $v, w \in t^*$  such that  $v(x)=1, w \geq 1_{(u^c)^{-1}\{1\}}, v \cap w = 0$ .

(a) Since  $v, w \in t^*$  then  $v^{-1}(0, 1], w^{-1}(0, 1] \in I_0(t^*)$  and  $x \in v^{-1}(0, 1]$

(b) Since  $w \geq 1_{(u^c)^{-1}\{1\}}$  then  $w^{-1}(0, 1] \supseteq (1_{(u^c)^{-1}\{1\}})^{-1}(0, 1]$ .

(c) And  $v \cap w = 0$ , mean  $(v \cap w)^{-1}(0, 1] = v^{-1}(0, 1] \cap w^{-1}(0, 1] = \phi$ .

Now, we have

$$\begin{aligned} (1_{(u^c)^{-1}\{1\}})^{-1}(0, 1] &= \{ x : 1_{(u^c)^{-1}\{1\}}(x) \in (0, 1] \} \\ &= \{ x : 1_{(u^c)^{-1}\{1\}}(x) = 1 \} \\ &= \{ x : x \in (u^c)^{-1}\{1\} \} \\ &= \{ x : u^c(x) = 1 \} \\ &= \{ x : u(x) = 0 \} \\ &= \{ x : x \notin V^c \} \\ &= \{ x : x \in V \} \\ &= V. \end{aligned}$$

Put  $W = v^{-1}(0, 1]$  and  $W^* = w^{-1}(0, 1]$ , then  $x \in W, W^* \supseteq V$  and  $W \cap W^* = \phi$ . Hence it is clear that  $(X, I_0(t^*))$  is Regular.

**Theorem 5.10:** Let  $(X, \tau^*)$  be a supra fuzzy topological space  $A \subseteq X$ , and  $\tau^*_A = \{u/A : u \in \tau^*\}$ , then  $1_{((u/A)^c)^{-1}\{1\}}(x) = (1_{(u^c)^{-1}\{1\}}/A)(x)$ .

**Proof:** Let  $w$  be a closed supra fuzzy set in  $\tau^*_A$ , i.e.,  $w \in \tau^*_A^c$ , then  $u/A = w^c$ , where  $u \in \tau^*$ .

Now, we have

$$\begin{aligned} 1_{((u/A)^c)^{-1}\{1\}}(x) &= \begin{cases} 0 & \text{if } x \notin ((u/A)^c)^{-1}\{1\} \\ 1 & \text{if } x \in ((u/A)^c)^{-1}\{1\} \end{cases} \\ &= \begin{cases} 0 & \text{if } x \notin \{y : (u/A)^c(y) = 1\} \\ 1 & \text{if } x \in \{y : (u/A)^c(y) = 1\} \end{cases} \\ &= \begin{cases} 0 & \text{if } (u/A)^c(x) < 1 \\ 1 & \text{if } (u/A)^c(x) = 1 \end{cases} \\ &= \begin{cases} 0 & \text{if } w(x) < 1 \\ 1 & \text{if } w(x) = 1 \end{cases} \end{aligned}$$

$$\begin{aligned} \text{Again, } 1_{(u^c)^{-1}\{1\}}(x) &= \begin{cases} 0 & \text{if } x \notin (u^c)^{-1}\{1\} \\ 1 & \text{if } x \in (u^c)^{-1}\{1\} \end{cases} \\ &= \begin{cases} 0 & \text{if } x \notin \{y : u^c(y) = 1\} \\ 1 & \text{if } x \in \{y : u^c(y) = 1\} \end{cases} \\ &= \begin{cases} 0 & \text{if } u^c(x) < 1 \\ 1 & \text{if } u^c(x) = 1 \end{cases} \end{aligned}$$

$$\begin{aligned} \text{Now, } (1_{(u^c)^{-1}\{1\}}/A)(x) &= \begin{cases} 0 & \text{if } (u^c/A)(x) < 1 \\ 1 & \text{if } (u^c/A)(x) = 1 \end{cases} \\ &= \begin{cases} 0 & \text{if } (u/A)^c(x) < 1 \\ 1 & \text{if } (u/A)^c(x) = 1 \end{cases} \end{aligned}$$

$$= \begin{cases} 0 & \text{if } w(x) < 1 \\ 1 & \text{if } w(x) = 1 \end{cases}$$

Hence it is clear that  $1_{((u/A)^c)^{-1}\{1\}}(x) = (1_{(u^c)^{-1}\{1\}} / A)(x)$ .

**Theorem 5.11:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $A \subseteq X$  and  $t^*_A = \{ u/A : u \in t^* \}$ , then

(a)  $(X, t^*)$  is  $\alpha$ -SFR (i) implies  $(A, t^*_A)$  is  $\alpha$ -SFR (i).

(b)  $(X, t^*)$  is  $\alpha$ -SFR (ii) implies  $(A, t^*_A)$  is  $\alpha$ -SFR (ii).

(c)  $(X, t^*)$  is  $\alpha$ -SFR (iii) implies  $(A, t^*_A)$  is  $\alpha$ -SFR (iii).

(d)  $(X, t^*)$  is  $\alpha$ -SFR(iv) implies  $(A, t^*_A)$  is  $\alpha$ -SFR(iv).

**Proof:** Let  $(X, t^*)$  be  $\alpha$ -SFR (i). We have to prove that  $(A, t^*_A)$  is  $\alpha$ -SFR (i). Let  $w$  be a closed fuzzy set in  $t^*_A$ , and  $x^* \in A$  such that  $w(x^*) < 1$ . This implies that  $w^c \in t^*_A$  and  $w^c(x^*) > 0$ . So there exists an  $u \in t^*$  such that  $u/A = w^c$  and clearly  $u^c$  is closed in  $t^*$  and  $u^c(x^*) = (u/A)^c(x^*) = w(x^*) < 1$ , i.e.,  $u^c(x^*) < 1$ . Since  $(X, t^*)$  is  $\alpha$ -SFR (i), for  $\alpha \in I_1$ , there exist  $v, v^* \in t^*$  such that  $v(x^*) = 1, v^* \geq 1_{(u^c)^{-1}\{1\}}$  and  $v \cap v^* \leq \alpha$ . Since  $v, v^* \in t^*$ , then  $v/A, v^*/A \in t^*_A$  and  $v/A(x^*) = 1, v^*/A \geq (1_{(u^c)^{-1}\{1\}}/A)$  and  $v/A \cap v^*/A = (v \cap v^*)/A \leq \alpha$ .

But  $1_{(u^c)^{-1}\{1\}}/A = 1_{((u/A)^c)^{-1}\{1\}} = 1_{w^{-1}\{1\}}$ , then  $v^*/A \geq 1_{w^{-1}\{1\}}$ . Hence it is clear that  $(A, t^*_A)$  is  $\alpha$ -SFR (i).

The proofs of (b), (c) and (d) are similar.

**Theorem 5.12:** Let  $(X, T)$  be a topological space and  $T^*$  be associated supra topology with  $T$ . Consider the following statements:

1)  $(X, T^*)$  is a Regular space.



- 2)  $(X, \omega(T^*))$  is  $\alpha$ -SFR (i).
- 3)  $(X, \omega(T^*))$  is  $\alpha$ -SFR (ii).
- 4)  $(X, \omega(T^*))$  is  $\alpha$ -SFR (iii).
- 5)  $(X, \omega(T^*))$  is  $\alpha$ -SFR (iv).

Then the following are true:

- (a) (1) implies (2) implies (3) implies (5) implies (1),
- (b) (1) implies (2) implies (4) implies (5) implies (1).

**Proof:** First, suppose that  $(X, T^*)$  be regular space. We shall prove that  $(X, \omega(T^*))$  is  $\alpha$ -SFR (i). Let  $w$  be a closed supra fuzzy set in  $\omega(T^*)$  and  $x \in X$  such that  $w(x) < 1$ , then  $w^c \in \omega(T^*)$  and  $w^c(x) > 0$ . Now we have  $(w^c)^{-1}(0, 1] \in T^*$ ,  $x \in (w^c)^{-1}(0, 1]$ . Also it is clear that  $[(w^c)^{-1}(0, 1)]^c = w^{-1}\{1\}$  be a closed in  $T^*$  and  $x \notin w^{-1}\{1\}$ . Since  $(X, T^*)$  is Regular, then there exist  $V, V^* \in T$  such that  $x \in V, V^* \supseteq w^{-1}\{1\}$  and  $V \cap V^* = \phi$ . But by the definition of lower semi continuous functions  $1_V, 1_{V^*} \in \omega(T^*)$  and  $1_V(x) = 1, 1_{V^*} \supseteq 1_{w^{-1}\{1\}}, 1_V \cap 1_{V^*} = 1_{V \cap V^*} = 0$ . Put  $u = 1_V$  and  $v = 1_{V^*}$ , then, we have  $u(x) = 1, v \supseteq 1_{w^{-1}\{1\}}$  and  $u \cap v \leq \alpha$ . Hence  $(X, \omega(T^*))$  is  $\alpha$ -SFR (i).

We can easily be shown that (2) implies (3), (3) implies (5), (2) implies (4), (4) implies (5).

We therefore prove that (5) implies (1).

Let  $(X, \omega(T^*))$  be  $\alpha$ -SFR (iv). Let  $x \in X$  and  $V$  be a closed set in  $T^*$ , such that  $x \notin V$ . This implies that  $V^c \in T^*$  and  $x \in V^c$ . But from the definition of  $\omega(T^*)$ ,  $1_{V^c} \in \omega(T^*)$ , and  $(1_{V^c})^c = 1_V$  closed in  $\omega(T^*)$  and  $1_V(x) = 0$ . Since  $(X, \omega(T^*))$  is  $\alpha$ -SFR (iv), for  $\alpha \in I_1$ , there exist  $u, v \in \omega(T^*)$  such that  $u(x) > \alpha, v \supseteq 1_{(1_V)^{-1}\{1\}} = 1_V$  and  $u \cap v \leq \alpha$ .

Since  $u, v \in \omega(T^*)$ , then  $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in T^*$  and  $x \in u^{-1}(\alpha, 1]$ . Since  $v \geq 1_v$ , then  $v^{-1}(\alpha, 1] \supseteq (1_v)^{-1}(\alpha, 1] = V$ , and  $u \cap v \leq \alpha$  implies  $(u \cap v)^{-1}(\alpha, 1] = u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ . Then by definition,  $(X, T^*)$  is Regular space.

Thus it is seen that  $\alpha$ -SFR (p) is a good extension of its topological counter part (p = i, ii, iii, iv).

**Theorem 5.13:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be continuous, one-one, onto and open map, then

- (a)  $(X, t^*)$  is  $\alpha$ -SFR (i) implies  $(Y, s^*)$  is  $\alpha$ -SFR (i).
- (b)  $(X, t^*)$  is  $\alpha$ -SFR (ii) implies  $(Y, s^*)$  is  $\alpha$ -SFR (ii).
- (c)  $(X, t^*)$  is  $\alpha$ -SFR (iii) implies  $(Y, s^*)$  is  $\alpha$ -SFR (iii).
- (d)  $(X, t^*)$  is  $\alpha$ -SFR (iv) implies  $(Y, s^*)$  is  $\alpha$ -SFR (iv).

**Proof:** Suppose  $(X, t^*)$  be  $\alpha$ -SFR (i). For  $w \in (s^*)^c$  and  $p \in Y$  such that  $w(p) < 1$ ,  $f^{-1}(w) \in (t^*)^c$  as  $f$  is continuous and  $x \in X$  such that  $f(x) = p$  as  $f$  is one-one and onto. Hence  $f^{-1}(w)(x) = w(f(x)) = w(p) < 1$ . Since  $(X, t^*)$  is  $\alpha$ -SFR (i), for  $\alpha \in I_1$ , then there exist  $u, v \in t^*$  such that  $u(x) = 1, v(y) = 1, y \in \{f^{-1}(w)\}^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . This implies that

$$f(u)(p) = \{ \text{Sup } u(x) : f(x) = p \} = 1,$$

$$\text{and } f(v)(f(y)) = \{ \text{Sup } v(y) \} = 1 \text{ as } f(f^{-1}(w)) \subseteq w \Rightarrow f(y) \in w^{-1}\{1\}$$

$$\text{and } f(u \cap v) \leq \alpha \text{ as } u \cap v \leq \alpha \Rightarrow f(u) \cap f(v) \leq \alpha.$$

Now, it is clear that for every  $f(u), f(v) \in s^*$  such that  $f(u)(x) = 1, f(v)(f(y)) = 1, f(y) \in w^{-1}\{1\}$  and  $f(u) \cap f(v) \leq \alpha$ . Hence  $(Y, s^*)$  is  $\alpha$ -SFR (i).

Similarly (b), (c) and (d) can be proved.

**Theorem 5.14:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be a continuous, one-one, onto and closed map then,

- (a)  $(Y, s^*)$  is  $\alpha$  - SFR (i) implies  $(X, t^*)$  is  $\alpha$  - SFR (i).
- (b)  $(Y, s^*)$  is  $\alpha$  - SFR (ii) implies  $(X, t^*)$  is  $\alpha$  - SFR (ii).
- (c)  $(Y, s^*)$  is  $\alpha$  - SFR (iii) implies  $(X, t^*)$  is  $\alpha$  - SFR (iii).
- (d)  $(Y, s^*)$  is  $\alpha$  - SFR (iv) implies  $(X, t^*)$  is  $\alpha$  - SFR (iv).

**Proof:** Suppose  $(Y, s^*)$  be  $\alpha$  - SFR (i). For  $w \in (t^*)^c$  and  $x \in X$  with  $w(x) < 1$ , then  $f(w) \in (s^*)^c$  as  $f$  is closed and we find  $p \in Y$  such that  $f(x) = p$  as  $f$  is one-one. Now we have  $f(w)(p) = \{\text{Sup } w(x) : f(x) = p\} < 1$ . Since  $(Y, s^*)$  is  $\alpha$  - SFR (i), for  $\alpha \in I_1$ , then there exist  $u, v \in s^*$  such that  $u(f(x)) = 1, v(y) = 1, y \in (f(w))^{-1}\{1\}$  and  $u \cap v \leq \alpha$ . This implies that  $f^{-1}(u), f^{-1}(v) \in t^*$  as  $f$  is continuous and  $u, v \in s^*$ . Now  $f^{-1}(u)(x) = u(f(x)) = u(p) = 1$  and  $f^{-1}(v)(q) = v(f(q)) = v(y) = 1$  as  $f(q) = y, y \in (f(w))^{-1}\{1\}$ , i.e.,  $f(p) \in (f(w))^{-1}\{1\} \Rightarrow q \in w^{-1}\{1\}$  and  $f^{-1}(u) \cap f^{-1}(v) = f^{-1}(u \cap v) \leq \alpha$  as  $u \cap v \leq \alpha$ . Now we observe that  $\exists f^{-1}(u), f^{-1}(v) \in t^*$  such that  $f^{-1}(u)(x) = 1, f^{-1}(v)(q) = 1, q \in w^{-1}\{1\}$  and  $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$ . Hence  $(X, t^*)$  is  $\alpha$  - SFR (i).

Similarly, (b), (c) and (d) can be proved.

# CHAPTER 6

## Normal Spaces in Supra Fuzzy Topology

### 6.1 Introduction

In this chapter, we introduce and study normal property in supra fuzzy topological spaces and obtain their several features.

**Definition 6.2:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is  $\alpha$  - SFN(i) space if and only if for all  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq \alpha$ , there exist  $u, u^* \in t^*$  such that  $u(x) = 1$ , for all  $x \in w^{-1}\{1\}$ ,  $u^*(y) = 1$ , for all  $y \in w^{*-1}\{1\}$  and  $u \cap u^* = 0$ .
- (b)  $(X, t^*)$  is  $\alpha$  - SFN (ii) space if and only if for all  $w, w^* \in (t^*)^c$  with  $w \cap w^* = 0$ , there exist  $u, u^* \in t^*$  such that  $u(x) = 1$ , for all  $x \in w^{-1}\{1\}$ ,  $u^*(y) = 1$ , for all  $y \in w^{*-1}\{1\}$  and  $u \cap u^* \leq \alpha$ .
- (c)  $(X, t^*)$  is a  $\alpha$  - SFN(iii) space if and only if for all  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq \alpha$ , there exist  $u, u^* \in t^*$  such that  $u(x) = 1$  for all  $x \in w^{-1}\{1\}$ ,  $u^*(y) = 1$ , for all  $y \in w^{*-1}\{1\}$  and  $u \cap u^* \leq \alpha$ .

**Theorem 6.3:** The following implications are true:

- (a)  $(X, t^*)$  is 0 - SFN (i) if and only if  $(X, t^*)$  is 0 - SFN (ii).
- (b)  $(X, t^*)$  is 0 - SFN (ii) if and only if  $(X, t^*)$  is 0 - SFN (iii).

**Proof:** Suppose  $(X, t^*)$  be  $\alpha$  -SFN (i). We have to show that  $(X, t^*)$  is 0 -SFN (ii). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* = 0$ . This implies that  $w \cap w^* \leq 0$ . Since  $(X, t^*)$  is 0 - SFN(i),

for  $\alpha \in I_1$ , there exist  $u, u^* \in t^*$  such that  $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$  and  $u \cap u^* = 0$ . Then we can write it as  $u \cap u^* \leq 0$ . Hence by definition,  $(X, t^*)$  is  $\alpha$ -SFN (ii).

Conversely, suppose that  $(X, t^*)$  is 0-SFN (ii). We have to prove that  $(X, t^*)$  is -FN (i). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq 0$ , i.e.,  $w \cap w^* = 0$ . Since  $(X, t^*)$  is 0-FN(ii), there exist  $u, u^* \in t^*$  such that  $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ , and  $u \cap u^* \leq 0$ , i.e.,  $u \cap u^* = 0$ . Hence by definition,  $(X, t^*)$  is 0-SFN (i).

Next, suppose that  $(X, t^*)$  is 0-SFN (ii). We have to prove that  $(X, t^*)$  is 0-SFN (iii). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq 0$ , i.e.,  $w \cap w^* = 0$ . Since  $(X, t^*)$  is 0-SFN (ii), there exist  $u, u^* \in t^*$  such that  $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$  and  $u \cap u^* \leq 0$ . Hence by definition,  $(X, t^*)$  is  $\alpha$ -SFN (iii).

Conversely, suppose that  $(X, t^*)$  is 0-SFN (iii). We have to prove that  $(X, t^*)$  is 0-SFN (ii). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* = 0$ , i.e.,  $w \cap w^* \leq 0$ . Since  $(X, t^*)$  is 0-SFN (iii), there exist  $u, u^* \in t^*$  such that  $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$  and  $u \cap u^* \leq 0$ . Hence by definition  $(X, t^*)$  is  $\alpha$ -SFN (ii).

**Theorem 6.4:** Let  $(X, t^*)$  be a supra topological space and  $\alpha, \beta \in t^*$  with  $0 \leq \alpha \leq \beta < 1$ , then

(a)  $(X, t^*)$  is  $\beta$ -SFN (i) implies  $(X, t^*)$  is  $\alpha$ -SFN (i).

(b)  $(X, t^*)$  is  $\alpha$ -SFN (ii) implies  $(X, t^*)$  is  $\beta$ -SFN (ii).

**Proof:** First, suppose that  $(X, t^*)$  is  $\beta$  - SFN (i). We have to prove that  $(X, t^*)$  is  $\alpha$  -SFN (i). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq \alpha$ . Since  $0 \leq \alpha \leq \beta < 1$ , then  $w \cap w^* \leq \beta$ . Since  $(X, t^*)$  is  $\beta$  - SFN(i), for  $\beta \in I_1$ , there exist  $u, u^* \in t^*$  such that  $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$  and  $u \cap u^* = 0$ . Thus by definition,  $(X, t^*)$  is a  $\alpha$  - SFN (i).

Next, suppose that  $(X, t^*)$  is  $\alpha$  - SFN (ii). We have to prove that  $(X, t^*)$  is  $\beta$  - SFN (ii). Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* = 0$ . Since  $(X, t^*)$  is  $\alpha$  -SFN(ii), there exist  $u, u^* \in t^*$  such that  $u(x) = 1, x \in w^{-1}\{1\}, u^*(y) = 1, y \in w^{*-1}\{1\}$  and  $u \cap u^* \leq \alpha$ . Since  $0 \leq \alpha \leq \beta < 1$ , then  $u \cap u^* \leq \beta$ . Then we have  $(X, t^*)$  is  $\beta$  - SFN (ii).

**Theorem 6.5:** Let  $(X, t^*)$  be a supra fuzzy topological space, and  $I_\alpha(t^*) = \{u^{-1}(\alpha, 1] : u \in t^*\}$ , then  $(X, t^*)$  is 0 - SFN(iii) space implies  $(X, I_0(t^*))$  is Normal space.

**Proof:** Suppose that  $(X, t^*)$  be a  $\alpha$  - FN (iii) space. We have to prove that  $(X, I_0(t^*))$  is Normal space. Let  $V, V^*$  be closed set in  $I_0(t^*)$ , and  $V \cap V^* = \phi$ . Then  $V^c, V^{*c} \in I_0(t^*)$  and  $(V \cap V^*)^c = V^c \cup V^{*c} = X$ . Since  $V^c, V^{*c} \in I_0(t^*)$ , then there exist  $u, u^* \in t^*$  such that  $V^c = u^{-1}(0, 1]$  and  $V^{*c} = u^{*-1}(0, 1]$  and  $u^{-1}(0, 1] \cup u^{*-1}(0, 1] = V^c \cup V^{*c} = X$ . Hence  $(u \cup u^*)^{-1}(0, 1] = X$ . Now we find  $u^c, u^{*c} \in (t^*)^c$  such that  $((u \cup u^*)^{-1}(0, 1])^c = \phi$ . This implies that  $(u \cup u^*)^c = u^c \cap u^{*c} = 0$ . Since  $(X, t^*)$  is 0 - SFN(iii), there exist  $v, v^* \in t^*$  such that  $v \geq 1_{(u^c)^{-1}\{1\}}, v^* \geq 1_{(u^{*c})^{-1}\{1\}}, v \cap v^* = 0$ . But from the definition of  $I_0(t^*)$ ,  $v^{-1}(0, 1], v^{*-1}(0, 1] \in I_0(t^*)$ , and we get  $v^{-1}(0, 1] \supseteq 1_{(u^c)^{-1}\{1\}}(0, 1], v^{*-1}(0, 1] \supseteq 1_{(u^{*c})^{-1}\{1\}}(0, 1]$ ,  $(v \cap v^*)^{-1}(0, 1] = \phi$ . Put  $W = v^{-1}(0, 1], W^* = v^{*-1}(0, 1]$  in

$I_0(t^*)$ . Then finally we find,  $W \supseteq V$ ,  $W^* \supseteq V^*$  and  $W \cap W^* = \phi$ . Hence by definition,  $(X, I_0(t^*))$  is Normal space.

**Theorem 6.6:** Suppose  $(X, T^*)$  be supra topological space, then consider the following statements:

- (a)  $(X, T^*)$  be a Normal space
- (b)  $(X, \omega(T^*))$  be  $\alpha$  - SFN (i) space.
- (c)  $(X, \omega(T^*))$  be  $\alpha$  - SFN (ii) space.
- (d)  $(X, \omega(T^*))$  be  $\alpha$  - SFN (iii) space.

Then (a) implies (b); (b) implies (c) and (c)  $\Rightarrow$  (a).

**Proof:** First suppose that  $(X, T^*)$  be a Normal space. We have to prove that  $(X, \omega(T^*))$  be  $\alpha$  - SFN (i) space. Let  $w, w^*$  be closed in  $\omega(T^*)$  and  $w \cap w^* \leq \alpha$ . Then we have  $w^c, w^{*c} \in \omega(T^*)$  and  $(w \cap w^*)^c \geq 1 - \alpha > 0$ . But from the definition of  $\omega(T^*)$ ,  $(w^c)^{-1}(0, 1], (w^{*c})^{-1}(0, 1] \in T^*$ . Now we see that  $((w \cap w^*)^c)^{-1}(0, 1] = X$ , and we also see that  $((w^c)^{-1}(0, 1])^c = w^{-1}\{1\}$  and  $((w^{*c})^{-1}(0, 1])^c = (w^*)^{-1}\{1\}$  be closed in  $T^*$ . Now  $((w \cap w^*)^c)^{-1}(0, 1])^c = (w \cap w^*)^{-1}\{1\} = w^{-1}\{1\} \cap w^{*-1}\{1\} = \phi$ . Since  $(X, t^*)$  is supra fuzzy Normal, then there exist  $V, V^* \in T^*$  such that  $V \supseteq w^{-1}\{1\}, V^* \supseteq w^{*-1}\{1\}$  and  $V \cap V^* = \phi$ . But from the definition of  $\omega(T^*)$ ,  $1_V, 1_{V^*} \in \omega(T)$ ,  $1_V \geq 1_{w^{-1}\{1\}}, 1_{V^*} \geq 1_{(w^*)^{-1}\{1\}}$  and  $1_{V \cap V^*} = 0$ . Hence  $(X, \omega(T^*))$  is  $\alpha$  - SFN (i) space.

We can easily show that (b)  $\Rightarrow$  (c)  $\Rightarrow$  (d).

We, therefore, prove that (d)  $\Rightarrow$  (a).

Suppose  $(X, \omega(T^*))$  is  $\alpha$  - SFN (iii). We have to prove that  $(X, T^*)$  is Normal space. Let  $V, V^* \in (T^*)^c$  and  $V \cap V^c = \phi$ . Then we have  $1_V, 1_{V^*}$  be closed in  $\omega(T^*)$  and  $1_{V \cap V^*} = 0$ .

Since  $(X, w(T^*))$  is  $\alpha$ -SFN (iii), then there exist  $u, u^* \in w(T^*)$  such that  $u \geq 1_v, u^* \geq 1_{v^*}$  and  $u \cap u^* \leq \alpha$ . But from the definition of  $w(T^*)$ ,  $u^{-1}(\alpha, 1], u^{*-1}(\alpha, 1] \in T^*$  and  $u^{-1}(\alpha, 1] \supseteq (1_v)^{-1}(\alpha, 1] = V, u^{*-1}(\alpha, 1] \supseteq (1_{v^*})^{-1}(\alpha, 1] = V^*$  and  $u^{-1}(\alpha, 1] \cap u^{*-1}(\alpha, 1] = (u \cap u^*)^{-1}(\alpha, 1] = \phi$ . Hence by definition,  $(X, T^*)$  is Normal space.

Thus it is seen that  $\alpha$ -SFN (p) is a good extension of its topological counterpart. (p = i, ii, iii).

**Theorem 6.7:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be a continuous, one-one, onto and open map then,

- (a)  $(X, t^*)$  is  $\alpha$ -SFN (i) implies  $(Y, s^*)$  is  $\alpha$ -SFN(i).
- (b)  $(X, t^*)$  is  $\alpha$ -SFN (ii) implies  $(Y, s^*)$  is  $\alpha$ -SFN (ii).
- (c)  $(X, t^*)$  is  $\alpha$ -SFN(iii) implies  $(Y, s^*)$  is  $\alpha$ -SFN(iii).

**Proof:** Suppose  $(X, t^*)$  be  $\alpha$ -SFN (i). We have to prove that  $(Y, s^*)$  is  $\alpha$ -SFN (i). Let  $w, w^* \in (s^*)^c$  with  $w \cap w^* \leq \alpha$  then  $f^{-1}(w), f^{-1}(w^*) \in (t^*)^c$  as  $f$  is continuous. Now  $f^{-1}(w \cap w^*) \leq \alpha$  as  $w \cap w^* \leq \alpha \Rightarrow f^{-1}(w) \cap f^{-1}(w^*) \leq \alpha$ . Since  $(X, t^*)$  is  $\alpha$ -SFN(i), for  $\alpha \in I_1$ , then there  $u, u^* \in t^*$  such that  $u(x) = 1, x \in (f^{-1}(w))^{-1}\{1\}$ ,  $u^*(y) = 1, y \in (f^{-1}(w^*))^{-1}\{1\}$  and  $u \cap u^* = 0$ . This implies that  $f(u), f(v) \in s$  as  $f$  is open. Now  $f(u)(p) = \{ \text{Sup } u(x) : f(x) = p \}$ ,  $f^{-1}(p) \in (f^{-1}(w))^{-1}\{1\}$ , i.e.,  $f(u)(p) = 1$ ,  $p \in w^{-1}\{1\}$  and  $f(u^*)(q) = \{ \text{Sup } u^*(y); f(y) = q \}$ ,  $f^{-1}(q) \in (f^{-1}(w^*))^{-1}\{1\}$ , i.e.,  $f(u^*)(q) = 1, q \in w^{*-1}\{1\}$  and  $f(u) \cap f(u^*) = f(u \cap u^*) = 0$  as  $u \cap u^* = 0$ . Now it is clear that there are  $f(u), f(u^*) \in s^*$  such that  $f(u)(p) = 1, p \in w^{-1}\{1\}$ ,  $f(u^*)(q) = 1, q \in w^{*-1}\{1\}$  and  $f(u) \cap f(u^*) = 0$ . Hence  $(Y, s^*)$  is  $\alpha$ -SFN (i).

Similarly, (b) and (c) can be proved.



**Theorem 6.8:** Let  $(X, t^*)$  and  $(Y, s^*)$  be two supra fuzzy topological spaces and  $f: X \longrightarrow Y$  be continuous, one-one, onto and closed map then,

(a)  $(Y, s)$  is  $\alpha$ -SFN(i) implies  $(X, t^*)$  is  $\alpha$ -SFN(i).

(b)  $(Y, s^*)$  is  $\alpha$ -SFN(ii) implies  $(X, t^*)$  is  $\alpha$ -SFN(ii).

(c)  $(Y, s^*)$  is  $\alpha$ -SFN (iii) implies  $(X, t^*)$  is  $\alpha$ -SFN (iii).

**Proof:** Suppose  $(Y, s^*)$  be  $\alpha$ -SFN (i). We have to prove that  $(X, t^*)$  is  $\alpha$ -SFN(i).

Let  $w, w^* \in (t^*)^c$  with  $w \cap w^* \leq \alpha$ , then  $f(w), f(w^*) \in s^c$  as  $f$  is closed and

$f(w) \cap f(w^*) = f(w \cap w^*) \leq \alpha$ , as  $w \cap w^* \leq \alpha$ . Since  $(Y, s^*)$  is  $\alpha$ -SFN(i), for  $\alpha \in I_1$ ,

then there exist  $u, u^* \in s^*$  such that  $u(x) = 1, x \in (f(w))^{-1}\{1\}, u^*(y) = 1, y \in (f(w^*))^{-1}\{1\}$

and  $u \cap u^* = 0$ . This implies that  $f^{-1}(u), f^{-1}(u^*) \in t^*$  as  $f$  is continuous and  $u, u^* \in s^*$ .

Now  $f^{-1}(u)(p) = u(f(p)) = u(x) = 1$  as  $f(p) = x \in (f(w))^{-1}\{1\}$

i.e.,  $f^{-1}(u)(p) = 1, p \in w^{-1}\{1\}$

$f^{-1}(u^*)(q) = u^*(f(q)) = u^*(y) = 1$  as  $f(q) = y \in (f(w^*))^{-1}\{1\}$

i.e.,  $f^{-1}(u^*)(q) = 1, q \in w^{*-1}\{1\}$

and  $f^{-1}(u) \cap f^{-1}(u^*) = f^{-1}(u \cap u^*) = 0$ .

Then we have there exist  $f^{-1}(u), f^{-1}(u^*) \in t^*$  such that  $f^{-1}(u)(p) = 1, p \in w^{-1}\{1\}$ ,

$f^{-1}(u^*)(q) = 1, q \in w^{*-1}\{1\}$  and  $f^{-1}(u) \cap f^{-1}(u^*) = 0$ . Hence  $(X, t^*)$  is  $\alpha$ -SFN (i).

Similarly, (b) and (c) can be proved.

# CHAPTER 7

## $R_0$ and $R_1$ Spaces in Supra Fuzzy Topology

### 7.1 Introduction

We introduce and study some  $R_0$  and  $R_1$  properties in supra fuzzy topological spaces and obtain their several features in this chapter.

**Definition 7.2:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is an  $\alpha$ - $R_0$  (i) space if and only if for all  $x, y \in X$  with  $x \neq y$ , whenever there exist  $u \in t^*$  with  $u(x) = 1$  and  $u(y) \leq \alpha$ , then there exist  $v \in t^*$  with  $v(x) \leq \alpha$  and  $v(y) = 1$ .
- (b)  $(X, t^*)$  is an  $\alpha$ - $R_0$  (ii) space if and only if for all  $x, y \in X$ ,  $x \neq y$ , whenever  $u \in t^*$  with  $u(x) = 0$  and  $u(y) > \alpha$ , then there exists  $v \in t^*$  with  $v(x) > \alpha$  and  $v(y) = 0$ .
- (c)  $(X, t^*)$  is an  $\alpha$ - $R_0$  (iii) space if and only if for all  $x, y \in X$  with  $x \neq y$ , whenever there exists  $u \in t^*$  with  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ , then there exists  $v \in t^*$  with  $0 \leq v(y) \leq \alpha < v(x) \leq 1$ .
- (d)  $(X, t^*)$  is an  $R_0$  (iv) space if and only if for all  $x, y \in X$  with  $x \neq y$ , whenever there exists  $u \in t^*$  with  $u(x) < u(y)$ , then there exists  $v \in t^*$  with  $v(x) > v(y)$ .

The following examples show that  $\alpha$ - $R_0$  (i),  $\alpha$ - $R_0$  (ii),  $\alpha$ - $R_0$  (iii) and  $R_0$  (iv) are all independent.

**Example 7.3:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 1, u(y) = 0$  and  $v(x) = 0.45, v(y) = 1$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.55$ ,  $(X, t^*)$  is  $\alpha$ - $R_0$ (i), but  $(X, t^*)$  is not  $\alpha$ - $R_0$  (ii).

**Example 7.4:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 0, u(y) = 1$  and  $v(x) = 0.73, v(y) = 0$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.63$ ,  $(X, t^*)$  is  $\alpha - R_0$  (ii), but  $(X, t^*)$  is not  $\alpha - R_0$  (i).

**Example 7.5:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 1, u(y) = 0$  and  $v(x) = 0.32, v(y) = 0.69$ . Consider the supra fuzzy topology generated by  $\{0, u, v, 1, \text{Constant}\}$ . Then by definition, for  $\alpha = 0.52$ ,  $(X, t^*)$  is  $\alpha - R_0$  (iii), but  $(X, t^*)$  is not  $\alpha - R_0$  (i) and  $(X, t^*)$  is not  $\alpha - R_0$  (ii).

**Example 7.6:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 1, u(y) = 0$  and  $v(x) = 0.24, v(y) = 0.48$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.53$ ,  $(X, t^*)$  is  $\alpha - R_0$  (iv), but  $(X, t^*)$  is not  $\alpha - R_0$  (i),  $(X, t^*)$  is not  $\alpha - R_0$  (ii), and  $(X, t^*)$  is not  $\alpha - R_0$  (iii).

**Example 7.7:** Let  $X = \{x, y, z\}$  and  $u, v, w \in I^X$  are defined by  $u(x) = 1, u(y) = 1, u(z) = 0$  and  $v(x) = 0, v(y) = 0, v(z) = 1$  and  $w(x) = 0.92, w(y) = 0.52, w(z) = 0$ . Consider the fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, w, 1, \text{Constants}\}$ . Then for  $\alpha = 0.63$ , it can easily shown that  $(X, t^*)$  is  $\alpha - R_0$  (i) and  $(X, t^*)$  is  $\alpha - R_0$  (ii). But we observe  $(X, t^*)$  is not  $\alpha - R_0$  (iii), and  $(X, t^*)$  is not  $R_0$  (iv), Since  $w(x) > \alpha \geq w(y)$  but there does not exist  $q \in t^*$  such that  $q(x) \leq \alpha < q(y)$ .

**Example 7.8:** Let  $X = \{x, y, z\}$  and  $u, v \in I^X$  are defined by  $u(x) = 0.83, u(y) = 0.41, u(z) = 0.36$ , and  $v(x) = 0.36, v(y) = 0.83, v(z) = 0.24$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then by definition, for  $\alpha = 0.5$ ,  $(X, t^*)$  is  $\alpha - R_0$  (iii), but  $(X, t^*)$  is not  $R_0$  (iv), since  $u(y) > u(z)$  but we have no  $q \in t^*$  such that  $q(y) < q(z)$ .

**Theorem 7.9:** Prove that  $(X, t^*)$  is  $0-R_0$  (ii) if and only if  $(X, t^*)$  is  $0-R_0$  (iii).

**Proof:** The proof is trivial.

**Theorem 7.10:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $I_\alpha(t^*) = \{u^{-1}(\alpha, 1] \mid u \in t^*\}$ . Then the following is true:

- (a)  $(X, t^*)$  is  $\alpha-R_0$  (iii) space if and only if  $(X, I_\alpha(t^*))$  is  $R_0$  space.
- (b) if  $(X, t^*)$  is  $\alpha-R_0$  (i) space, then  $(X, I_\alpha(t^*))$  is not  $R_0$  space and conversely.
- (c) if  $(X, t^*)$  is  $\alpha-R_0$  (ii) space then  $(X, I_\alpha(t^*))$  is not  $R_0$  space and conversely.
- (d) if  $(X, t^*)$  is  $R_0$  (iv), then  $(X, I_\alpha(t^*))$  is not  $R_0$  space and conversely.

**Proof:** Let  $(X, t^*)$  be  $\alpha-R_0$  (iii). We have to prove that  $(X, I_\alpha(t^*))$  is  $R_0$ . Let  $x, y \in X$  with  $x \neq y$  and  $M \in I_\alpha(t^*)$  with  $x \in M, y \notin M$  or  $x \notin M, y \in M$ . Suppose that  $x \in M, y \notin M$ . We can write,  $M = u^{-1}(\alpha, 1]$ , for some  $u \in t^*$ . Then we have  $u(x) > \alpha, u(y) \leq \alpha$ , i.e.,  $0 \leq u(y) \leq \alpha < u(x) \leq 1$ . Since  $(X, t^*)$  is  $\alpha-R_0$ (iii),  $\alpha \in I_1$ , then there exists  $v \in t^*$  such that  $0 \leq v(x) \leq \alpha < v(y) \leq 1$ , i.e.,  $v(x) \leq \alpha, v(y) > \alpha$ . It follows that  $x \notin v^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$  and also  $v^{-1}(\alpha, 1] \in I_\alpha(t^*)$ . Thus  $(X, I_\alpha(t^*))$  is  $R_0$ .

Conversely, suppose that  $(X, I_\alpha(t^*))$  is  $R_0$ . We have to prove that  $(X, t^*)$  is  $\alpha-R_0$  (iii). Let  $x, y \in X$  with  $x \neq y$  and  $u \in t^*$  with  $0 \leq u(x) \leq \alpha < u(y) \leq 1$ , i.e.,  $u(x) \leq \alpha, u(y) > \alpha$ , it follows that  $x \notin u^{-1}(\alpha, 1], y \in u^{-1}(\alpha, 1]$ , and  $u^{-1}(\alpha, 1] \in I_\alpha(t)$ , for every  $u \in t^*$ . Since  $(X, I_\alpha(t^*))$  is  $R_0$ , then there exists  $M \in I_\alpha(t^*)$  such that  $x \in M, y \notin M$ . We can write  $M = v^{-1}(\alpha, 1]$ , where  $v \in t^*$ , it follows that  $v(x) > \alpha, v(y) \leq \alpha$ , i.e.,  $0 \leq v(y) \leq \alpha < v(x) \leq 1$ . Thus  $(X, t^*)$  is  $\alpha-R_0(t^*)$ , i.e., (a) is proved.

**Example 7.11:** Let  $X = \{x, y, z\}$  and  $u, v \in I^X$  are defined by  $u(x) = 1, u(y) = 0, u(z) = 0.8$  and  $v(x) = 0, v(y) = 1, v(z) = 0.7$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then for  $\alpha = 0.6$ , we have,  $(X, t^*)$  is

$\alpha - R_0$  (i). Now,  $I_\alpha(t^*) = \{X, \Phi, \{x, z\}, \{y, z\}, \{z\}\}$ . It is observed that  $(X, t^*)$  is not  $R_0$  space, since  $y, z \in X$ ,  $y \neq z$  and  $\{x, z\} \in I_\alpha(t^*)$ , with  $z \in \{x, z\}$ ,  $y \notin \{x, z\}$ , but no such  $U \in I_\alpha(t^*)$  with  $x \notin U, y \in U$ .

**Example 7.12:** Let  $X = \{x, y, z\}$  and  $u, v \in I^X$  are defined by  $u(x) = 0.3, u(y) = 0, u(z) = 0.8$ , and  $v(x) = 0.8, v(y) = 1, v(z) = 0$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constants}\}$ . Then, for  $\alpha = 0.5$ , we have,  $(X, t^*)$  is  $\alpha - R_0$  (ii) and  $(X, t^*)$  is also  $R_0$  (iv). Now  $I_\alpha(t^*) = \{X, \Phi, \{z\}, \{y\}, \{y, z\}\}$ . It is observed that  $(X, I_\alpha(t^*))$  is not  $R_0$  space, since  $x, y \in X, x \neq y$  and  $\{y\} \in I_\alpha(t^*)$  with  $x \notin \{y\}, y \in \{y\}$ , but no such  $U \in I_\alpha(t^*)$  with  $x \in U, y \notin U$ .

**Example 7.13:** Let  $X = \{x, y\}$  and  $u, v, w \in I^X$  are defined by  $u(x) = 1, u(y) = 0, v(x) = 0.4, v(y) = 0.9, w(x) = 0.7, w(y) = 0.3$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, w, 1, \text{Constants}\}$ . Then for  $\alpha = 0.6$ , we have,  $(X, t^*)$  is not  $\alpha - R_0$  (i) and  $(X, t^*)$  is not  $\alpha - R_0$  (ii). Now,  $I_\alpha(t^*) = \{X, \Phi, \{x\}, \{y\}\}$ . Then we see that  $I_\alpha(t^*)$  is a topology on  $X$  and  $(X, I_\alpha(t^*))$  is  $R_0$ .

**Example 7.14:** Let  $X = \{x, y\}$  and  $u, v \in I^X$  are defined by  $u(x) = 0.4, u(y) = 0.5, v(x) = 0.3$ , and  $v(y) = 0.4$ . Consider the supra fuzzy topology  $t^*$  on  $X$  generated by  $\{0, u, v, 1, \text{Constant}\}$ . Then, for  $\alpha = 0.6$ , we have  $(X, t^*)$  is not  $\alpha - R_0$  (iv). Now,  $I_\alpha(t^*) = \{X, \Phi\}$ . Then  $I_\alpha(t^*)$  is a topology on  $X$  and  $(X, I_\alpha(t^*))$  is  $R_0$ .

Hence the proof is complete.

**Theorem 7.15:** Let  $(X, T^*)$  be a supra topological space. Then  $(X, T^*)$  is  $R_0$ , if and only if  $(X, w(T^*))$  is  $\alpha - R_0$  (p), where  $p = i, ii, iii, iv$ .

**Proof:** Let  $(X, w(T^*))$  be  $\alpha - R_0$  (i). Let  $x, y \in X$  with  $x \neq y$  and  $U \in T^*$  with  $x \in U, y \notin U$ . But  $1_U \in w(T^*)$  and  $1_U(x) = 1, 1_U(y) = 0$ . Now, we have  $1_U \in w(T^*)$  with

$1_U(x) = 1, 1_U(y) \leq \alpha$ . Since  $(X, w(T^*))$  is  $\alpha - R_0$  (i), there exists  $v \in w(T^*)$  such that  $v(x) \leq \alpha, v(y) = 1$ . Then  $x \notin v^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$  as  $v(x) \leq \alpha, v(y) = 1$  and also there exists  $v^{-1}(\alpha, 1] \in T^*$ . Thus  $(X, T^*)$  is  $R_0$ -space.

Conversely, suppose that  $(X, T^*)$  be a  $R_0$ -space. We have to prove that  $(X, w(T^*))$  is  $\alpha - R_0$ (i). Let  $x, y \in X$  with  $x \neq y$  and there exists  $u \in w(T^*)$  such that  $u(x) = 1, u(y) \leq \alpha$ . Then  $x \in u^{-1}(\alpha, 1], y \notin u^{-1}(\alpha, 1]$  as  $u(x) = 1, u(y) \leq \alpha$ . Hence  $u^{-1}(\alpha, 1] \in T^*$ . Since  $(X, T^*)$  is  $R_0$ , then there exists  $V \in T^*$  such that  $x \notin V, y \in V$ , but  $1_V \in w(T^*)$  and  $1_V(x) = 0, 1_V(y) = 1$ , i.e., there exists  $1_V \in w(T^*)$  such that  $1_V(x) \leq \alpha, 1_V(y) = 1$ . Thus  $(X, w(T^*))$  is  $\alpha - R_0$ (i).

Hence  $(X, T^*)$  is  $R_0$  if and only if  $(X, w(T^*))$  is  $\alpha - R_0$ (i).

In the same way, we can prove that

- (a)  $(X, T^*)$  is  $R_0$  if and only if  $(X, w(T^*))$  is  $\alpha - R_0$ (ii).
- (b)  $(X, T^*)$  is  $R_0$  if and only if  $(X, w(T^*))$  is  $\alpha - R_0$ (iii).
- (c)  $(X, T^*)$  is  $R_0$  if and only if  $(X, w(T^*))$  is  $R_0$ (iv)

Thus it is seen that  $\alpha - R_0$  (p) is a good extension of its topological counterpart (p = i, ii, iii, iv).

**Theorem 7.16:** Let  $(X, t^*)$  be a supra fuzzy topological space and  $A \subseteq X$ ,  $t^*_{A} = \{u/A : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is an  $\alpha - R_0$  (i) if and only if  $(A, t^*_A)$  is an  $\alpha - R_0$  (i).
- (b)  $(X, t^*)$  is an  $\alpha - R_0$  (ii) if and only if  $(A, t^*_A)$  is an  $\alpha - R_0$  (ii).
- (c)  $(X, t^*)$  is an  $\alpha - R_0$  (iii) if and only if  $(A, t^*_A)$  is an  $\alpha - R_0$  (iii)
- (d)  $(X, t^*)$  is an  $R_0$ (iv) if and only if  $(A, t^*_A)$  is an  $R_0$  (iv).

**Proof:** Suppose that  $(X, t^*)$  is  $\alpha - R_0$  (i). Then for  $x, y \in A$ , with  $x \neq y$  and  $u \in t^*_A$  such that  $u(x) = 1, u(y) \leq \alpha$ , then also  $x, y \in X, x \neq y$ . But we can write  $u = w/A$ , where  $w \in t^*$  and hence  $w(x) = 1, w(y) \leq \alpha$ . Since  $(X, t^*)$  is  $\alpha - R_0$  (i), then there exists  $m \in t^*$  such that  $m(x) \leq \alpha, m(y) = 1$ . But from the definition  $m/A \in t^*_A$ , for every  $m \in t^*$  and  $m/A(x) \leq \alpha, m/A(y) = 1$ . Thus  $(A, t^*_A)$  is  $\alpha - R_0$  (i). i.e., (a) proved.

Similarly (b), (c) and (d) can be proved.

**Theorem 7.17:** Given  $(X_i, t^*_i), i \in \Lambda$  be supra fuzzy topological spaces and  $X = \prod_{i \in \Lambda} X_i$  and  $t^*$  be a product supra fuzzy topology on  $X$ . Then

(a)  $\forall i \in \Lambda, (X_i, t^*_i)$  is  $\alpha - R_0$  (i) if and only if  $(X, t^*)$  is  $\alpha - R_0$  (i).

(b)  $\forall i \in \Lambda, (X_i, t^*_i)$  is  $\alpha - R_0$  (ii) if and only if  $(X, t^*)$  is  $\alpha - R_0$  (ii).

(c)  $\forall i \in \Lambda, (X_i, t^*_i)$  is  $\alpha - R_0$  (iii) if and only if  $(X, t^*)$  is  $\alpha - R_0$  (iii).

(d)  $\forall i \in \Lambda, (X_i, t^*_i)$  is  $R_0$  (iv) if and only if  $(X, t^*)$  is  $R_0$  (iv).

**Proof:** Let  $(X_i, t^*_i), i \in \Lambda$  be  $\alpha - R_0$  (i). We have to prove that  $(X, t^*)$  is  $\alpha - R_0$  (i). Let  $x, y \in X$ , with  $x \neq y$  and  $u \in t^*$  such that  $u(x) = 1, u(y) \leq \alpha$ . But we have  $u(x) = \min\{u_i(x_i) : i \in \Lambda\}$  and  $u(y) = \min\{u_i(y_i) : i \in \Lambda\}$  and hence we can find an  $u_i \in t^*_i$  and  $x_i \neq y_i$  such that  $u_i(x_i) = 1$  and  $u_i(y_i) \leq \alpha$ . Since  $(X_i, t^*_i), i \in \Lambda$  is  $\alpha - R_0$  (i),  $\alpha \in I_1$ , then there exist  $v_i \in t^*_i$ , such that  $v_i(x_i) \leq \alpha, v_i(y_i) = 1$ . But  $\pi_i(x) = x_i$  and  $\pi_i(y) = y_i$  and hence  $v_i(\pi_i(x)) \leq \alpha, v_i(\pi_i(y)) = 1$ . It follows that there exists  $u_i \circ \pi_i \in t^*$  such that  $(v_i \circ \pi_i)(x) \leq \alpha, (v_i \circ \pi_i)(y) = 1$ . Thus  $(X, t^*)$  is  $\alpha - R_0$  (i).

Conversely, suppose that  $(X, t^*)$  is  $\alpha - R_0$  (i). We have to prove that  $(X_i, t^*_i), i \in \Lambda$ , is  $\alpha - R_0$  (i). Let for some  $i \in \Lambda, a_i$  be a fixed element in  $X_i$ , suppose that  $A_i = \{x \in X = \prod_{i \in \Lambda} X_i / x_j = a_j \text{ for some } i \neq j\}$ . So that  $A_i$  is the subset of  $X$ , and this implies that  $(A_i, t^*_{A_i})$  is also the subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha - R_0$  (i), then

$(A_i, t^*_{A_i})$  is also  $\alpha - R_0(i)$  and  $A_i$  is a homeomorphic image of  $X_i$ . Thus  $(X_i, t^*_{X_i})$  is  $\alpha - R_0(i)$ , i.e., (a) is proved.

Similarly, (b), (c) and (d) can be proved.

**Definition 7.18:** Let  $(X, t^*)$  be a fuzzy topological space and  $\alpha \in I_1$ . Then

- (a)  $(X, t^*)$  is said to be  $\alpha - R_1(i)$  if and only if for all  $x, y \in X$  with  $x \neq y$  whenever there exists  $w \in t^*$  with  $w(x) \neq w(y)$ , then there exist  $u, v \in t^*$  such that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ .
- (b)  $(X, t^*)$  is said to be  $\alpha - R_1(ii)$  if and only if for all  $x, y \in X$  with  $x \neq y$  whenever there exists  $w \in t^*$  with  $w(x) \neq w(y)$ , then there exist  $u, v \in t^*$  such that  $u(x) = v(y) = 1$  and  $u \cap v \leq \alpha$ .
- (c)  $(X, t^*)$  is said to be  $\alpha - R_1(iii)$  if and only if for all  $x, y \in X$  with  $x \neq y$  whenever there exists  $w \in t^*$  with  $w(x) \neq w(y)$  then there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v = 0$ .
- (d)  $(X, t^*)$  is said to be  $\alpha - R_1(iv)$  if and only if for all  $x, y \in X$  with  $x \neq y$  whenever there exists  $w \in t^*$  with  $w(x) \neq w(y)$  then there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ .

**Lemma 7.19:** The following implication is true:

- (a)  $(X, t^*)$  is  $\alpha - R_1(i)$  implies  $\alpha - R_1(ii)$  implies  $\alpha - R_1(iii)$  but  $\alpha - R_1(iv)$  does not imply  $\alpha - R_1(iii)$ . So, not imply  $\alpha - R_1(i)$  and  $\alpha - R_1(ii)$ .
- (b)  $(X, t^*)$  is  $\alpha - R_1(i)$  does not imply  $\alpha - R_1(iv)$  and also  $\alpha - R_1(ii)$  and  $\alpha - R_1(iii)$  does not imply  $\alpha - R_1(iv)$ .

**Proof:** The proof is trivial.



**Theorem 7.20:** Suppose that  $(X, t^*)$  is a supra fuzzy topological space and  $0 \leq \alpha \leq \beta < 1$ , then

- (a)  $(X, t^*)$  is  $\alpha - R_1$ (ii) implies  $(X, t^*)$  is  $\beta - R_1$ (ii).
- (b)  $(X, t^*)$  is  $\beta - R_1$ (iii) implies  $(X, t^*)$  is  $\alpha - R_1$ (iii).
- (c)  $(X, t^*)$  is  $0 - R_1$ (iii) if and only if  $(X, t^*)$  is  $0 - R_1$ (iv).

**Proof:** (a) Suppose that  $(X, t^*)$  is a  $\alpha - R_1$ (ii) space. We have to prove that  $(X, t^*)$  is a  $\beta - R_1$ (ii) space. Let  $x, y \in X, x \neq y$  and  $w \in t^*$  such that  $w(x) \neq w(y)$ . Since  $(X, t^*)$  is  $\alpha - R_1$ (ii), for some  $\alpha \in I_1$ , there exist  $u, v \in t^*$ , such that  $u(x) = 1 = v(y)$  and  $u \cap v \leq \alpha$ . Since  $0 \leq \alpha \leq \beta < 1$ . Thus, we have  $u \cap v \leq \beta$ . Hence  $(X, t^*)$  is a  $\beta - R_1$ (ii) space.

Similarly, we can prove that.

$(X, t^*)$  is  $\beta - R_1$ (iii) space implies  $(X, t^*)$  is  $\alpha - R_1$ (iii) space.

$(X, t^*)$  is  $0 - R_1$ (iii) space if and only if  $(X, t^*)$  is  $0 - R_1$ (iv) space.

**Theorem 7.21:** Let  $(X, t^*)$  be a supra fuzzy topological space, and  $I_\alpha(t^*) = \{ u^{-1}(\alpha, 1) / u \in t^* \}$ , then

- (a)  $(X, t^*)$  is an  $\alpha - R_1$ (ii) implies  $(X, I_\alpha(t^*))$  is a  $R_1$ .
- (b)  $(X, t^*)$  is an  $\alpha - R_1$ (iii) implies  $(X, I_\alpha(t^*))$  is a  $R_1$ .
- (c)  $(X, t^*)$  is an  $\alpha - R_1$ (iv) implies  $(X, I_\alpha(t^*))$  is a  $R_1$ .

**Proof:** (c) Suppose that  $(X, t^*)$  is  $\alpha - R_1$ (iv) space. We have to prove that  $(X, I_\alpha(t^*))$  is a  $R_1$  space. Let  $x, y \in X$ , with  $x \neq y$  and  $M \in I_\alpha(t^*)$  such that  $x \in M, y \notin M$  or  $x \notin M, y \in M$ . Suppose that  $x \in M, y \notin M$ . We can write  $M = w^{-1}(\alpha, 1]$ , where  $w \in t^*$ . Now we have  $w(x) > \alpha, w(y) \leq \alpha$ , i.e.,  $w(x) \neq w(y)$ . Since  $(X, t^*)$  is  $\alpha - R_1$ (iv),  $\alpha \in I_1$ , then there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . It

follows that there exist  $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t^*)$  with  $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$  as  $u(x) > \alpha, v(y) > \alpha$ . and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ , since  $u \cap v \leq \alpha$ . Thus we have,  $(X, t^*)$  is  $R_1$ .

Similarly, (a) and (b) can be proved.

**Theorem 7.22:** Suppose that  $(X, T^*)$  be a supra topological space. Consider the following statements:

- (1)  $(X, T^*)$  be a  $R_1$  space.
- (2)  $(X, w(T^*))$  be an  $\alpha - R_1(i)$  space.
- (3)  $(X, w(T^*))$  be an  $\alpha - R_1(ii)$  space.
- (4)  $(X, w(T^*))$  be an  $\alpha - R_1(iii)$  space.
- (5)  $(X, w(T^*))$  be an  $\alpha - R_1(iv)$  space.

Then

- (a) (1)  $\Rightarrow$  (2)  $\Rightarrow$  (3)  $\Rightarrow$  (5)  $\Rightarrow$  (1)
- (b) (1)  $\Rightarrow$  (2)  $\Rightarrow$  (4)  $\Rightarrow$  (5)  $\Rightarrow$  (1).

**Proof:** First, suppose that  $(X, T^*)$  be a  $R_1$  space. We have to prove that  $(X, w(T^*))$  be a  $\alpha - R_1(i)$  space. Let  $x, y \in X$  with  $x \neq y$  and  $m \in w(T^*)$  such that  $m(x) \neq m(y)$  i.e., either  $m(x) < m(y)$  or  $m(x) > m(y)$ . Suppose that  $m(x) < m(y)$ . Then it is clear that  $m^{-1}(m(x), 1] \in T^*$  as  $m \in w(T^*)$  and  $x \notin m^{-1}(m(x), 1], y \in m^{-1}(m(x), 1]$ . Since  $(X, T^*)$  is  $R_1$  space then there exist  $U, V \in T^*$  such that  $x \in U, y \in V$  and  $U \cap V = \phi$ . Since  $1_U, 1_V$  are lower semi continuous function from  $X$  into  $I$ , then  $1_U, 1_V \in w(T^*)$ , and it is clear that  $1_U(x) = 1, 1_V(y) = 1$  and  $1_U \cap 1_V = 0$ , i.e.,  $(X, w(T^*))$  is  $\alpha - R_1(i)$  space. Hence  $(X, w(T^*))$  is also an  $\alpha - R_1(ii)$  space.

We have already proved (3)  $\Rightarrow$  (5) and (4)  $\Rightarrow$  (5).

Finally, we shall prove that (5)  $\Rightarrow$  (1).

Suppose that  $(X, w(T^*))$  is  $\alpha$ - $R_1$  (iv). We have to prove that  $(X, T^*)$  is  $R_1$  space. Let  $x, y \in X$  with  $x \neq y$  and  $M \in T^*$  such that  $x \in M, y \notin M$  or  $x \notin M, y \in M$ . Suppose that  $x \in M, y \notin M$ . But  $1_M$  is lower semi continuous function from  $X$  into  $I$ , so  $1_M \in w(T^*)$  and  $1_M(x) = 1, 1_M(y) = 0$ , i.e.,  $1_M(x) \neq 1_M(y)$ . Since  $(X, w(T^*))$  is  $\alpha$ - $R_1$  (iii),  $\alpha \in I_1$  then  $\exists u, v \in w(T^*)$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . Now it is observed that  $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in T^*$  such that  $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$  and  $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ . Thus  $(X, T^*)$  is  $R_1$  space.

The proof is now complete.

Thus it is seen that  $\alpha$ - $R_1(p)$  is a good extension of its topological counter part ( $p = i, ii, iii$ ).

**Theorem 7.23:** Let  $(X, t^*)$  be a supra fuzzy topological space,  $A \subseteq X$  and  $t^*_A = \{u/A : u \in t^*\}$ , then

- (a)  $(X, t^*)$  is  $\alpha$ - $R_1$  (i) implies  $(A, t^*_A)$  is  $\alpha$ - $R_1$  (i).
- (b)  $(X, t^*)$  is  $\alpha$ - $R_1$  (ii) implies  $(A, t^*_A)$  is  $\alpha$ - $R_1$  (ii).
- (c)  $(X, t^*)$  is  $\alpha$ - $R_1$  (iii) implies  $(A, t^*_A)$  is  $\alpha$ - $R_1$  (iii).
- (d)  $(X, t^*)$  is  $\alpha$ - $R_1$  (iv) implies  $(A, t^*_A)$  is  $\alpha$ - $R_1$  (iv).

**Proof:** (d) Suppose that  $(X, t^*)$  is  $\alpha$ - $R_1$  (iv). We have to prove that  $(A, t^*_A)$  is  $\alpha$ - $R_1$  (iv). Let  $x, y \in A$  with  $x \neq y$  then  $x, y \in X$  and  $x \neq y$ . Consider  $m \in t^*_A$  with  $m(x) \neq m(y)$ . Then  $m$  can be written as  $w/A$ , where  $w \in t^*$  and hence  $w(x) \neq w(y)$ . Since  $(X, t^*)$  is  $\alpha$ - $R_1$  (iv),  $\alpha \in I_1$ , then there exist  $u, v \in t^*$  such that  $u(x) > \alpha, v(y) > \alpha$  and  $u \cap v \leq \alpha$ . But we know  $u/A \in t^*_A$ , for every  $u \in t^*$ . Now we observed that

$u/A(x) > \alpha$ ,  $v/A(y) > \alpha$  and  $u/A \cap v/A \leq \alpha$ , since  $u \cap v \leq \alpha$ . Thus  $(A, t^*_A)$  is  $\alpha$ - $R_1$ (iv).

Similarly (a), (b) and (c) can be proved.

**Theorem 7.24:** Given  $(X_i, t^*_i)$ ,  $i \in \Lambda$  be supra fuzzy topological spaces and  $X = \prod_{i \in \Lambda} X_i$  and  $t^*$  be the product supra topology on  $X$ , then

- (a)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (i) if and only if  $(X, t^*)$  is  $\alpha$ - $R_1$ (i).
- (b)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (ii) if and only if  $(X, t^*)$  is  $\alpha$ - $R_1$ (ii).
- (c)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (iii) if and only if  $(X, t^*)$  is  $\alpha$ - $R_1$ (iii).
- (d)  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (iv) if and only if  $(X, t^*)$  is  $\alpha$ - $R_1$ (iv).

**Proof:** Suppose that  $\forall i \in \Lambda$ ,  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (iv). We have to prove that  $(X, t^*)$  is  $\alpha$ - $R_1$ (iv). Let  $x, y \in X$  with  $x \neq y$  and  $w \in t^*$  with  $w(x) \neq w(y)$ . But we have  $w(x) = \min \{ w_i(x_i) : i \in \Lambda \}$ ,  $w(y) = \min \{ w_i(y_i) : i \in \Lambda \}$ . Hence we can find at least one  $w_i \in t^*_i$  and  $x_i, y_i \in X_i$ , with  $w_i(x_i) \neq w_i(y_i)$ . Since  $(X_i, t^*_i)$ ,  $i \in \Lambda$  is  $\alpha$ - $R_1$ (iv),  $\alpha \in I_1$ , then there exist  $u_i, v_i \in t^*_i$  such that  $u_i(x_i) > \alpha$ ,  $v_i(y_i) > \alpha$  and  $u_i \cap v_i \leq \alpha$ . But we have  $\pi_i(x) = x_i$  and  $\pi_i(y) = y_i$  and hence  $u_i(\pi_i(x)) > \alpha$ ,  $v_i(\pi_i(y)) > \alpha$ . It follows that there exist  $u_i \circ \pi_i, v_i \circ \pi_i \in t^*$  such that  $(u_i \circ \pi_i)(x) > \alpha$ ,  $(v_i \circ \pi_i)(y) > \alpha$  and  $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) \leq \alpha$ . Thus  $(X, t^*)$  is  $\alpha$ - $R_1$ (iv).

Conversely, suppose that  $(X, t^*)$  is  $\alpha$ - $R_1$ (iii). We have to prove that  $(X_i, t^*_i)$ ,  $i \in \Lambda$ , is  $\alpha$ - $R_1$ (iv). Let for some  $i \in \Lambda$ ,  $a_i$  be a fixed element in  $X_i$ , suppose that  $A_i = \{ x \in X = \prod_{i \in \Lambda} X_i / x_j = a_j \text{ for some } i \neq j \}$ . So that  $A_i$  is a subset of  $X$ , and hence  $(A_i, t^*_{A_i})$  is also a subspace of  $(X, t^*)$ . Since  $(X, t^*)$  is  $\alpha$ - $R_1$ (iv), then  $(A_i, t^*_{A_i})$  is also  $\alpha$ - $R_1$ (iii). Now we have  $A_i$  is homomorphic image of  $X_i$ . Hence  $(X_i, t^*_i)$  is  $\alpha$ - $R_1$ (iv) i.e., (c) is proved.

Similarly, (a), (b) and (c) can be proved.

## BIBLIOGRAPHY

1. Abd EL-Monsef, M.E., and Ramadan, A. E.: On fuzzy supra topological spaces, Indian J. Pure and Appl. Math. 18(4)(1987), 322-329.
2. Ajmal, N.; Tyagi , B. K. : A Characterization of regular fuzzy space; Mat. Vesnik 41(1989) no. 2, 65-70.
3. Ali, D. M. : On Certain Separation and Connectedness Concepts in Fuzzy Topology(PhD Thesis), 1987.
4. Ali, D. M.: A note on fuzzy regularity concepts; Fuzzy sets and systems. 35(1990), 101-104.
5. Ali, D. M.; Wuyts, P.; Srivastava, A. K. : On the  $R_0$  - property in fuzzy topology; Fuzzy sets and systems. 38(1990) no. 1, 97-113.
6. Ali, D. M.: A note on some  $FT_2$  concepts; Fuzzy sets and systems. 42(1991) no. 3, 381-386.
7. Ali, D. M.: A note on  $T_0$  and  $R_0$  fuzzy topological spaces; Pro. Math. Soc. B.H.U. Vol. 3(1987), 165-167.
8. Ali, D. M.: Some Remarks on  $\alpha$ - $T_0$ ,  $\alpha$ - $T_1$ ,  $\alpha$ - $T_2$  fuzzy topological spaces; The Journal of fuzzy Mathematics, Los Angeles, Vol. 1, No. 2(1993), 311-321.
9. Ali, D. M.: Some other types of fuzzy connectedness; Fuzzy sets and systems. 46(1992), 55-61.
10. Ali, D. M.: On the  $R_1$ - property in fuzzy topology; Fuzzy sets and systems. 50(1992), 97-101.
11. Allam , A. A.; Abt. E1, K. M. ; Nehad N. Morsi: On Fuzzy neighbourhood spaces: Fuzzy sets and systems. 41 (2), (1991), 201-212.

12. Arya, S. P.; Nour, T. M.: Separation axioms for bi -topological spaces; Jour. Pure Appl. Math. 19(1), (1988) 42-50.
13. Azad, K. K.: On Fuzzy semi- continuity, Fuzzy almost continuity and Fuzzy weakly continuity; J. Math. Anal. Appl. 82(1) (1981) 14-32.
14. Bennet , G.; Kalton, N. J. : FK-spaces containing co; Duke Math Jour. 39(1972), 561-582.
15. Bhaumik, R. N.; Mukherjee, A.: Some more results on completely induced fuzzy topological spaces, Fuzzy sets and systems. 50 (1992) 113-117.
16. Buck, R. C. : Operator algebras and dual spaces. Proc. Amer. Math. Soc. 3(1952), 681-687.
17. Carmer , J. H. : L – topologische Raume ( un published ) Universitat Bremen (1986) .
18. Chakravarty, K. K.; Ahsanullah, T. M. G. : Fuzzy topology on fuzzy sets and tolerance topology; Fuzzy sets and systems. 45 (1992) 103-108.
19. Chang, C. L.: Fuzzy topological spaces; J. Math. Anal Appl. 24(1968), 182-192.
20. Chattopadhyaya, K. C.; Hazra, R. N.; Samanta, S. K.: Gradation of openness fuzzy topology; Fuzzy sets and systems. 49 (1992) 1237-242.
21. Choubey, A.: Some investigations in Categorical Fuzzy topology, Thesis, Banaras Hindu University, Varanasi, 1995.
22. Cooper, J. B.: The strict topology and space with mixed topologies; Proc. Amer. Math. Soc. 30 (1971), 583-592.
23. Cooper, J. B.: Saks space and applications to Functional Analysis; North Holland Publishers 1978.

- 
- 
24. Cutler, D. R.; Reilly, I. L: A comparison of some Hausdorff notions in fuzzy topological spaces; *Comput. Math. Appl.* 19(1990) no. 11, 97-104.
  25. Das, N. R.; Das, P. : Mixed topological groups. *Indian J. Pure appl. Math.* 22(4), (1991), 323-329.
  26. Das, N. R. ; Baishya, P. C.: Mixed fuzzy topological spaces. *The Journal of Fuzzy Math.* Vol. 3 No. December 4, 1995.
  27. Das, N. R. ; Baishya, P. C. : On open maps, closed maps and fuzzy continuous maps in fuzzy bi-topological spaces. (Communicated)
  28. Devi, R.; Sampathkumar S. and Caldes, M. : *General Mathematics*, Vol. 16, Nr. 2 (2008), 77-84.
  29. Dube, K. K.; Misra, D. N. : Some localized separation axioms and their applications; *M. R.* 59(3) (1979)
  30. Dubois, D.; Prade, H. : *Fuzzy sets and systems. Theory and Applications*; Academic Press New York 1980.
  31. Erceg, M. A. : Functions Equivalence relations, Quotient spaces and subjects in Fuzzy set theory; *Fuzzy sets and systems*, 3(1), (1980) , 75-92.
  32. Eroglue, M. S.: Topological representation for fuzzy topological spaces; *Fuzzy sets and systems.* 42(1991), 335-362.
  33. Evert, J.: On normal fuzzy topological spaces; *Mathematica (cluj)* 31(54) (1989) no. 1, 39-45.
  34. Fath Alla, M. A. ; Abd El . Hakeim, K. M. : Mapping with fuzzy closed graphs and strongly fuzzy closed graphs; *J. Inst. Math. Comput. Sci. Math. Ser* 4(1991) no. 1, 61-67.

- 
- 
35. Fora, Ali Ahmed: Separation axioms, subspace and product spaces in fuzzy topology; Arab Culf J. Sci. Res. 8(1990) no.3, 1-16.
  36. Foster, D. H.: Fuzzy topological groups; J. Math. Anal. Appl. 67(1979), 549-567.
  37. Ganguly, S.; Saha, S. : On separation axioms and separations of connected sets in fuzzy topological spaces.; Bull. Cal. Math. Soc. 79(1987), 215-225.
  38. Ghanim, M. H.; Kerre, E. E.; Mashhour, A. S. : Separation axioms, subspace and sums in fuzzy topology; J. Math. Anal. Appl. 102(1984), 189-202.
  39. Ghanim, M. H.; Morsi, N. N.:  $\alpha$  - axioms in fuzzy topology; Simon Stevin 63(1989) no. 3-4, 193-208.
  40. Ghanim, M. H.: Pseudo- closure operators in fuzzy topological spaces ; Fuzzy sets and systems. 39(1991) no. 3, 339-346.
  41. Goguen, J. A.: L-fuzzy sets; J. Math. Anal. Appl. 18(1967), 145-174.
  42. Goguen, J. A.: The fuzzy Tychonoff Theorem; J. Math. Anal. Appl. 43(1973), 734-742.
  43. Hazra, R. N.; Samanta, S. K.; Chattopadhyaya : Fuzzy topology redefined ; Fuzzy sets and systems. 45(1992) , 79-82.
  44. Hossain, M. S. and Ali. D. M.: On  $T_0$  fuzzy topological spaces, *J. Math and Math. Sci.* Vol. 24(2009), 95-102.
  45. Hossain, M. S. and Ali, D. M.: On  $T_1$  fuzzy topological spaces, *Ganit. J. Bangladesh Math. Soc.* 24(2004), 99-106.
  46. Hossain, M. S. and Ali, D. M.: On  $T_2$  fuzzy topological spaces, *J. Bangladesh Academy of Science*, 29(2), 2005, 201-208.
  47. Hossain, M. S. and Ali, D. M.: On Fuzzy Regular topological spaces, R.U. Studies. Part B. Journal of Science, Vol. 34, 2006, 25-36.



- 
- 
48. Hutton, B.; Reilly, I. L. : Separation axioms in fuzzy topological spaces; Dept. Math. Univ. of Auckland, Report No. 55, March 1974.
  49. Hutton, B. : Normality in fuzzy topological spaces; J. Math. Anal. Appl. (1975), 74-79.
  50. Hutton, B.: Uniformities on fuzzy topological spaces; J. Math. Anal. Appl. 58(1977), 559-571.
  51. Hutton, B.: Products of fuzzy topological spaces Topology; Appl. 11(1980), 59-67.
  52. Katsaras, A. K. : Fuzzy neighborhood structures and fuzzy quasi uniformities; Fuzzy sets and systems. 29(1989) no. 2, 187-199.
  53. Kaufmann, A.: Introduction to the theory of fuzzy subsets. Volume I; Academic Press, I. N. C. (1975).
  54. Kaufmann, A.; Gupta, M. M : Introduction to fuzzy Arithmetic Theory and Application. Van Nostrand Reinhold Company. New York.
  55. Kelley, J. L.: General Topology; Van Nostrand, Princeton, NJ, 1955.
  56. Kerre, E. E.; Ottey, P. L.: On  $\alpha$ -generated fuzzy topologies; Fasc. Math. No. 19 (1990), 127-134.
  57. Lipschutz, S. : General, Topology; Copyright © 1965, by the Schaum publishing Company.
  58. Lowen, R.: Topologies floues; C. R. Acad Sc. Parish 278 series A (1974), 925-928.
  59. Lowen, R. : Initial and final topologies and fuzzy Tychonoff Theorem; J. Math. Anal. Appl. 58(1977), 11-21.
  60. Lowen, R. : Fuzzy topological spaces and fuzzy compactness; J. Math. Anal. Appl. 56(1976), 621-633.

61. Macho, Stadler, M. ; De Prada Vicente M. A. : Strong separation and strong countability in fuzzy topological spaces; Fuzzy sets and systems. 43(1991) no. 1, 95-116.
62. Malghan, S. R.; Benchalli, S. S.: On fuzzy topological spaces; Glasnic Mathematicki, 16(36) (1981), 313-325.
63. Malghan, S. R.; Benchalli, S. S. : On open Maps , closed Maps and Local compactness in fuzzy topological spaces; J. Math. Anal. Appl. 99(2) (1984), 338-349.
64. Mashhour, A.S., Allam, A.A., Mahmoud, F.S. and Khedr, F.H.: On supra topological spaces, Indian J. Pure and Appl. Math. 14(4)(1983), 502-510.
65. Mashhour, A. S.; Morsi, Nehad, N; El. Tantawy, O. A. : Fuzzy neighborhood syntopogenous structure; Fuzzy sets and systems. 43(1991) no. 2, 219-234.
66. Mashhour, A. S. ; Morsi, N, N. : On regularity axioms in fuzzy neighborhood spaces; Fuzzy sets and systems. 44(1991) , 265-271.
67. Md. Fazlul Hoque, Hossain M. S. , Ali D. M.: Some Features of  $\alpha$ - $T_0$  Spaces in Supra Topology, J. Bangladesh Academy of Sciences (Communicated).
68. Md. Fazlul Hoque, Hossain M. S. , Ali D. M.: Some Features of  $\alpha$ - $T_1$  Spaces in Supra Topology, J. Mechanics of Continua and Mathematical Sciences (Accepted).
69. Md. Fazlul Hoque, Hossain M. S. , Ali D. M.: Some Features of  $\alpha$ - $T_2$  Spaces in Supra Topology (Communicated).
70. Md. Fazlul Hoque, Hossain M. S. , Ali D. M.: Some Features of Regular Spaces in Supra Topology (Communicated).
71. Md. Fazlul Hoque, Hossain M. S. , Ali D. M.: Some Features of Normal Spaces in Supra Topology (Communicated).

- 
72. Ming, Pu. Pao ; Ming, Liu Ying : Fuzzy topology I. Neighborhood Structure of a fuzzy point and Moore - Smith convergence; J. Math. Anal. Appl. 76 (1980), 571-599.
  73. Ming, Pu. Pao ; Ming, Liu Ying : Fuzzy topology II. Product and Quotient Spaces; J. Math. Anal. Appl. 77 (1980), 20-37.
  74. Ming Sheng Ying: A new approach for fuzzy topology I; Fuzzy sets and systems. 39(1991), 303-321.
  75. Ming Sheng Ying: A new approach for fuzzy topology II; Fuzzy sets and systems. 49(1992), 221-232.
  76. Morsi, Nehad N.: The Urysohn Lemma for fuzzy neighbourhood spaces; Fuzzy sets and systems. 39(1991) no. 3, 347-360.
  77. Morsi, Nehad N. : Dual fuzzy neighbourhood spaces I; Fuzzy sets and systems. 44(1991) , 245-263.
  78. Mukherjee, M. N.; Sinha, S. P.: On some near fuzzy countinuous functions between fuzzy topological spaces ; Fuzzy sets and systems. 34 (1990) no. 2, 245-254.
  79. Mukherjee, M. N. ; Ghosh, B. : Fuzzy semi- regularzation topologies and Fuzzy submaximal spaces; Fuzzy sets and systems. 49(1991) , 283-294.
  80. Nanda, S.; S. Dang, A. Behera : The fuzzy topological complementation Theorem; Jour. Fuzzy Math, 1(2) (1993), 303-310.
  81. Patronis, T.; Stravrinou, P.: Fuzzy equivalence and the resulting topology; Fuzzy sets and systems. 46(1992), 237-243.
  82. Petricevic , zлата :  $R_0$  and  $R_1$  axioms in fuzzy topology; Mat. Vesnik 41(1989), 21-28.

- 
- 
83. Petricevic , zлата : Some separation axioms in fuzzy spaces; zb, Rad. No. 4(1990), 25 – 33 .
  84. Ramadan, A. A.; Abd Alla , M. A. : On smooth pre-proximity spaces; Fuzzy sets and systems. 48(1992), no 1, 117 – 121 .
  85. Ramadan, A. A.: Smooth topological spaces; Fuzzy sets and systems. 48(1992), 371-375.
  86. Rodabaugh , S. E. ; The Hausdorff Separation axiom for fuzzy topological spaces. Top. Appl (1980), 319 – 334 .
  87. Rudin , W. ; Real and complex analysis . Copyright © 1966, 1974, by McGraw – hill Inc.
  88. Sentilles , F. Dannis : Bounded continuous functions on a completely regular space; Trans . Amer. Math. Soc. Vol. 168(1972).
  89. Shostak, A. P. : Two decades of fuzzy topology : the main ideas , concepts and results. (Russian). Uspekhi. Mat. Nauk 44(1989) 99-147.
  90. Singal, M. K. ; Prakash, N. : Fuzzy pre-open sets and fuzzy pre separation axioms; Fuzzy sets and systems. 44(1991), 273-281.
  91. Singal, M. K.; Rajvanshi, N.: Regularly open sets in fuzzy topological spaces; Fuzzy sets and systems. 50(1992), 343-353.
  92. Sinha, S. P.: Fuzzy normality and some of its weaker forms; Bull. Korean. Math. Soc. 28(1991) no. 1, 89-97.
  93. Sinha, S. P.: Separation axioms in fuzzy topological spaces; Fuzzy sets and systems. 45(1992) , 261-270.
  94. Sostak, A. P.: On the neighborhood structure of fuzzy topological spaces; Zb, Rad. No. 4(1990) , 7-14.

- 
- 
95. Sostak , Alexander; Dzhajanbajev : On completeness and pre-compactness spectra of fuzzy sets in fuzzy uniform spaces; Comment. Math. Univ. Car. 31 (1990) no. 1, 149-157.
  96. Srivastava , R. ; Lal, S. N. ; Srivastava, A. K. : Fuzzy  $T_1$ - topological spaces; J. Math. Anal. Appl.102 (1984), 442-4448.
  97. Srivastava , A. K.; Ali , D. M. : A note on K. K. Azad's fuzzy hausdorffness concepts; Fuzzy sets and Systems. 42 (1991), 363-367.
  98. Srivastava , A. K.; Ali , D. M. : A comparison of some  $FT_2$  concepts; Fuzzy sets and Systems. 23 (1987), 289-294.
  99. Stadler, M. M.; De prada Vicente M. A.: Strong separation and strong countability in fuzzy topological spaces; Fuzzy sets and System. 43 (1991) no. 1, 95-116.
  100. Warner, M. W.; Mclean, R. G.: On compact Hausdorff  $L$ -fuzzy spaces ; Fuzzy Sets and Systems. 56 (1993) , 103-110.
  101. Warren, R. H.: Neighbourhoods, bases and continuity in fuzzy topological spaces; Rocky Mountain J. Math. 8 (1978). 459-470.
  102. Warren, R. H.: Fuzzy topologies characterized by neighborhood systems; Rocky Mountain J. Math. 9 (4) (1997), 761-764.
  103. Weiss, M. D.: Fixed points, separation and induced topologies for fuzzy sets; J. Math . Anal. Appl . 50 (1975), 142-150.
  104. Wong, C. K.: Covering properties of fuzzy topological spaces; J. Math. Anal. Appl. 43 (1973), 697-704.
  105. Wong, C. K.: Fuzzy topology: Product and Quotient Theorem. J. Math. Anal. Appl. 45 (1974) 512-521.

- 
106. Wong, C. K.: Fuzzy points and local properties of Fuzzy topology; J. Math. Anal. Appl. 46 (1974), 316-328.
  107. Wuyts, P., Lowen, R.: On separation axioms in fuzzy topological spaces; Fuzzy neighborhood spaces and Fuzzy Uniform spaces; J. Math. Anal. Appl. 93 (1973), 27-41.
  108. Xu, Luo Shan: The strong Hausdorff Separation property in L- fuzzy topological spaces; Mohu. Xitong . Yu. Shuxue 5 (1991) no. 2, 25-29.
  109. Ying, Ming Sheng: A new approach for fuzzy topology I; Fuzzy Sets and Systems. 39 (1991) no. 3, 303-321.
  110. Zhang, Ghong Quan: Fuzzy continuous function and its properties; Fuzzy Sets and Systems. 43 (1991), 159-171.
  111. Zadeh, L. A.: Fuzzy sets. Information and control 8 (1965), 338-353.

Rajshahi University Library  
Documentation Section  
Document No... D... 3366  
Date... 11.4.12