University of Rajshahi	Rajshahi-6205	Bangladesh.	
RUCL Institutional Repository		http://rulrepository.ru.ac.bd	
Department of Botany		MPhil Thesis	_

1992

Studies on Water Relations in Mustard(Brass/Ca Juncea L.)

Mondal, Ranajit Kumar

University of Rajshahi

http://rulrepository.ru.ac.bd/handle/123456789/949 Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

STUDIES ON WATER RELATIONS IN MUSTARD (BRASSICA JUNCEA L.)



M. PHIL. THESIS

BY

Ranajit Kumar Mondal

B.Sc. (Hons.), M.Sc.

CROP PHYSIOLOGY LABORATORY DEPARTMENT OF BOTANY UNIVERSITY OF RAJSHAHI RAJSHAHI, BANGLADESH

JULY, 1992

STUDIES ON WATER RELATIONS IN MUSTARD(*BRASSICA JUNCEA* L.)

A THESIS

SUBMITTED TO THE UNIVERSITY OF RAJSHAHI IN FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF PHILOSOPHY

ΒY

D-1749

RANAJIT KUMAR MONDAL B.Sc.(Hona.), M.Sc.

JULY, 1992

CROP PHYSIOLOGY LABORATORY DEPARTMENT OF BOTANY UNIVERSITY OF RAJSHAHI RAJSHAHI, BANGLADESH



.

DECLARATION

.

I, Ranajit Kumar Mandal, hereby declare that the whole of the work now submitted as a thesis for the degree of Master of Philosophy to the University of Rajshahi is the result of my own investigation.

٠.

0.7.92

(Professor Nishit Kumar Paul) Countersigned by Supervisor

.

- .

•

Ranajit Kunier Mondal (Ranajit Kumar Mondal)

Candidate

CERTIFICATE

I, Ranajit Kumar Mondal, hereby certify that the work embodied in this thesis has not been submitted in substance of any degree, and is not being concurrently submitted in candidature for any other degree.

Kaul 30.7.92

(Professor Nishit Kumar Paul) Countersigned by Supervisor

Ranajit Kumar Mondal)

Candidate

CONTENTS

.

р	A	ð	P
-	a	ĸ	С.

v

ACKNOWLEDGEMENT	vii
INDEX OF TABLES	ix
INDEX OF FIGURES	xi
ABSTRACT	xv
GENERAL INTRODUCTION	1
REVIEW OF LITERATURE	3
EXPERIMENT 1	19
INTRODUCTION	19
MATERIALS AND METHODS	21
RESULTS	38
Choice of Growth Analysis Technique	38
Total Dry Matter	39
Leaf Area Index	44
Leaf Area Duration	46
. Crop Growth Rate	50
Relative Growth Rate	53
Net Assimilation Rate	53
Leaf Area Ratio	56
Relative Leaf Growth Rate	59
Specific Leaf Area	. 64
Leaf Weight Ratio	64
Correlation Coefficients Between Growth Attributes and Yield	67
Distribution of Dry Matter	75
Relative Leaf Water Content	78

Chlorophyll, Proline and Sugar Contents	78
Relative Leaf Water Content and Time At Which Stomata Closed	81
Transpiration Rates	88
Stomatal Characters	88
Anatomical Characters of Leaves	90
Characters Related to Flowering	92
Flowering Pattern	94
Pod Abortion	94
Pod Growth	99
Seed Growth	99
Yield and Yield Components, Consumptive Water Use and Water Use Efficiency	99
Correlation Coefficients Between Yield and Yield Components	108
DISCUSSION	111
EXPERIMENT 2	132
INTRODUCTION	132
MATERIALS AND METHODS	134
RESULTS	136
Total Dry Matter	136
Distribution of Dry Matter	139
Yield and Yield Components	. 139
Simple Correlation Coefficients Between Seed Yield and Its Components	145
DISCUSSION	149
REFERENCES	154

.

ACKNOWLEDGEMENT

۰,

The author wishes to express his heartiest gratitude and indebtedness to Professor Nishit Kumar Paul, Department of Botany, Rajshahi University, for his valuable advice, active guidance and careful supervision throughout the experimental work and for his helpful criticism and suggestions in preparing the manuscript.

The author thanks to the Chairman, Department of Botany, Rajshahi University, for providing facilities to carry out this work.

The laboratory facilities of Professors O.I. Joarder, M.I. Zuberi, M.M. Hossain, M.S. Rahman and M.A. Khaleque, all from the Department of Botany, Rajshahi University, are gratefully acknowledged. Thanks are due to the Chairman, Department of Bio-Chemistry, Rajshahi University, for providing some chemicals and laboratory facilities.

The author expresses his indebtedness to Mrs. P.B. Kundu, Associate Professor, Department of Botany, Rajshahi University, for her inspiration and help throughout the investigation. For computer facilities, grateful thanks are due to the authority of Computer Centre, Rajshahi University. Thanks are also due to the Meteorological Office, Sampur, Rajshahi, for delivering meteorological data. The author acknowledges the cooperation of Mr. M.R. Khan and Miss. N. Nahar, thesis fellows, Department of Botany, Rajshahi University.

Deep and sincere gratitudes are also due to his parents, brothers and sisters and other well-wishers for their blessings throughout the whole period of the study.

The author is grateful to the University of Rajshahi, Rajshahi, Bangladesh, for the award of a fellowship enabling him to carry out this study.

Finally, the author is thankful to Mr. M. R. Islam for his careful compose of the manuscript.

THE AUTHOR

- Table 1.1. Reduction in mean squares due to successive terms40of two mustard cultivarsas influenced by soilmoisture.
- Table 1.2. Constants obtained from quadratic fitted curves of 43 log_e total dry matter and log_e leaf area of two mustard cultivars as influenced by soil moisture.
- Table 1.3. Mean squares from the analysis of variance of TDM, 48 LAI, LAD and some other growth attributes of two mustard cultivars as influenced by soil moisture (Calculated from quadratic curve-fitted values).
- Table 1.4. Simple correlation coefficients between growth 74 attributes and yield of irrigated and rainfed plants.
- Table 1.5. Mean values of chlorophyll, proline and sugar 82 contents at the flowering and the mid-pod filling stages of two mustard cultivars as influenced by soil moisture.
- Table 1.6. Mean values of relative leaf water content and 87 time at which stomata closed of two mustard cultivars as influenced by soil moisture at the flowering and the mid-pod filling stages.
- Table 1.7. Mean values of transpiration rates of two mustard 89 cultivars as influenced by soil moisture at the flowering and the mid-pod filling stages.
- Table 1.8. Mean values of stomatal characters of two mustard 91 cultivars as influenced by soil moisture.
- Table 1.9. Mean values of some anatomical characters of leaves 92 of two mustard cultivars as influenced by soil moisture.
- Table 1.10.Mean values of some characters at flowering stage 93 of two mustard cultivars as influenced by soil moisture.
- Table 1.11.Mean values of pod abortion on main branch and 93 other branches of two mustard cultivars as influenced by soil moisture.

- Table 1.12.Mean values of yield and yield components and 105 consumptive water use and water use efficiency of two mustard cultivars as influenced by soil moisture.
- Table 1.13.Simple correlation coefficients between yield and 110 yield components of the irrigated and rainfed plants.
- Table 2.1. Mean squares from the analysis of variance of total 137 dry matter (TDM) of two mustard cultivars as influenced by soil moisture stress at specific stages of growth.
- Table 2.2. Mean squares from the analysis of variance of yield 141 and yield components of two mustard cultivars as influenced by soil moisture stress at specific stages of growth.
- Table 2.3. Mean values of yield and yield components of two 146 mustard cultivars as influenced by soil moisture stress at specific stages of growth.
- Table 2.4. Correlation coefficients between seed yield and its 148 components as influenced by soil moisture stress at specific stages of growth.

INDEX OF FIGURES

Figure	0.1.	Standard curve for proline.	28
Figure	0.2.	Standard curve for glucose.	30
Figure	0.3.	Some meteorological parameters during the crop period.(a) Maximum, minimum and soil tempera- ture (⁰ C) and total rainfall (mm), (b) Relative humidity (%) and sunshine hour (h).	36
Figure	1.1.	Effect of soil moisture on total dry matter of two mustard cultivars at different stages of growth.	45
Figure	1.2.	Influence of soil moisture on leaf area index (LAI) of two mustard cultivars at different stages of growth.	47
Figure	1.3.	Effect of soil moisture on leaf area duration (LAD) of two mustard cultivars at different stages of growth.	49
Figure	1.4a.	Influence of soil moisture on crop growth rate (CGR) of two mustard cultivars at different stages of growth (Classical technique).	51
Figure	1.4b	Influence of soil moisture on crop growth rate (CGR) of two mustard cultivars at different stages of growth (Functional technique).	52
Figure	1.5a	Effect of soil moisture on relative growth rate (RGR) of two mustard cultivars at different stages of growth (Classical technique).	54
Figure	1.5b	Effect of soil moisture on relative growth rate (RGR) of two mustard cultivars at different stages of growth (Functional technique).	55
Figure	1.6a	Influence of soil moisture on net assimilation rate(NAR) of two mustard cultivars at different stages of growth (Classical technique).	57
Figure	1.6b	Influence of soil moisture on net assimilation rate(NAR) of two mustard cultivars at different stages of growth (Functional tecnique).	58
Figure	1.7a	.Influence of soil moisture on leaf area ratio	60

Figure 1.7a.Influence of soil moisture on leaf area ratio 60 (LAR) of two mustard cultivars at different stages of growth (Classical technique).

Page

- Figure 1.7b.Influence of soil moisture on leaf area ratio 61 (LAR) of two mustard cultivars at different stages of growth (Functional technique).
- Figure 1.8a. Soil moisture effect on relative leaf growth 62 rate(RLGR) of two mustard cultivars at different stages of growth (Classical technique).
- Figure 1.8b. Soil moisture effect on relative leaf growth 63 rate(RLGR) of two mustard cultivars at different stages of growth (Functional technique).
- Figure 1.9a. Effect of soil moisture on specific leaf area 65 (SLA) of two mustard cultivars at various stages of growth (Classical technique).
- Figure 1.9b. Effect of soil moisture on specific leaf area 66 (SLA) of two mustard cultivars at various stages of growth (Functional technique).
- Figure 1.10a.Effect of soil moisture on leaf weight ratio 68 (LWR) of two mustard cultivars at different stages of growth (Classical technique).
- Figure 1.10b.Effect of soil moisture on leaf weight ratio 69 (LWR) of two mustard cultivars at different stages of growth (Functional technique).
- Figure 1.11. RGR and NAR plotted against log plant dry 70 weight, based on fitted quadratic relations between log (dry weight) and log LA (leaf area) with time.
- Figure 1.12. LAR and RLGR plotted against log plant dry 71 wieght, based on fitted quadratic relations between log W (dry weight) and log LA (leaf area) with time.
- Figure 1.13. SLA and LWR plotted against log plant dry 72 weight, based on fitted quadratic relations between log W (dry weight) and log LA (leaf area) with time.
- Figure 1.14. Influence of soil moisture on distribution of 76 dry matter(DM) of Sambal at successive stages of growth.
- Figure 1.15. Influence of soil moisture on distribution of 77 dry matter(DM) of Daulat at successive stages of growth.

- Figure 1.16a.Relative leaf water content (RLWC) of two 79 mustard cultivars as influenced by soil moisture at the flowering stage.
- Figure 1.16b.Relative leaf water content (RLWC) of two 80 mustard cultivars as influenced by soil moisture at the mid-pod filling stage.
- Figure 1.17a.Effect of soil moisture on relative leaf water 83 content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at the flowering stage of Sambal.
- Figure 1.17b.Effect of soil moisture on relative leaf water 84 content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at the flowering stage of Daulat.
- Figure 1.17c.Effect of soil moisture on relative leaf water 85 content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at the mid-pod filling stage of Sambal.
- Figure 1.17d.Effect of soil moisture on relative leaf water 86 content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at the mid-pod filling stage of Daulat.
- Figure 1.18a.Influence of soil moisture on cumulative number 95 of flowers opened on main axis.
- Figure 1.18b.Influence of soil moisture on cumulative number 96 of flowers opened on primary branch 1.
- Figure 1.18c.Influence of soil moisture on cumulative number 97 of flowers opened on primary branch 2.
- Figure 1.18d.Influence of soil moisture on cumulative number 98 of flowers opened on primary branch 3.
- Figure 1.19. Effect of soil moisture on pod growth of two 100 mustard cultivars at successive stages of growth.
- Figure 1.20. Effect of soil moisture on seed growth of two 101 mustard cultivars at different stages of growth.
- Figure 2.1. Effect of soil moisture stress at specific 138 stages of growth on total dry matter (TDM) of two mustard cultivars.

Figure 2.2. Influence of soil moisture stress at specific 140 stages of growth on distribution of dry matter (DM) of two mustard cultivars.

.

ABSTRACT

Effect of soil moisture on growth, yield and water use of two mustard (*Brassica juncea* L.) cultivars (Sambal and Daulat) was studied. The two moisture levels were irrigation at every 10 days intervals and rainfed control.

Total dry matter (TDM), leaf area index (LAI) and leaf area duration (LAD) were significantly higher in the irrigated plants than that of the rainfed plants. The increase of TDM production was slow at the early vegetative phases but increased rapidly with the advancement of the growth period. Starting from a lower value, LAI and LAD reached a peak and then declined gradually.

Among the growth attributes, relative growth rate (RGR), leaf area ratio (LAR), relative leaf growth rate (RLGR) and leaf weight ratio declined throughout with the increasing time and plant weight. Specific leaf area (SLA) showed no clear pattern but crop growth rate (CGR) increased slightly with fluctuations. Net assimilation rate (NAR) increased at the later stages of growth. Significantly higher CGR, LAR, SLA and LWR were found in the irrigated plants of both the cultivars than that of the rainfed plants.

Simple correlation coefficients indicated that in the irrigated plants, seed yield was positively correlated with the preflowering LAI and CGR and the post-flowering NAR. In the rainfed plants, seed yield was positively correlated with LAI and CGR at the post-flowering stage and negatively correlated with the postflowering NAR and the pre-flowering LAR.

Percentage of dry matter (DM) accumulation for stem and petiole was lower at the early stages, at the middle stages it increased to a peak and then gradually decreased with time. Percentage of DM accumulation in the leaves was found highest at the initial stage and gradully decreased with time and ultimately it became zero.Dry matter partitioning for pods gradually increased with plant age. No clear effect of soil moisture on partitioning of dry matter was found.

At the flowering stage, the irrigated plants had significantly higher relative leaf water content (RLWC)than the rainfed plants at 8 A.M. in both the cultivars. At the mid-pod filling stage, the irrigated plants had significantly higher RLWC at 8 A.M. and 4 P.M. than the rainfed plants of Sambal. The highest RLWC was observed at 8 A.M., but at the midday it decreased and later part of the day it increased gradually. Soil moisture had no significant effect on chlorophyll a,b and total chlorophyll. Proline and sugar contents of leaves were significantly higher in the rainfed plants than that of the irrigated plants at both the flowering and mid-pod filling stages. Accumulation of proline and sugar was higher at the mid-pod filling stage than the flowering stage.

Time taken to stomatal closure and total transpiration rate at the flowering stage were significantly higher in the irrigated plants. The number of epidermal cells was significantly lower and stomatal index was significantly higher in the irrigated plants on the adaxial surface of leaves. Leaf thickness was significantly higher in the irrigated plants than that of the rainfed plants. Mean cross-sectional areas of palisade and spongy parenchyma cells were unaffected by soil moisture.

Plant height at first flowering and duration of flowering were significantly higher in the irrigated plants. Irrigated plants had always higher number of flowers than the rainfed plants. No significant effect of soil moisture was found on pod abortion. Soil moisture had little effect on seed growth of both the cultivars.

Significantly higher plant height, length of main axis, number of pods/plant, number of seeds/pod, pod weight/plant, 1000-seed weight and seed yield were found in the irrigated plants than that of the rainfed plants. In the irrigated condition, Daulat yielded more seeds than Sambal but in the rainfed condition it yielded less, though the differences were not significant. The irrigated plants had significantly higher consumptive water use and lower water use efficiency than the rainfed plants.

Simple correlation coefficients indicated that seed yield in the irrigated plants was positively correlated with the number of pods/plant, pod weight/plant, number of seeds/pod and harvest index, whereas in the rainfed plants it was related positively with plant height, number of pods on main axis, number of pods/ plant, pod weight/plant and harvest index. Oil content of the rainfed seeds was positively correlated with the number of seeds/pod and negatively with plant height and number of pods on main axis.

Effect of soil moisture stress at differnt stages of growth of two mustard cultivars (Sambal and Daulat) was studied. The five treatments were rainfed control (T_0) , pre-flowering stress (T_1) , flowering to mid-pod filling stress (T_2) , post mid-pod filling stress (T_3) and uniformly irrigation throughout the growing season (T_4) . The treatments T_1 , T_2 , T_3 and T_4 of Sambal received 50,70,50 and 80 mm of irrigation water, respectively. The corresponding values of Daulat were 40, 40, 50 and 70 mm. The highest reduction of total dry matter (TDM) was found in continuously stressed (T_0) plants. TDM production was greatly reduced when soil moisture stress was imposed during pre-flowering (T_1) and flowering to pod filling (T_2) stages. It was not affected significantly in post mid-pod filling stress (T_3) . No significant effect of soil moisture stress on partitioning of dry matter was found.

The severest effect was found in continuous soil moisture stress (T_0) for seed yield and most of the yield components. Seed yield and almost all the yield components were highly reduced when moisture stress was imposed during the pre-flowering (T_1) and the flowering to pod filling (T_2) stages. Soil moisture stress at the post mid-pod filling stage had no significant effect on yield and its components.

GENERAL INTRODUCTION

GENERAL INTRODUCTION

Mustard (*Brassica juncea* L.) is a crop of increasing importance in Bangladesh as its utilization for both the edible oil content and the protein content of the meal. *Brassica* oil is also used as lighting and hair oil in rural areas. A small fraction of oil cake is used for manuring. Oil-meed *Brassica* is cultivated in about 3, 18, 000 hectares in Bangladesh (Anon., 1990), which is about 64% of the land planted for oil meds (Kaul and Das, 1986). Mustard is cultivated in about 25% of the total area in which *Brassica* is grown (Rahman, 1978) and the average yield is about 698 kg/ha (Anon., 1990).

Like any other crops, growth and yield of mustard is under the control of environmental factors. In a strict sense mustard is an irrigated crop (Andrews, 1972) since seed yield is greatly increased by the presence of adequate soil moisture (Banerjee *et al.*, 1967; Bhan, 1979; Jain and Jain, 1979; Reddy and Sinha, 1979; Joarder *et al.*, 1979; Bajpai *et al.*, 1981). Oil content and oil yield in mustard are increased with soil moisture treatment (Banerjee *et al.*, 1967; Bhati and Rathor, 1982; Chaniara and Damor, 1982). However, it is widely cultivated in the winter under rainfed conditions in Bangladesh. Average potential evapotranspiration during the growing period (November to February) of mustard was estimated to be 106.33 mm, whereas mean rainfall during that period was 12.28 mm (Manalo, 1975), therefore a moisture stress of 94.05 mm occurred. Thus, erratic and limited rainfall seriously limits production of mustard in Bangladesh.

There are many physiological processes that are modified by water stress and that may contribute to loss of productivity (Thurling, 1974; Richards and Thurling, 1978 a, b, c; Clarke and Simpson, 1978 a, b, c; Reddy and Sinha, 1987). As mustard is grown under areas relaying on moisture stored in the soil to support the crop, an improved understanding of crop responses to water stress would be helpful to manipulate higher yield.

Since rainfall during the winter season is inadequate and uncertain, mustard requires supplemental water for its proper growth and development, otherwise, water stress is likely to develop and reduce crop yields. The yield reduction does not only depend on the magnitude and duration of water stress but also on the stages of growth and development of plants. All stages of growth are not equally sensitive to moisture stress. Hence, it is essential to determine the specific stages of growth to soil moisture stress for proper irrigation scheduling in order to optimize crop production under limited water resources condition. The following two experiments were conducted in the present investigation :

Experiment 1. Effect of soil moisture on morphophysiological characters and yield of mustard (*Brassica juncea* L.).

Experiment 2. Effect of soil moisture stress at specific stages of growth on dry matter production and yield of mustard (*Brassica juncea* L.).



REVIEW OF LITERATURE

Maini, Sandhu and Johal (1964) reported that one post-sowing irrigation gave high response both in seed yield and dry matter production of Toria. While two irrigations did not add much further to crop production. One post-sowing irrigation, as in the case of seed yield, registered marked increases in the growth of the plant while two irrigations appeared to be superfluous except that plant height continued to increase still further, highlighting the influence of water on extension growth. Under such conditions the number of primary and secondary branches suffered a setback possibly on account of development of the individual branch.

Banerjee, Das and Bhattacharjee (1967) studied the irrigation effect on yield and oil content of Toria and its economics. On the basis of seed yield, the differences in number of irrigation were significant. On an average, two irrigations yielded the highest (741 kg/ha) with 49% increased yield over the control (no irrigation). One irrigation stood second (651 kg/ha) with 31% increased yield over the control. There was a fall off in the response with three irrigations and the increased yield was only 19% over the control. The yield of control plot was very poor and yielded on an average 497 kg/ha. There was a slight increase in oil content up to one irrigation. The oil content declined a little with increasing number of irrigations.

Interrelationships between various yield components and yield of yellow sarson under conditions of soil drought were studied by Nathawat, Jain, Bagrecha and Bhargava (1969). A negative correlation between primary and secondary branches was found. The number of pods/plant was positively correlated with primary branches. Primary branches were negatively correlated with yield. The number of pods/plant increased the yield significantly.

Denna (1970) carried out an investigation on transpiration and the waxy bloom in Brassica oleracea. The stomatal and cuticular transpiration rates and quantity of wax per unit area of leaf surface were determined for seven glaucous and non-glaucous sibling lines. There were no statistically significant differences in the stomatal transpiration rates of glaucous and non-glaucous lines, but there were highly statistically significant differences between the two classes of lines in terms of cuticular transpiration. Rubbing the surfaces of leaves to remove lightly adhering wax deposits, such as the waxy bloom, significantly increased the cuticular but not the stomatal transpiration rates of both glaucous and non-glaucous plants. There was no appreciable correlation between the quantity of wax per unit leaf area and the loss of water through cuticular transpiration among either the glaucous or non-glaucous lines. It was concluded that the function of cuticular waxes in limiting cuticular transpiration is a product of the architecture of the quantity of wax per unit area of leaf surface.

The effect of irrigation, plant population and fertilizer was studied with 3 levels of each factor on yield and quality of Indian rape by Wankhede, Ahlawat and Sahni (1970). They observed that under 3 irrigations the growth and fruiting capacity of the plants increased further with 4 irrigations, whereas under 4 irrigations the branches/plant were reduced and the number of capsules/plant was also fewer. Three irrigations were optimum and gave about 6% higher seed yield than 2 and 4 irrigations. Irrigation had no significant effect on oil content. However, with 4 irrigations slightly higher percentage of oil was obtained.

Response of mustard to varying irrigation levels, spacing and fertilizer application was studied by Singh and Tomar (1971). They found that there was definite increase in yield due to application of irrigations over control. During 1969-70, a significant response to irrigation levels was recorded. Highest seed yield was recorded where irrigation was applied at every 40 days which was significantly superior over all others. Second best production was recorded with irrigation given at every 55 days followed by that given every 25 days. Lowest yield was recorded for control plots where no irrigation was given.

Irrigation requirement of raya and effect of irrigation applied at different growth stages in combination with three levels of nitrogen was studied by Mathur and Tomar (1972). Irrigation increased raya yield significantly. Irrigation at the pre-flowering stage was found to be the most beneficial. Two irrigations, first at the pre-flowering (40 to 45 days) and the second at the postflowering (80 to 85 days) stages were recommended. The irrigation requirement of raya crop was to be around 32 cm which may be applied through heavy pre-sowing and two post-sowing irrigations at the aforesaid stages.

Free proline accumulation was measured by Waldren and Teare (1974) in leaves of intact sorghum and soybean grown in growth chambers and subjected to normal drought stress. Stomatal diffusive resistance and leaf water potential were used to determine the degree of stress at the time of proline analysis. Free proline did not accumulate markedly in either species until each was severely stressed, indicating that proline is not a sensitive indicator of drought stress. Free proline accumulated under less stress in soybean than in sorghum. Since soybean is less drought resistant than sorghum, proline accumulation may be indicator of drought resistance or susceptibility.

A field experiment was conducted by Krogman and Hobbs (1975) on yield and morphological responses of rape to irrigation and fertilizer treatments. Rape was grown for 3 years under four levels of irrigation. Irrigation produced more than double yield by promoting greater plant growth, more pods, more seeds/pod and larger seeds than where irrigation was not applied. Maximum yield (about 3000 kg/ha) was obtained where irrigation maintained soil moisture in the upper half of the available range until pod ripening. Oil content was increased by irrigation ; irrigation increased oil yield from 368 to 986 kg/ha.

Blum and Ebercon (1976) explored a possible association

between free proline accumulation in water-stressed leaves and the drought resistance of various grain sorghum cultivars. Eight sorghum cultivars were grown in a growth chamber under a drying cycle and a subsequent recovery phase. Free proline, free ammonia, dark respiration rates and recovery rating were recorded, as well as desiccation tolerance of each cultivar when grown in the field. Proline accumulation was initiated in leaves of all cultivars as leaf water potential was reduced to -14 to -16 bars. Upon watering, free proline content was reduced abruptly. Cultivars differed significantly in maximum free proline accumulation. Free proline accumulation during water stress was correlated significantly with post-stress recovery rating. It was suggested that accumulated free proline in water-stressed sorghum leaves was related to the ability of cultivar to recover upon the relief of stress, possibly by way of proline's role as a source of respiratory energy in the recovering plant.

Clarke and Simpson (1978a) carried out an experiment to study the growth analysis of *Brassica napus* cv. Tower. The experiment was conducted in the field for two years under conditions with four planting densities and three water regimes. They noted that nonirrigated conditions caused a greater proportion of dry matter production to occur before flowering than after flowering, while the reverse occurred under irrigated conditions. Irrigation increased leaf area index (LAI). Maximum LAI occurred near the start of flowering and then declined rapidly. Maximum LAI was correlated with seed yield. Mean net assimilation rate (NAR) and mean crop growth rate (CGR) were influenced by irrigation. There was an increase in NAR during the ripening phase, suggesting increased photosynthetic efficiency. No evidence of a distinct optimum leaf area index was found.

Clarke and Simpson (1978b) studied the effect of irrigation and seeding rates on yield components of *Brassica napus* cv. Tower. The experiment was carried out in the field for two years under conditions of four planting densities and three water regimes. The number of branches/plant was scarcely affected by irrigation. The number of pods/plant was incressed by increased irrigation. The number of seeds/pod and 1000-seed weight also increased by irrigation. Yield was positively correlated with 1000-seed weight in both years. Yield component compensation was evident in the relationship of 1000-seed weight to pod number and number of seeds/pod.

Clarke and Simpson (1978c) reported that leaf osmotic potential was an indicator of crop water deficit and irrigation need in rapeseed. Leaf osmotic potential (L ψ_0) of field-grown rapeseed was measured under rainfed conditions and at two levels of irrigation. Seed yield of irrigated *B. napus* was substantially higher than that of the rainfed crop in both years. In 1975, the increase in yield of the low and high irrigation treatments were 89 and 213%, respectively. The corresponding increases for 1976 were 59 and 128%. Within year, seed yields of the three water regimes were significatly different. Henry and MacDonald (1978) conducted an experiment to study the effect of soil and fertilizer nitrogen and moisture stress on yield, oil and protein contents of rape. They observed that yield and oil content increased with irrigation treatment.

Richards (1978) studied the genetic variatin in physiological and physicochemical parameters associated with drought resistance between cultivars in *Brassica napus* and *B. campestris*. Significant variation in proline accumulation, chlorophyll stability, germination rate and percentage, relative turgidity, growth rates and water use efficiency were found in plants grown under droughted conditions.

Richards and Thurling (1978a) investigated the variation between and within species of rape seed (Brassica campestris and B. napus) in response to drought stress. They observed a highly significant variation between cultivars in B. campestris for every parameters except straw weight. In B. napus, the yield components, pods/plant, pods/main axis, branches/plant and seeds/pod were all Thousand-seed weight significantly reduced by drought. was significantly affected by drought only in the late-flowering cultivars. With the exception of 1000-seed weight, yield component means in the treatment imposed from the commencement of pod-filling were significantly higher than those in the stem elongation or flowering treatments. Differences between means of the two latter treatments were not significant, which suggested that the plants

were more sensitive to drought at flowering than at stem elongation.

Bhan (1979) studied the effect of soil moisture and nitrogen on mustard under Gangetic alluvium of Uttar Pradesh. Two varieties viz., Varuna and Appressed mutant, 4 soil moisture regimes and 4 levels of nitrogen showed profound variation in different growth and yield attributes. Varuna was found significantly superior to Appressed mutant in yield and yield contributing characters like siliquae/plant, length of siliqua and 1000-seed weight. Seed yield obtained with irrigation 75% available soil moisture depletion (ASMD) was found at par with irrigation after 60% ASMD and significantly superior to irrigation after 90% ASMD and unirrigated (control) plots. Plant height, secondary branches, siliquae/plant and 1000-seed weight were also improved significantly due to irrigation. The interactions of variety and irrigation showed a significant effect on seed yield indicating great responsiveness of Varuna to irrigation.

The response of Indian mustard to levels of irrigation and fertility in light-textured soil of Rajsthan was studied by Jain and Jain (1979). They observed an increase in yield with an increase in the irrigation over minimum level, but the differences were not significant. The overall average yield of 1893 kg/ha was with 3 irrigations. A further increase in irrigation gave an additional yield of only 100 kg/ha.

Joarder, Paul and Ghose (1979) conducted an investigation on the effect of irrigation and fertilizer on mustard. Three cultivars of mustard were grown on field plots with five levels of fertilizers and with or without irrigation. Irrigation gave 59% more seed yield than no irrigation. Higher seed yield in the irrigated plots were accompanied by increased pods/plant, seeds/pod and number of primary and secondary branches. The most significant increase (83%) was noted in the number of seeds/pod produced by late flowers under irrigation, whereas the increase for earlyformed pods was only 6%. Increase in more than 20% for primary and secondary branches and pods/plant were also obtained with irrigation, but there were no significant differences in 1000-seed weight of early and late-flowered pods, though the seeds of irrigated plots were significantly heavier than the non-irrigated ones. Under irrigated conditions plants produced more primary and secondary branches, more late flowers produced seed-bearing pods and seeds grew slightly larger than without irrigation.

Bhan, Balaraju and Vidya Ram (1980) carried out an experiment on water use, yield and quality of rape as influenced by spacing, irrigation and time of harvest. They found that irrigation had a pronounced effect on the seed yield and the characters contributing to it, like siliquae/plant, seeds/siliqua and 1000-seed weight. The maximum yield was obtained with two irrigations administered at flower initiation and pod development. With one irrigation, better result was obtained when it was given at flower initiation than pod development. The yield was stepped up to 9.8 q/ha with irrigation

Rajshahi University Library Document N. D-2644 Document N. D-2644 Date 7: 81 94 D-1744 at flower initiation from a mere 5.13 g/ha unirrigated control. The oil content in the seed was also highest (40.4%) with this treatment, which gave a difference of 2% over the unirrigated plot.

Bajpai, Singh and Chhipa (1981) studied the influence of sowing date and irrigation levels on growth and yield of mustard. The field study was conducted on 3 varieties, four sowing dates and three irrigation levels. The effect of moisture regimes on the yield was not found significant. But the plant height and number of pods/plant was influenced significantly due to moisture regimes and the maximum height and number of pods were recorded at irrigation water/cumulative pan evaporation (IW/CPE) ratio of 0.5.

Bhati and Rathor (1982) studied the response of Indian mustard to irrigation and nitrogen fertilizer. The treatment consisted of 3 levels of irrigation viz., no irrigation (I_{0}), one irrigation at flowering stage (I_1) and two irrigations, one at flowering and the other at seed setting stage (I_2). The experiment showed that irrigation treatment did not affect the seed yield, oil content and oil yield significantly yet a consistent increasing trend was observed. Except the number of siliquae/plant and stover yield, all other yield attributting characters also showed nonsignificant change due to irrigation. Studies on moisture depletion pattern from different soil depths and consumptive use indicate that the amount of water depleted from 0-15, 15-30 and 30-45 cm soil depth was 7.52, 5.43, 5.12; 10.01, 7.24, 6.60 and 11.64, 8.15, 7.09 cm in I_0 , I_1 and I_2 treatments, respectively. The total consumptive water use thus obtained with different levels of irrigation was of the order of 18.07, 23.85 and 26.88 cm under I_0 , I_1 and I_2 treatmetns, respectively.

Chaniara and Damor (1982) carried out a field experiment to study the effect of irrigation intervals and various levels of nitrogen and phosphorus on the yield of mustard variety Varuna. The treatment consisted of 3 levels of irrigation, 3 levels of nitrogen and two levels of phosphorus. The experiment indicated that the differences due to irrigation interval were significant in respect of number of siliquae/plant, weight of seeds/plant, seed yield and oil yield/ha. The average seed yield realised by 3, 5 and 7 irrigations during the crop growth period at 35, 25 and 15 days interval was 1686, 1893 and 1974 kg/ha, respectively. Maximum seed yield was recorded by giving 5 irrigations at 25 days interval. This may be attributed to higher number and weight of seeds/plant.

Singh (1983) investigated the response of mustard and chickpea to soil moisture in soil profile and plant population on aridisols. He obseved that the mean seed yield was 19.53 q/ha with high soil moisture and 13.66 q/ha with low soil moisture, the increase being 43% with soil moisture. An additional supply of 1 cm/1 m water in the soil profile gave 110.7 kg/ha greater mean seed yield of mustard. Seed yield of chickpea increased 23.3% with high soil moisture over low moisture. Harvest index in general increased with the supply of moisture. In chickpea, high moisture gave 21.9% more straw yield. The harvest index in chickpea increased with an increase in soil profile moisture.

Khan and Agarwal (1985) worked on the effect of sowing methods, moisture stress and nitrogen levels on growth, yield components and seed yield of mustard. When compared with moisture stress on IW/CPE ratio of 0.4 enhanced all the growth and yield characters like plant height, branches/plant, dry matter, siliquae/plant, seeds/siliqua, 1000-seed weight and seed yield and a ratio of 0.6 further increased plant height and dry matter.

Mandal, Ray and Dasgupta (1986) made an investigation on the effect of water used by wheat, chickpea and mustard grown as sole crop and intercrop. They reported that with increase in the number of irrigations, there was an increase in the accumulation of dry matter and leaf area index (LAI). The grain yield of wheat showed an increasing trend with the increase in the level of irrigation. In chickpea, the differences between 2, 3 and 4 irrigations were not very high. Likewise, the difference between 3 and 4 irrigations was less in mustard.

Haque, Ahmed and Rahman (1987) worked on irrigation scheduling for optimum yield of M-248 (Brassica juncea) and M-12 (Brassica campestris). Crops were grown in the field and in the lysimeter (M-248 only) to investigate the effect of irrigation at particular stages of growth on the yield and water use efficiency. Significant increase in yield was found for most of the irrigation treatments compared with no irrigation. The yield increase was highly significant for two irrigations, one at the early vegetative stage and the other at the initial pod formation stage. The field and lysimetric results were almost identical in nature. The seed yield of 1103 and 1204 kg/ha was obtained for two irrigations in M-12 and M-248, respectively and the correspoding values for no irrigation were 269 and 416 kg/ha.

Reddy and Sinha (1987) studied the effect of irrigation and NP fertilization on the yield of Indian mustard. They observed that irrigation at 0.6 IW/CPE and 0.3 IW/CPE produced 17.9 and 16.4 q/ha while the unirrigated treatment gave 15 q/ha seed. Irrigation at o.6 IW/CPE produced significantly higher yield over no irrigation, but the difference in seed yield between 0.3 IW/CPE and 0.6 IW/CPE was not significant. Similarly, the difference in yield between 0.3 IW/CPE over no irrigation treatment was also not significant. It was the combined effect of increased number of siliquae/plant and number of seeds/siliqua which resulted in higher seed yield under irrigation. One irrigation at 0.3 IW/CPE, which coincided with was found critical for moisture. siliqua formation stage, Irrigation at this stage mainly increased the siliqua size and seed size boosting total seed yield.

A field experiment was conducted by Singh, Gupta and Moinbasha (1987) to study the effect of moisture needs of Indian mustard-chickpea intercropping. They observed that with 1 and 2 irrigations, the grain yield of mustard increased by 4.19 and 4.93
q/ha over no irrigation in the first year and 3.41 and 4.46 q/ha in the second year. Further, the mustard yield with 2 irrigations was not significantly better than 1 irrigation. The plants utilized the maximum amount of water (213 and 230 mm) when they were irrigated twice instead of once (200.9 and 211.3 mm). Two irrigations caused 15.4 and 52.4 mm (mean) more moisture use than 1 irrigation and no irrigation, respectively. Moisture use efficiency increased with 1 irrigation by 0.18 kg/mm/ha over 2 irrigations and 0.57 kg/mm/ha over no irrigation. Thus, 1 irrigation produced 8.61 kg grain/mm/ha as compared with 8.43 and 8.04 kg grain/mm/ha with 2 irrigations and no irrigation, respectively.

Reddy, Sinha and Hedge (1988) conducted field experiments during rabi seasons of 1983-84 and 1984-85 to study the effect of irrigation, nitrogen and phosphorus fertilization on consumptive water use, yield, water use efficiency and moisture extraction by mustard. Irrigation at 0.6 IW/CPE produced significantly higher seed yield (21.6 g/ha) over no irrigation (18.3 g/ha), but the differences between the treatments 0.3 IW/CPE and 0.6 IW/CPE or 0.3 IW/CPE and no irrigation were not significant. Irrigation increased the number and size of the siliqua and seed yield per plant.

Sarkar, Ahsan, Sarker and Hossain (1989) studied the effect of nitrogen and irrigation at different stages of growth of mustard variety Sambal. There were four levels of irrigation : at (a) vegetative (25 DAS), (b) flowering (45 DAS), (c) siligua-filling stage (65 DAS) and (d) rainfed control. They observed that the plant height was increased due to irrigation but the same at different stages of crop growth did not influence the height significantly. The plants irrigated at flowering stage produced the highest yield of seed. The stover yield was not significantly influenced by the irrigation at different stages of growth. The number of filled-siliquae/plant, the number of seeds/siliqua and 1000-seed weight increased due to irrigation at various stages of growth.

Sharma and Kumar (1989) conducted a field experiment to study the effect of irrigation on growth analysis, yield and water use in Indian mustard. They found that LAI, CGR, RGR, NAR and RLGR were significantly higher under 0.6 irrigation depth/cumulative pan evaporation ratio than under rainfed codition. An increase in the frequency of irrigation gave more branches/plant, siliquae/plant, seeds/siliqua, 1000-seed weight and seed yield in both the years. The moisture use from the soil decreased by 53.3 and 75.8% under 0.4 and 0.6 irrigation depth/cumulative pan evaporation ratio compared with rainfed condition.

A field experiment was conducted by Raja and Bishnoi (1990) to study root characters, evapotranspiration, water use efficiency, moisture extraction pattern and plant water relations of rape genotypes under varying irrigation schedules. They observed that increased irrigation frequency increased evapotranspiration but decreased water use efficiency. The relative water content and osmotic potential of the leaves increased with more frequent irrigation, but plant water retention capacity decreased.

A field experiment was carried out by Sarker, Ahmed, Talukdar and Khan (1990) to study the influence of irrigation on the yield and water use of mustard mutant BINA-3. Irrigation treatments showed positive effect on the increase of yield of the mutant. The highest yield of 1.15 t/ha (17% higher over control) was obtained with one irrigation at the pre-flowering stage. Crop water use efficiency decreased with the increase of irrigation frequency. The results thus revealed that one irrigation at the pre-flowering stage (35-40 DAS) was sufficient for obtaining the optimum yield of the mustard mutant.

EXPERIMENT 1

EFFECT OF SOIL MOISTURE ON MORPHO-PHYSIOLOGICAL CHARACTERS AND YIELD OF MUSTARD(BRASSICA JUNCEA L.)

INTRODUCTION

Seed yield of mustard is increased by the presence of adequate soil moisture (Mathur and Tomar, 1972; Bajpai *et al.*, 1981; Bhati and Rathor, 1982; Mandal *et al.*, 1986; Reddy and Sinha, 1987). Water stress leads direct changes in the physical environment of the crops, and these changes may subsequently affect crop physiology. As the soil dries, the water potential decreases and so does the soil hydraulic conductivity. Thus it is more difficult for plants to extract water and, as a consequence the plant water potential tends to decrease (Gardner, 1960).

Water deficit in plants generally leads to reduced leaf water potential and stomatal closure, as manifested from an increased leaf resistance to transpiration. The effect of depletion and replenishment of soil water on transpiration are of specific importance to water use and its efficiency in crop production. The relative rates of absorption and transpiration determine plant's internal water balance, which directly affects the physiological and biochemical processes of plant growth (Teare and Kanemasu, 1972). Shortage of water in soil may cause a decrease in nutrient supply. Mobility of ions in the soil decreases as the soil dries and the ability of roots to take up some ions may also decrease (Dunhum and Nye, 1976). The consequent decrease in nutrient uptake may affect a wide range of physiological processes.

Variation in dry matter accumulation and yield influenced by

soil moisture may be related to factors, such as leaf area index (LAI), crop growth rate (CGR), net assimilation rate (NAR) and leaf area ratio (LAR).

Crop plants accumulate free proline in their leaves when exposed to moderate to severe water stress (Palfi et al., 1973). The physiological significance of this metabolic response to water is, however, contentious. Free sugars are also accumulated in the stressed plants (Stewart, 1971; Narasimha Rao and Shiv Raj, 1985). It is suggested that proline and sugar accumulating potential could serve as an index of drought resistance.

Stomata are recognized as the main site of transfer of water vapour and Co_2 between a plant and its environment (Milthorpe, 1969). There are few reports which indicate that stomatal frequency of leaves is affected by soil moisture (Ciha and Brun, 1975; Nerkar *et al.*, 1981).

The main objective of the present experiment was to study the effect of soil moisture on crop growth, development, dry matter production and partitioning, crop water use and yield of mustard.

MATERAIALS AND METHODS

Two mustard (*Brassica jencea* L.) cultivars viz., Sambal and Daulat were used as the plant material for the experiment.

The experiment was carried out in the Botanical Research Field of Rajshahi University. The soil of the experimental field was silty loam having a pH 7.5, low in organic carbon (0.44%), total N (0.43%) available P (15 ppm) and K (82 ppm). The field capacity of the soil was 35%. The field was prepared after repeated ploughing. Before sowing, a basal dose of nitrogen (60 kg/ha), phosphate (30kg/ha) and potash (30 kg/ha) were added. The experiment was arranged in a split plot design with three replications. After preparation of the field it was divided into two main plots. The main plots were treated as treatment plots. Each main plot was three replications. Each replication again subdivided into comprised 8 rows of 4 m long and 30 cm apart. Six rows for each cultivar were taken as experimental rows and two bordering rows were considered as non-experimental. The distance between plants within a row was approximatly 10 cm. Seeds were sown on I November, 1990. Two levels of irrigation treatment were adopted, viz., (i) irrigation at 10 days interval throughout the growing period (irrigated) and (ii) no irrigation (rainfed). On each occasion 10 mm of water was applied to the irrigated field with sprinklers. The irrigated Sambal received 80 mm and Daulat received 70 mm of irrigation water.

Twelve days after sowing (DAS), the seedlings were thinned to uniform and desirable number of plants. Necessary cultural practices, such as weeding and spraying of insecticides were done for normal growth and development of the plants.

Growth Analysis

Eleven harvests were taken at equal intervals of 7 days for growth analysis. Four plants were selected for each cultivar from each treatment for each harvest. The first harvest was taken at 20 DAS.At each harvest, the plants were cut just below the cotyledonary node and tops were separated into leaves, stem + petiole, flowers + buds and pods (when present). Before weighing, the plant parts were dried separately in an oven at about 85 $^{\circ}$ C for 24 hours till they reached constant weight.

Leaf area was measured by the disc method. Twenty discs were used in calculating the area. The area of the cork borer used for taking the discs was 0.5675 cm^2 . Discs were weighed after ovendrying and leaf area was calculated by using following formula:

Area of dics x Weight of leaves Weight of discs

Both the harvest interval method (classical technique) and the curve-fitting method (functional technique) were followed to

determine different growth attributes. From the dry weight of different plant parts and leaf area data, the following growth attributes were calculated between two successive harvests by following the technique of growth analysis (Radford, 1967) :

1. Crop growth rate (CGR) =
$$\frac{W_2 - W_1}{t_2 - t_1}$$
2. Relative growth rate (RGR) =
$$\frac{\log_e W_2 - \log_e W_1}{t_2 - t_1}$$
3. Net assimilation rate (NAR) =
$$\frac{(W_2 - W_1) (\log_e LA_2 - \log_e LA_1)}{(LA_2 - LA_1) (t_2 - t_1)}$$
4. Leaf area ratio (LAR) =
$$\frac{(\log_e W_2 - \log_e W_1) (LA_2 - LA_1)}{(W_2 - W_1) (\log_e LA_2 - \log_e LA_1)}$$
5. Relative leaf growth rate (RLGR) =
$$\frac{\log_e LA_2 - \log_e LA_1}{t_2 - t_1}$$
6. Leaf area duration (LAD) =
$$\frac{(LA_1 + LA_2)}{2} (t_2 - t_1)$$

The following attributes were calculated separately for each harvest :

7. Leaf area index (LAI) =
$$\frac{\text{Leaf area}}{\text{Ground area}}$$
8. Specific leaf area (SLA)=
$$\frac{\text{Leaf area}}{\text{Leaf dry weight}}$$
9. Leaf weight ratio (LWR) =
$$\frac{\text{Leaf dry weight}}{\text{Total plant dry weight}}$$

Where, W_2 and W_1 are the tatal dry weights, LA_2 and LA_1 are the

leaf area per plant at t_1 and t_1 , the latter and the former harvest, respectively.

In the curve-fitting method, polynomial functions were fitted to natural logarithmic values of total dry weight and total leaf area, using a scientific calculator. The log_e transformation was made in order to render the variance homogeneous with time (Hughes and Freeman, 1967). The selection of appropriate polynomial regression model was made by 'lack of fit' method of Nicholls and Calder (1973). The second degree polynomial curves have been used in all respects. The second degree formulae for growth attributes in the functional technique used were as follows :

1. RGR = b + 2 ct 2. LAR = Antilog_e ($\log_e LA - \log_e W$) 3. NAR = RGR/LAR 4. RLGR = b[/] + 2 c[/]t

Where, b and c are constants of dry weight and b[/] and c[/] are constants of leaf area per plant obtained from polynomial equations and t is the instanteneous time of harvest. From similar curves fitted to \log_e total dry weights, \log_e leaf weights and \log_e leaf area, it was possible to derive values for specific leaf area (SLA) and leaf weight ratio (LWR).

Relative Leaf Water Content (RLWC)

Relative leaf water content (RLWC) was determined from the fully matured 3rd or 4th leaf. The leaves were collected at 8 A.M., 12 Noon and 4 P.M. and 20 discs were taken from leaves of each replication of each cultivar of each treatment. Their fresh weights were taken immediately and were sinked into water kept in petri dish for 3 hours. After 3 hours when the cells of the discs became fully turgid they were taken off from water and after drying with blotting paper their turgid weights were determined. Then the discs were dried in an oven and weighed. The relative leaf weter content was calculated from the following formula (Weatherly, 1950):

RLWC was determined at both the flowering and the mid-pod filling stages.

Chlorophyll Content

Chlorophyll contents were estimated at both the flowering and the mid-pod filling stages. Three discs, each of 0.5675 cm² area were taken from three different positions of the 4th leaf. Chlorophyll was extracted with 80% aquous acetone using a morter and pestle to grind the tissues. The suspension was decanted into centrifuge tubes and centrifuged it for 3 minutes. The clear green solution was then decanted from the colourless residue and made up to 10 ml with 80% acetone. The optical density (O.D.) of this solution was determined against 80% acetone as blank using a Spectrophotometer at 645 and 663 nm. The chlorophyll a and b were determined according to the formulae given by Mackinney (1941) and later used by Arnon (1949) as follows :

Chlorophyll a = 12.717 $A_2 - 2.584 A_1 = mg$ Chl. a per litre. Chlorophyll b = 22.869 $A_1 - 4.670 A_2 = mg$ Chl. b per litre.

Where A_1 and A_2 are O.D. at wavelengths of 645 and 663 nm, respectively. Amount of chlorophyll per unit of area was calculated in the following way :

mg Chl. a or b per litre x 10

1000 x leaf area

Estimation of Proline

Proline estimation of the leaves was done according to Bates et al. (1973). For estimation of proline the reagents required were (1) 3% aquous sulphosalicylic acid, (2) Glacial acetic acid, (3) Toluene, (4) Proline (AR) and (5) Acid ninhydrin reagent [6M orthophosphoric acid (20 ml) + glacial acetic acid (30 ml) + ninhydrin (1.25 g)].

At first 0.5 g fresh leaf was homogenized in 10 ml of 3% aquous sulphosalicylic acid and centrifuged it for 6 minutes. The clear solution was separated and 2 ml of it was reacted with 2 ml of acid ninhydrin and 2 ml of glacial acetic acid in a test tube. Then it was boiled in a boiling water bath for 1 hour and the reaction was terminated in an ice bath. The reaction mixture was then extracted with 4 ml of toluene, mixed vigorously with stirring for 15-20 seconds. The chromophore containing proline - toluene was separated with a separating funnel and warmed to room temperature. The optical density (O.D.) at 520 nm was read using toluene as blank. Proline content was determined from the standard curve (Figure 0.1). Proline (AR) was used for the preparation of standard curve.Proline content of leaves was determined at the flowering and mid-pod filling stages.

Stock solution of 1 millimole was prepared by dissolving 0.1151 g of proline (AR)in distilled water and made up to 1000 ml.By successive dilution 2 ml of solution containing 0.0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 and 0.9 micromole concentrations were prepared and reacted with acid ninhydrin and glacial acetic acid as described earlier. The colour was read at 520 nm and the 0.D. was plotted against concentration (Figure 0.1).

Estimation of Sugar

Estimation of total free sugar of the leaves was done according to the anthrone method. For estimation of sugar, the reagents required were (1) 0.2% anthrone reagent (200 mg of anthrone in 100ml of conc. H_2So_i), (2) Ethyl alcohol and (3) Standard glucose solution (10 mg of glucose in 100 ml of distilled water).



At first 1 g of dry leaf was homogenized with distilled water. The mixture was filtered with a double-layered muslin cloth. The filtrate was diluted with twice volume of ethyl alcohol and was kept in a cool place overnight. In the next morning, the mixure was centrifuged and the clear solution was separated. The volume of the extract was evaporated to about one-fourth of the volume over a water bath and cooled. This reduced the volume of the extract and was then transferred to a 100 ml volumetric flask and made up to the mark with distilled water.

One ml of the distilled extract from each sample was pipetted into test tubes. Four ml of the anthrone reagent was added to each of this solution and mixed well.Glass marbles were placed on top of each tube to prevent evaporation. The tubes were placed in boiling water bath for 10 minutes, then removed and cooled. A reagent blank was prepared by 1 ml of water and 4 ml of reagent in a test tube and treated similarly. The optical density (O.D.) of the blue green solution was measured at 620 nm using a Spectrophotometer. Sugar content was determined from the standard curve (Figure 0.2). Sugar content was determined at both the flowering and mid-pod filling stages.

The standard curve of glucose was prepared by taking 0.0, 0.1, 0.2, 0.4, 0.6, 0.8 and 1 ml of standard glucose solution in different test tubes containng 0.0, 10, 20, 40, 60, 80 and 100 μ g of glucose, respectively and made the volume up to 1 ml with



Figure 0.2. Standard curve for glucose.

distilled water. Four ml of anthrone reagent was added to each test tube and mixed well. All these solutions were treated similarly as described above. The colour was read at 620 nm and the O.D. was plotted against concentration (Figure 0.2).

The Relation Between Leaf Water Content and Transpiration Rate

Hygen's (1951) quick weighing method for the analysis of transpiration of detached leaves was used for the comparison of the relationship between leaf water status and stomatal behaviour.Leaf samples were collected on sunny morning in polythene bags. The cut end of the petiole of the leaves was kept into water in beakers under bell jars in a humid condition for about 3 hours. Leaves were then rapidly dried with filter paper and weighed. The water loss was followed for 3 hours by recording fresh weight(FW) at intervals of 10 minutes for the first 90 minutes and then at intervals of 30 minutes. Between weighings, the leaves were kept in semi-controlled environment (light intensity 2.37 klux, temperature 25 °C and R.H. 50%) with their adaxial surface uppermost. The leaves were drawn on chromatographic paper to determine leaf area and the dry weight (DW) of the leaves was taken at the end of the experiment. The initial weight was considered as the turgid weight (TW). Relative leaf water content (RLWC) at any point during a drying cycle was calculated from the following formula :

$$FW - DW$$

$$RLWC = ----- x 100$$

$$TW - DW$$

The curve is divisible into 3 phases. The first and final linear phases when the stomata were opened (stomatal loss) and closed (cuticular loss), and a curvilinear phase during the time of stomatal closure. The point of intersection of the two linear portions of the curve has been taken as indicating the relative water content at stomatal closure (Jarvis and Jarvis, 1963). Stomatal and cuticular water loss have been calculated from the transpiration decline curves. These were determined at both the flowering and mid-pod filling stages.

Stomatal Characters and Leaf Anatomy

The number of stomata and the number of epidermal cells were counted and stomatal pore length was measured on the adaxial and abaxial surfaces by applying Quickfix (Wembley Enterprises, India) on the leaf segments. The number of stomata and epidermal cells of five random focuses was counted under a compound microscope with 15 x 40 magnification and subsequently converted to the number per mm^2 of leaf area. The stomatal pore lengths were measured and converted to μm . From the number of stomata and epidermal cells stomatal index was also calculated according to the following formula :

For anatomical study, leaf segments were fixed in acetic-alcohol fixative (3 parts of 70% ethyl alcohol and 1 part

glacial acetic acid) for about twelve days and then transferred to 70% ethyl alcohol. Transverse sections of the colourless segments were cut by a blade and were projected by a camera lucida and outline of the cells of the leaf tissues was drawn on paper. From the outline, leaf thickness was determined and was converted into μ m. The outline of five random cells of palisade parenchyma and five spongy parenchyma were also drawn on chromatographic paper and the cross-sectional area was determined and then converted into μ m².

Characters Related to Flowering

Days taken to first flowering were noted. Plant height and the number of leaves at first flowering were also recorded.Duration of flowering i.e., days taken from first flowering to last flowering was also noted.

Flowering Pattern and Pod Abortion

The cumulative number of flowers opened on the main axis and primary branches 1-3 was counted on alternate days. Pod abortion was also calculated on the main axis and primary branches 1-3 according to the following formula :

Pod abortion (%) =

Number of flowers opened finally - Number of pods matured Number of flowers opened finally

Pod and Seed Growth

For determining pod and seed growth, 8 harvests were taken at equal intervals of 5 days. The first harvest was taken at 10 days after flowering (DAF). Twenty pods were taken from the lower part of the main branch per replication per cultivar per treatment. Then the pods were dried in an oven at about 85 °C for 24 hours and weighed.

For calculating seed growth, the seeds of a pod collected from mid-position of the main branch were also dried and weighed.

Yield and Yield Components

The final harvest was done for Sambal at 100 and Daulat at 95 Days after sowing. Six plants per replication per cultivar per treatment were harvested and measurments of yield and yield components were made. The following characters were recorded :

Plant height at maturity, length of main axis, number of branches/plant, number of pods on main axis, number of pods/plant, number of seeds/pod, pod weight/plant, straw weight, 1000-seed weight, seed-husk weight ratio, harvest index, seed yield/ha and oil content of seeds. Harvest index was calculated as follows :

Economic yieldSeed yieldHarvest index % =_______ x 100 =_______ x 100Biological yieldTotal dry weight

Oil content of seeds was estimated by the Soxhlet method using petroleum ether (boiling point 40-60 $^{\circ}$ C) as solvent.

Soil Moisture Determination

Soil moisture was determined gravimetrically from five successive soil layers (0-20, 20-40, 40-60, 60-80, 80-100 cm) beginning from 15 DAS up to the final harvest at 10 days intarvals. The moisture content values were used to compute consumptive water use (Dastane, 1972) according to the following formula :

Consumptive water use (CWU) = IR + ER + $\sum_{i=1}^{n} \frac{Mb_i - Me_i}{100} \times A_i \times D_i$

Where IR =irrigation water, ER = rainfall, Mb_i = moisture % at the beginning of the season of the ith layer of the soil, Me_i = Moisture % at the end of the season of the ith layer of the soil, A_i = apparent specific gravity of the ith layer of soil, D_i = depth of the ith layer of the soil within root zone and n = no. of soil layers in the root zone D.

Water Use Efficiency (WUE)

WUE was computed by using the following formula : WUE (kg/ha/mm) = Seed yield (kg/ha)/consumptive water use (mm).



Figure. 0.3. Meteorological information during the crop period. A. Max=maximum,Min=minimum and S=soil temperature(°C) and total rainfall(histogram).B. Relative humidity at 9 A.M.(continuous line), 6 P.M.(dotted line) and sunshine hour (0-0).

Meteorological Data

Meteorological data were collected from the Regional Meteorological Station, Sampur, Rajshahi. The meteorological station is about 1 km from the experimental field. The information like maximum and minimum temperatures, humidity, rainfall, soil temperture and sunshine hours were collected (Figure 0.3).Total rainfall during the experimental period was 38 mm.

Statistical Analysis of Data

The experimental design was a split plot design and the analysis of variance was done accordingly. Least significance difference (LSD) values at 5% level were calculated, where the value of variance ratio (F) for treatment effect in case of figures and treatment, cultivar and interaction effects in case of tables were significant. Simple correlation coefficients were calculated between pair of characters.

The following notations have been used to specify statistical significance throughout the thesis :

```
* = Significant at p = 0.05
```

```
** = Significant at p = 0.01
```

```
*** = Significant at p = 0.001 and
```

```
NS = Non-significant
```

Vertical bars = LSD values at 5% level between treatments within a cultivar.

RESULTS

Choice of Growth Analysis Technique

Growth analysis has long been established as a standard technique for the study of plant growth and productivity. Out of the two approaches followed, the classical one relies strictly on the harvest data sets where parameters like mean growth rate (RGR) between a pair of harvests are taken into consideration, while the second one uses regression procedures. In the latter, a suitable mathematical function represented by a smooth curve is fitted to the recorded values of plant dry weight (W) or leaf area (LA) so that it approximates the real growth curve. However, the major difficulty encountered in the classical approach is of the sampling error of the primary data which seriously obliterates the derived functions. In the regression analysis, on the other hand, looser confidence limits are acceptable; however, the major problem lies in the choice of a suitable function that represents the best approximation of the observed growth. Amongst the various statistical models proposed, polynomial exponential model has been the most popular one because the simple mathematical and statistical properties of the model enable it to be fitted to data by an exact and relatively straightforward models of linear regression. After reviewing different methods it is concluded that use of polynomials is preferable in growth analysis. A polynomial exponential model has great potential in a purely empirical approach to the study of plant growth where the objective is to assess and compare either genetic or environmental influences. In the present experiment both

the classical and the polynomial exponential techniques were followed.

The residual sum of squares were calculated for linear, quadratic and cubic equations fitted to \log_e total dry matter per unit ground area, $\log_e LAI$, \log_e total dry matter per plant, \log_e leaf area, and \log_e leaf dry weight as a function of time (Table 1.1.).The changes in LAI, leaf area and leaf weight were adequatedly described by second degree polynomial in the two cultivars (p = 0.01), but in the case of total dry matter per unit ground area and total dry matter per plant a third degree polynomial was adequate. For comparison the second degree polynomial curves have been used in all the cases .

Coefficients obtained from quadratic fitted curves of \log_e leaf area of two mustard cultivars as influenced by soil moisture are given in Table 1.2. For all the equations the a,b, a' and b' coefficients were positive and c and c' coefficients were negative. In all the cases the coefficients for the irrigated values were greater than the rainfed values, except for c and c', where, for Sambal these were equal and for Daulat the irrigated c and c' were smaller than that of the rainfed treatment .

Total Dry Matter (TDM)

Effect of soil moisture on total dry matter of two mustard cultivars at different stages of growth is shown in Figure 1.1.

.

Table 1.1. Reduction in mean squares due to successive terms of two mustard cultivars as influenced by soil moisture.

 \log_e Total dry Matter (g m⁻²)

Sources of	df	Mean squares							
variation		Sam	bal	Daulat					
		Irrigated	Rainfed	Irrigated	Rainfed				
Reduction due to linear	1	21.331**	19.287**	20.989**	17.680**				
Deviation from linear	9	0.386	0.349	0.541	0.394				
Reduction due to quadratic	1	3.217**	2.810**	4.231**	2.893**				
Deviation from quadratic	8	0.033	0.041	0.080	0.081				
Reduction due to cubic	1	0.210**	0.228**	0.611**	0.563**				
Deviation from cubic	7	0.007	0.014	0.004	0.012				

Contd

Table 1.1. (Continued).

Sources of variation	df	Mean squares							
		Sam	ba 1	Daulat					
		Irrigated	Rainfed	Irrigated	Rainfed				
Reduction due to linear	1	1.581	0.163	0.169	1.035				
Deviation from linear	7	0.762	0.918	1.108	0.945				
Reduction due to quadratic	1	5.003**	5.838**	7.394**	5.377**				
Deviation from quadratic	6	0.056	0.098	0.060	0.207				
Reduction due to cubic	1	0.010	0.156	0.089	1.078				
Deviation from cubic	5	0.065	0.087	0.054	0.323				

log_e LAI

log_e Total Dry Matter (mg/plant)

Sources of variation	df	Mean squares							
		Samb	al	Daul	lat				
		Irrigated	Rainfed	Irrigated	Rainfed				
Reduction due to linear	1	20.360**	19.262**	22.470**	17.776**				
Deviation from linear	9	0.018	0.349	0.498	0.397				
Reduction due to quadratic	1	2.830**	2.811**	3.990**	2.918**				
Deviation from guadratic	8	0.019	0.041	0.063	0.081				
Reduction due to cubic	1	0.130**	0.231**	0.300*	0.564**				
Deviation from cubic	7	0.003	0.014	0.027	0.013				

Contd

Table 1.1. (Continued).

Sources of df		Mean squares							
variation		Samb	a]	Daulat					
		Irrigated	Rainfed	Irrigated	Rainfed				
Reduction due to linear	1	1.542	0.125	0.178	1.004				
Deviation from linear	7	0.743	0.858	1.059	0.909				
Reduction due to quadratic	1	4.881**	5.066**	6.949**	5.437**				
Deviation from quadratic	6	0.053	0.157	0.077	0.154				
Reduction due to cubic	1	0.015	0.137	0.095	0.706				
Deviation from cubic	5	0.061	0.160	0.074	0.044				

log_e Leaf Area (cm²/plant)

log, Leaf Weight (mg/plant)

Sources of	df	Mean squares								
Variation		Saut	bal	Daulat						
		Irrigated	Rainfed	Irrigated	Rainfed					
Reduction due to linear	1	1.115	0.063	0.124	1.051					
Deviation from linear	7	0.725	1.046	1.011	1.049					
Reduction due to quadratic	1	4.903**	6.370**	6.673**	5.793**					
Deviation from quadratic	6	0.029	0.159	0.067	0.259					
Reduction due to cubic	1	0.015	0.009	0.250	1.452					
Deviation from cubic	5	0.031	0.189	0.031	0.020					

Table 1.2. Constants obtained from quadratic fitted curves of log total dry matter and log leaf area of two mustard cultivars as influenced by soil moisture.

	a	b	c
	Sam	bal	
Irrigated	5.3429	0.1436	-0.0012
Rainfed	5.1368	0.1415	-0.0012
	Dau	lat	
Irrigated	5.1479	0.1620	-0.0014
Rainfed	5.0264	0.1407	-0.0012

log, Total Dry Matter (g/plant)

log_e Leaf Area (cm²/plant)

	\mathbf{a}^{\prime}	ь/	c/
	Sar	nbal	
Irrigated	3.4617	0.1669	-0.0026
Rainfed	3.4516	0.1529	-0.0026
	Dau	lat	
Irrigated	3.5914	0.1796	-0.0031
Rainfed	3.5401	0.1336	-0.0027

Increase of total dry matter was slow at the early vegetative phases (20, 27, 34 and 41 DAS), but increased rapidly with the advancement of the growth period in both the cultivars and treatments. The irrigated plants had always greater total dry matter than the rainfed plants. Significant increase of total dry matter production with irrigation was observed at almost all the stages of growth except at 20, 76 and 90 DAS in Sambal and 20 DAS in Daulat. Daulat had relatively higher dry matter accumulation than Sambal in the irrigated condition but it had lower dry matter in the rainfed condition .

Mean squares from the analysis of variance of curve-fitted values of TDM, LAI, LAD and some other growth attributes of two mustard cultivars as influnced by soil moisture are presented in Table 1.3. In case of TDM (g m⁻²) significant effect was found for treatment, harvest date, cultivar x treatment and treatment x harvest date. The curves of TDM (Figure 1.1) indicated that they were not adequately fitted. The curve-fitting method yielded unusual curves for TDM, where there was a decline at the later stages of growth.

Leaf Area Index (LAI)

Effect of soil moisture on LAI showed that the pattern of its development was essentially similar in both the treatments and cultivars(Figure 1.2). In every cultivar and treatment starting from a lower value, LAI reached a certain peak and then declined to zero. In Sambal, both the irrigated and the rainfed LAI reached their peak



Figure 1.1. Effect of soil moisture on total dry matter (TDM) of two mustard cultivars at different stages of growth. O=irrigated and A=rainfed from original values; smooth line for irrigated and dotted line for rainfed from quadratic curve-fitted values.

45

at 55 DAS but in Daulat the irrigated LAI at 55 and rainfed LAI at 48 DAS. The irrigated plants had higher LAI at all the stages of growth. Significantly higher LAI was found in the irrigated plants for Sambal at 24, 48, 55, 62 and 76 DAS and for Daulat at all the stages of growth except at 20 and 76 DAS. The highest LAI was found in the irrigated Daulat and the lowest in the rainfed Daulat .

Mean squares from the analysis of variance of curve-fitted values of LAI indicated that the soil moisture, cultivar and harvest date effects were significant (Table 1.3). The interactions like cultivar x treatment, treatment x harvest date were also significant. LAI calculated by the harvest interval method showed a close a correspondence with the instantaneous values of LAI obtained from curve-fitting method (Figure 1.2). Maximum LAI was under-estimated in the fitted data in all cases, except in the irrigated Sambal, where it was over-estimated .

Leaf Area Duration (LAD)

The pattern of LAD was similar as in LAI (Figure 1.3). In Sambal, LAD reached its peak at (55-62) DAS in both the irrigated and the rainfed plants, whereas in Daulat, it reached its peak at (48-55)DAS in the irrigated plants and at (41-48)DAS in the rainfed plants. The LAD values of the irrigated plants were higher than that of the rainfed plants in both the cultivars. The irrigated plants had significantly greater LAD at (20-27), (27-34), (48-55) and (62-69) DAS in Sambal and in Daulat at all the stages of growth except



Figure 1.2. Influence of soil moisture on leaf area index (LAI) of two mustard cultivars at different stages of growth. O=irrigated and A=rainfed from original values; smooth line for irrigated and dotted line for rainfed from quadratic curve-fitted values.

Table 1.3. Mean squares from the analysis of variance of TDM, LAI, LAD and some other growth attributes of two mustard cultivars as influenced by soil moisture (calculated from quadratic curve-fitted values).

Sources of TD variation df	T	TDH (g m ⁻²⁾		LAI		LAD (days)		$CGR (g m^{-2} day^{-1})$		RGR $(g g^{-1} day^{-1})$	
	Nean squares	dſ	Nean squares	df	Wean squares	df	Kean squares	df	Kean squares		
Treatment (T)	1	280112***	1	1.3728***	I	67.29***	1	201.960***	1	950.59**	
Cultivar (C)	1	37	1	0.5560**	1	9.88*	1	216.640***	1	199.33	
Harvest (H)	10	165803***	8	0.1981**	8	31.46**	9	5.520	10	24205.93***	
TrC	1	9524	1	0.1381*	1	6.80*	1	0.002	1	522.20*	
CrH	10	520	8	0.0272	8	1.12	9	4.360	10	542.95**	
T x H	10	20771***	8	0.0695*	8	3.45*	9	2.350**	10	308.18**	
TxCxH	10	753	8	0.0146	8	0.94	9	7.010	10	78.60	

Sources of variation	df	Nean squares						
		NAR x 10 ⁻⁴ (g cm ⁻² day ⁻¹)	LAR (cm ² g ⁻¹)	$(cm^2 cm^{-2} day^{-1})$	SLA ($cn^2 g^{-1}$)	LWR $(g g^{-1})$		
Treatment (T)	1	0.00033**	323.94**	0.00360***	0.21	0.0062		
Cultivars (C)	1	0.00001	365.90*	0.00280***	5649.99***	0.0062		
Harvest (H)	8	0.01296***	464.93**	0.04224***	387.42	0.4125***		
TrC	1	0.00013*	61.80	0.00032*	1750.07*	0.0182*		
CxH	8	0.00002	76.01	0.00012	102.47	0.0093*		
ТхН	8	0.00003	63.47	0.00006	516.69	0.0007		
TICIH	8	0.00002	27.02	0.00005	189.58	0.0018		



Figure.1.3. Effect of soil moisture on leaf area duration (IAD) of two mustard cultivars at different stages of growth, O=irrigated and △=rainf rom original data; smooth line for irrigated and c d line for rainfed from quadratic curve-fitted ds

at (55-62) DAS. Irrigated Daulat showed the highest and rainfed Daulat showed the lowest LAD.LAD calculated by the classical method showed a close agreement with the LAD obtained from the curvefitted method (Figure 1.3). In the fitted data, all the maximum LAD values were under-estimated, except in the irrigated Sambal, where LAD was over-estimated .

Crop Growth Rate (CGR)

CGR fluctuated in both the cultivars and treatments (Figure 1.4a).Though it increased slightly at the early stages and declined at the later stages, yet no clear pattern was found. Highest CGR was found at (48-55) DAS in Sambal and at (41-48) and (34-41) DAS in the irrigated and the rainfed Daulat, respectively. The irrigated plants had higher CGR at all the stages of growth than the rainfed plants except at (69-76) DAS in Sambal.Significantly higher CGR was found at (62-69) and (76-83) DAS in Sambal and at all the stages of growth in Daulat except (34-41), (55-62) and (69-76) DAS in the irrigated plants than that of the rainfed plants. The highest CGR was observed in the irrigated Daulat and the lowest in the rainfed Daulat.

Mean squares from the analysis of variance of the curvefitted values of CGR shown in Table 1.3 indicate that soil moisture treatment differences were significant. Cultivar and treatment and treatment x harvest date interaction were also significant. CGR calculated from the quadratic-fitted data gave uncharacteristically negative values late in the season (Figure 1.4b). In Daulat, CGR in






Figure.1.4b. Influence of soil moisture on crop growth rate (CGR) of two mustard cultivars at different stages of growth (smooth line for irrigated and dotted line for rainfed from quadratic fitted values).

negative values late in the season (Figure 1.4b). In Daulat, CGR in the irrigated plants became more negative than the rainfed plants.

Relative Growth Rate (RGR)

Effect of soil moisture on RGR of two mustard cultivars at different stages of growth is shown in Figure 1.5a. In both the cultivars, RGR of both treatments declined with slight fluctuation. At the early stages, it declined sharply but at the later stages it declined slowly. In irrigated condition, Sambal had higer RGR at (34-41), (48-55) and (69-76) DAS and Daulat at (20-27), (27-34), (41-48) and (48-55) DAS. Irrigated Daulat had significatly higher RGR at (41-48) DAS than the rainfed plants. The highest RGR was observed in Daulat .

Analysis of variance of the curve-fitted values of RGR (Table 1.3) indicates that the effect of soil moisture was significant. The effect of harvest date and interaction items like cultivar x treatment, cultivar x harvest date and treatment x harvest date were also significant.RGR declined sharply throuhout from a maximum value (Figure 1.5b). RGR calculated from the quadratic-fitted data gave also uncharacteristically negative values at the later stages of growth .

Net Assimilation Rate (NAR)

The pattern of soil moisture effect exhibited that NAR



Figure 1.5a. Effect of soil moisture on relative growth rate (RGR) of two mustard cultivars at different stages of growth (by classical technique).



Figure 1.5b. Influence of soil moisture on relative growth rate (RGR) of two mustard cultivars at different stages of growth (Smooth line for irrigated and dotted line for rainfed from quadratic fitted values).

fluctuated in both the treatments of Sambal and Daulat (Figure 1.6a). Very little increase or depletion was found in Sambal up to (62-69) DAS and in Daulat upto (48-55) DAS and then increased very rapidly, except in the irrigated Sambal, where no such increase was observed. NAR values of the rainfed plants was higher at (27-34), (41-48), (48-55), (55-62) and (69-76) DAS in Sambal and at (34-41), (48-55), (55-62), (69-76) DAS in Daulat. The rainfed plants had significantly higher NAR at (69-76) DAS in Sambal and at (55-62), (62-69) and (69-76) DAS in Daulat. Maximum NAR value was found in rainfed Daulat .

Mean squares from the analysis of variance of the curvefitted data of NAR (Table 1.3) indicates that soil moisture treatment was highly significant. Harvest date differences were also highly significant. Cultivar and treatment interaction was significant. NAR calculated by the harvest interval method showed a close correspondence with the instantaneous values obtained from the curve-fitting method (Figures 1.6a and 1.6b). The rainfed values remained lower at the early stages but became higher at the later stages of growth .

Leaf Area Ratio (LAR)

LAR values of both the treatments of the two cultivars were found maximum at (20-27) DAS and then declined steadily with increasing plant age (Figure 1.7a). The irrigated plants had higher LAR at all the growth stages in Sambal except at (20-27) and (41-48)







Figure 1.6b. Effect of soil moisture at different stages of growth on net assimilation rate (NAR) of two mustard cultivars (Smooth line for irrigated and dotted line for rainfed values from quadratic fitted values).

DAS. In the irrigated plants significantly higher LAR was found at (27-34), (48-55) and (55-62) DAS in Sambal and at (48-55), (55-62) and (62-69) DAS in Daulat than the rainfed plants. Significantly lower LAR was observed at (20-27) in the irrigated Daulat.The highest LAR was found in the rainfed Daulat.

Highly significant effect of soil moisture on LAR was present (Table 1.3). Cultivar and harvest date differences were also significant. The pattern of LAR calculated from the curve-fitted data was closer to that calculated from the original data (Figures 1.7a and 1.7b).

Relative Leaf Growth Rate (RLGR)

The maximum RLGR was found at (20-27) DAS in both the treatments and cultivars and then declined gradually with fluctuations (Figure 1.8a). In the irrigated Sambal plants, RLGR became negative at (55-62)DAS and in the rainfed Sambal it became negative first at (41-48) DAS, then increased into positive value and further became negative at (55-62) DAS. In the irrigated Daulat plants, RLGR became negative at (48-55) DAS and in the rainfed plants it became negative at (41-48) DAS. The highest RLGR was found in the irrigated plants of both the cultivars. Irrigated plants had higher RLGR at (20-27), (41-48), (48-55) and (69-76) DAS in Sambal and (20-27), (41-48), (48-55) and (62-69) DAS in Daulat than the rainfed plants. Significantly higher RLGR in the irrigated plants was found at (27-34) and (48-55) DAS in Daulat and (69-76) DAS in Sambal. Significantly lower RLGR in the irrigated plants was observed at (55-62)DAS in Sambal and at (69-76) DAS in Daulat than the rainfed plants.













Figure.1.8b. Soil moisture effect on relative leaf growth rate (RLGR) of two mustard cultivars at various stages of growth (Smooth line for irrigated and dotted line for rainfed from quadratic fitted values).

Soil moisture treatment, cultivar and harvest date differences for RLGR, calculated from curve-fitted data were highly significant (Table 1.3). Cultivar x treatment effect was significant. Soil moisture effect on RLGR calculated from curve-fitted values showed a linear drift with increasing age (Figure 1.8b).

Specific Leaf Area (SLA)

There were no clear patterns of soil moisture effect on SLA (Figure 1.9a). The irrigated plants had significantly higher SLA than the rainfed plants at 34, 48, 55 and 62 DAS in Sambal and 34 and 55 DAS in Daulat. The irrigated Daulat had significantly lower SLA at 48 DAS than the rainfed plants .

Mean squares from the analysis of variance of the curvefitted data of SLA (Table 1.3) showed that the cultivar differences were highly significant. Cultivar x treatment was also significant. The curve-fitted values of SLA (Figure 1.9b) indicate that there was slight increase or depletion of SLA with increasing age. The irrigated plants had higher SLA at the middle stages of growth but it was lower at both the initial and terminal stages .

Leaf Weight Ratio (LWR)

Starting from a maximum value, LWR of both the treatments and cultivars declined sharply with increasing age (Figure 1.10a). The exception was found only at 27 DAS in the rainfed Daulat. Significantly higher LWR in the irrigated plants than the rainfed



Figure 1.9a. Effect of soil moisture on specific leaf area (SLA) of two mustard cultivars at various stages of growth (by classical technique).



plants was found at 48 and 76 DAS in Sambal and at 55, 62 and 69 DAS in Daulat. The irrigated Sambal had significantly lower LWR at 69 DAS .

Highly significant effect of harvest date was found on LWR (Table 1.3). The interactions between cultivar and treatment and cultivar and harvest date were also found significant. The pattern of LWR obtained from the curve-fitted values was closure to LWR obtained from the original values (Figures 1.10a and 1.10b). LWR also showed downward drifts with age (Figure 1.10b).

The values of RGR and NAR (Figure 1.11), LAR and RLGR (Figure 1.12) and SLA and LWR (Figure 1.13) plotted against \log_e plant dry weight, based on the fitted quadratic relationship between $\log_e W$ (dry weight) and $\log_e LA$ (leaf area) with time, are shown. The curves of RGR, LAR, RLGR and LWR with respect to \log_e plant dry weight in both the treatments and cultivars showed similar declining trends. Very little increase or decrease of NAR was found at the initial stages but it increased very sharply with increasing dry weight. The pattern of SLA with increasing dry weight was similar as was with increasing time.

Correlation Coefficients Between Growth Attributes and Yield

Simple correlation coefficients between growth attributes and yield at two soil moisture levels are given in Table 1.4.



Figure 1.10a. Effect of soil moisture on leaf weight ratio (LWR) of two mustard cultivars at different stages of growth (by classical technique).



Figure 1.10b. Soil moisture effect on leaf weight ratio (LWR) of two mustard cultivars at different growth stages (Smooth line for irrigated and dotted line for rainfed from quadratic fitted values).



with time (Smooth line for irrigated and dotted line for rainfed values).





Association between growth attributes and yield of irrigated plants

CGR at the pre-flowering stage was positively correlated with pre-flowering LAI and negatively correlated with the postflowering RGR. RGR at the post-flowering stage was negatively

correlated with LAI at the pre-flowering stage. NAR at the postflowering stage was positively correlated with the pre-flowering LAR and negatively correlated with the post-flowering LAR. LAR at the pre-flowering stage was positively correlated with the postflowering LAR. Seed yield was positively correlated with the preflowering LAI and CGR and the post-flowering NAR .

Association between growth attributes and seed yield of rainfed plants

CGR at the pre-flowering stage was positively correlated with RGR and NAR at the pre-flowering stage. CGR at the post-flowering stage was positively correlated with LAI (post-flowering) and negatively correlated with NAR (post-flowering) and LAR (preflowering).RGR at the pre-flowering stage was positively associated with NAR at the pre-flowering stage. The pre-flowering NAR was positively correlated with pre-flowering LAI. The post-flowering NAR was positively related with LAR (pre-flowering) and negatively with LAI (pre-flowering) and LAR (post-flowering).The pre-flowering LAR was negatively and the post-flowering LAR was positively associated with the post-flowering LAI. Seed yield was positively Table 1.4.Simple correlation coefficients between growth attributes and yield (upper diagonal irrigated and lower diagonal rainfed values).

	LAI		CGR		RGR		NAR		LAR		Seed yeild
	Pre- flow- ering (1)	Post- flow- ering (2)	Pre- flow- ering (3)	Post- flow- ering (4)	Pre- flow- ering (5)	Post- flow- ering (6)	Pre- flow- ering (7)	Post- flow- ering (8)	Pre- flow- ering (9)	Post- flow- ering (10)	(11)
1		-0.52	0.89*	-0.06	0.60	-0.81*	0.11	0.73	0.89*	-0.89*	0.82*
2	0.01		-0.74	0.46	-0,70	0.65	-0.53	-0.57	-0.55	0.70	0.01
3	0.78	-0.52		-0.38	0.73	-0.97**	0.48	0.51	0.76	-0.77	0.18
4	0.03	0.81*	-0.48		-0.51	0.37	-0.76	0.25	0.26	0.07	0.84*
5	0.71	-0.53	0.86*	-0.65		-0.56	0.76	0.48	0.39	-0.72	-0.29
6	0.51	0.64	0.06	0.72	-0.21		-0.44	-0.34	-0.71	0.61	-0.19
7	0.79*	-0.35	0.89*	-0.55	0.93**	0.06		-0.17	-0.11	-0.14	-0.77
8	-0.05	-0.97**	0.50	-0.80*	0.44	-0.54	0.34		0.82*	-0.92**	0.81*
9	0.04	-0.97**	0.57	-0.79*	0.60	-0.70	0.38	0.89*		-0.84*	0.67
10	0.50	0.84*	-0.02	0.77	-0.14	0.81*	0.05	-0.85*	-0.77		-0.45
11	-0.25	0.84*	-0,76	0.86*	-0.69	0.36	-0.70	-0.89*	-0.79*	0.60	

correlated with LAI and CGR at the post-flowering stage and negatively correlated with the post-flowering NAR and the preflowering LAR.

Distribution of Dry Matter

Effect of soil moisture on distribution of dry matter (DM) of two mustard cultivars at successive stages of growth is shown in Figures 1.14 and 1.15. At the early stages of growth only stem + petiole and leaves were present, whereas, at the later stages only stem and pods were present. Percentage of DM accumulation for stem + petiole was lower at the early stages, at the middle stages it reached a certain peak and then gradually declined with time. In Sambal the irrigated plants had peak stem + petiole DM (78%) at 62 DAS and the rainfed plants (78%) at 55 DAS. Daulat had its peak DM of stem + petiole (70.4%) at 55 DAS in the irrigated plants and (74%) also at 55 DAS in the rainfed plants.

Percentage of DM accumulation in the leaves revealed that it was highest at the initial stage and gradually declined with increasing age and at 83 DAS it became zero. The irrigated plants accumulated higher leaf DM than the rainfed plants at 20, 48, 55, and 76 DAS in Sambal and at 20,41, 55, 62, 69 and 76 DAS in Daulat.

Dry matter accumulation for buds + flowers started at 41 DAS and gradually increased up to several harvests and then declined with increasing time. The irrigated plants had higher percentage of



Figure 1.14. Influence of soil moisture on distribution of dry matter (DM) of Sambal at successive stages of growth.



Figure 1.15. Influence of soil moisture on distribution of dry matter (DM) of Daulat at successive stages of growth.

DM accumulation for buds + flowers than the rainfed plants at 55 and 62 DAS in Sambal.

Percentage of DM accumulation for pods was found lowest at 48 DAS and gradually increased with plant age. Higher percentage of DM accumulation was found in the rainfed plants at 62, 83 and 90 DAS in Sambal and at 55, 62 and 69 DAS in Daulat .

Relative Leaf Water Content (RLWC)

The effects of soil moisture on RLWC of two cultivars at different times of a day at the flowering and the mid-pod filling stages are given in Figures 1.16a and 1.16b. The irrigated plants had higher RLWC than that of the rainfed plants in all cases. At the flowering stage, the irrigated plants had significantly higher RLWC than the rainfed plants at 8 A.M. in both the cultivars. At mid-pod filling stage, the irrigated plants had significantly higher RLWC at 8 A.M. and 4 P.M.than the rainfed plants of Sambal. The highest RLWC was observed at 8 A.M. but at the midday it decreased and at the later part of the day it increased gradually.

Chlorophyll, Proline and Sugar Contents

Mean values of chlorophyll, proline and sugar contents of two cultivars as influenced by soil moisture are given in Table 1.5.



Figure 1.16a. Relative leaf water content (RLWC) of two mustard cultivars as influenced by soil moisture at flowering stage.



Figure 1.16b. Relative leaf water content (RLWC) of two mustard cultivars as influenced by soil moisture at mid-pod filling stage.

Soil moisture effect was not significant for chlorophyll a, chlorophyll b and total chlorophyll at both the flowering and midpod filling stages.Chlorophyll b of Daulat was significantly higher than that of Sambal plants at the flowering stage. At the mid-pod filling stage chlorophyll a ,b and total chlorophyll were slightly higher in the irrigated plants than the rainfed plants .

The rainfed plants had significantly higher proline than that of the irrigated plants at both the flowering and mid-pod filling stages. At the flowering stage, Sambal had significantly higher proline than Daulat. At the mid-pod filling stage such difference was not significant. Proline content was higher at the mid-pod filling stage than that of the flowering stage .

The leaves of the rainfed plants accumulated significantly higher quantity of sugars at both the flowering and mid-pod filling stages. Rainfed Sambal accumulated significantly higher sugars at both the stages. At the mid-pod filling stage, rainfed Sambal had significantly higher sugars than the rainfed Daulat. Accumulation of sugar was higher at the mid-pod filling stage than the flowering stage.

Relative Leaf Water Content (RLWC) and Time At Which Stomata Closed

The curves from which relative leaf water content and the time taken to stomatal closure were calculated by extrapolation are Table 1.5. Mean values of chlorophyll, proline and sugar contents at the flowering and the mid-pod filling stages of two mustard cultivars as influenced by soil moisture.

-	Chlorophyll a (mg dm ⁻²)			Chlorophyll b (mg dm ⁻²)			Total chlorophyll(mg dm ⁻²)		
	Samba l	Daulat	Mean	Samba 1	Daulat	Mean	Samba 1	Daulat	Mean
				Floweri	ng stage				
Irrigated	2.42	2.72	2.57	0.97	1.09	1.03	3.40	3.80	3.61
Rainfed	2.49	2.58	2.53	1.00	1.14	1.07	3.49	3.72	3.60
Mean	2.46	2.65		0.99	1.12		3.45	3.76	
LSD at 5%	SD at 5% (a)NS,(b)NS,(c)NS,(d)NS		(a)NS,(b)0.10,(c)NS,(d)NS			(a)NS,(b)NS,(c)NS,(d)NS			
			Mi	d-pod fi	lling sta	ge			
Irrigated	2.73	2.62	2.68	1.07	1.14	1.11	3.81	3.93	3.87
Rainfed	2.20	2.26	2.23	1.01	0.95	0.98	3.21	3.21	3.21
Mean	2.47	2.44		1.04	1.05		3.51	3.57	
LSD at 5%	(a)NS,(b))NS,(c)NS	,(d)NS	(a)NS,(b)NS,(c))	s,(d)NS	(a)NS,(1)NS,(c)NS	, (d) NS

Proline (_µ g (g ⁻¹ fresh weigh	nt of leaf)	Sugars content (mg g^{-1} dry weight)				
Samba l	Daulat	Mean	Samba 1	Daulat	Mean		
]	lowering stage	9				
68	60	64	8.1	10.1	9.1		
100	78	89	13.1	13.7	13.4		
84	69		10.6	11.9			
LSD at 5% (a)20,(b)14,(c)14,(d)13			(a)3.6,(b)NS,(c)NS,(d)2.3				
Mid-pod filling			tage				
145	121	133	12.4	14.9	13.7		
187	163	175	22.9	17.5	20.2		
166	142		17.6	16.2			
(a)31,(b)NS	,(c)NS,(d)40		(a)1.1,(b)NS,(c)3.6,(d)3.2				
	Proline (µg a Sambal 68 100 84 (a)20,(b)14 145 187 166 (a)31,(b)NS	Proline (µg g ⁻¹ fresh weigh Sambal Daulat 68 60 100 78 84 69 (a)20,(b)14,(c)14,(d)13 Mid- 145 121 187 163 166 142 (a)31,(b)NS,(c)NS,(d)40	Proline (µg g ⁻¹ fresh weight of leaf) Sambal Daulat Mean Sambal Daulat Mean Flowering stage 68 60 64 100 78 89 84 69	Proline (µg g ⁻¹ fresh weight of leaf) Sugars contained Sambal Daulat Mean Sambal Baulat Mean Sambal Sambal Flowering stage 68 60 64 8.1 100 78 89 13.1 84 69 10.6 (a)20,(b)14,(c)14,(d)13 (a)3.6,(b)N Mid-pod filling stage 145 121 133 12.4 187 163 175 22.9 166 142 17.6 (a)31,(b)NS,(c)NS,(d)40 (a)1.1,(b)N (a)1.1,(b)N (a)1.1,(b)N	Proline (μ g g^{-1} fresh weight of leaf)Sugars content (mg g^{-1} dry SambalSambalDaulatMeanSambalDaulatFlowering stage6860648.110.1100788913.113.7846910.611.9(a)20,(b)14,(c)14,(d)13(a)3.6,(b)NS,(c)NS,(d)2.3Mid-pod filling stage14512113312.414.918716317522.917.516614217.616.2(a)1.1,(b)NS,(c)3.6,(d)3.		

(a) = Difference between treatments

.

(b) = Difference between cultivars

.

(c) = Difference within treatment between cultivars

(d) = Difference within cultivar between treatments



Figure 1.17a. Effect of soil moisture on relative leaf water content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at flowering stage.



Figure 1.17c. Effect of soil moisture on relative leaf water content and stomatal closure showing extrapolations of the straight line parts intersecting at closure at mid-pod filling stage.



Figure 1.17d. Effect of soil moisture on relative leaf water content and stomatal closure showing extrapolation of the straight line parts intersecting at closure at mid-pod filling stage.

Table 1.6. Mean values of relative leaf water content (RLWC) and time at which stomata closed of two mustard cultivars as influenced by soil moisture at the flowering and the mid-pod filling stages.

	RLWC a closed	t which s (%)	stomata	Time taken to stomatal closure (min)			
	Sambal	Daulat	Mean	Sambal	Daulat	Mean	
4		Flowe	ering sta	ge			
Irrigated	76.00	73.70	74.85	66.00	75.00	70.50	
Rainfed	77.50	74.20	75.85	51.70	52.70	51.90	
Mean	76.80	74.00		58.90	63.90		
LSD at 5%	(a)NS,(b (d)NS)NS,(c)NS	IS, (a)14.50,(b) (d)18.70			c)NS,	
r 1		Mid-pod	filling	stage			
Irrigated	74.50	68.50	72.60	61.70	69.00	58.50	
Rainfed	72.80	71.50	73.30	54.00	67.00	60.50	
Mean	73.65	70.00		51.15	68.00		
LSD at 5%	(a)NS,(b)NS,(c)]	NS,(d)NS	(a)NS,(b)NS,(c)N	S,(d)NS	

(a)	=	Difference	between treatments
(b)	=	Difference	between cultivars
(c)	Ξ	Difference	within treatment between cultivars
(d)	=	Difference	within cultivar between treatments
shown in Figures 1.17a-d. The relative turgidity at the point of intersection was taken as a critical level for stomatal closure. Relative water content of leaves at points of stomatal closure and time at which stomata closed at the flowering and the mid-pod filling stages are given in Table 1.6. Treatment and cultivar effects were non-significant at both the flowering and mid-pod filling stages for both the characters, but at the flowering stage, time taken to stomatal closure was significantly higher in the irrigated plants.

Transpiration Rates

Mean values of transpiration rates of two cultivars as influenced by soil moisture at the flowering and mid-pod filling stages are presented in Table 1.7. At the flowering stage, soil moisture effect was found significant on total transpiration rate. Total transpiration rate was significantly higher in the irrigated plants at the flowering stage. Sambal had higher rate of total and cuticular transpiration at the flowering stage. At the mid-pod filling stage, soil moisture effect on total, stomatal and cuticular transpiration rates were found non-significant. The irrigated Sambal had singificantly higher cuticular rate than that of the rainfed plants. The stomatal transpiration rate in all the cases .

Stomatal Characters

Mean values of stomatal frequency, stomatal pore length, number

Table 1.7. Mean values of transpiration rates (mg cm⁻² h⁻¹) of two mustard cultivars as influenced by soil moisture at the flowering and the mid-pod filling stages.

	Total t	ranspirati	on rate	Stomata	l transpirat	tion rate	Cuticular	r transpirat	ion rate
	Sambal	Daulat	Nean	Sambal	Daulat	Nean	Sambal	Daulat	Nean
				Floweri	ng stage				
Irrigated	9.67	7.65	8.66	6.99	6.39	6.69	2.68	1.26	1.97
Rainfed	8.24	7.28	7.76	6.52	5.66	6.09	1.72	1.62	1.67
Nean	8.96	7.47		6.76	6.03		2.20	1.44	
LSD at 5%	(a)0.85,(b)1.25,(c)	NS,(d)1.42	(a)NS,(b)NS,(c)1.09,	(d)NS	(a)NS,(b)().71,(c)1.40	,(d)0.87
			H	id-pod fi	lling stage	:			
lrrigated	6.26	6.55	6.41	5.44	5.86	5.66	2.81	0.69	0.75
Rainfed	6.40	6.59	6.50	5.89	6.12	6.01	0.51	0.47	0.49
Wean	6.33	6.57		5.67	6.01		0.66	0.58	
LSD at 5%	(æ)NS,(b)NS,(d)NS,(d)NS	(a)NS,(b)NS,(c)NS,(c	I)NS	(a)NS,(b)N	IS,(c)0.21,(d)0.19

(a) = Difference between treatments

(b) = Difference between cultivars

.

(c) = Difference within treatment between cultivars

(d) = Difference within cultivar between treatments

of epidermal cells and stomatal index of both the adaxial and abaxial surfaces are given in Table 1.8. Although soil moisture effect was not significant, stomatal frequency of the abaxial surface was higher than that of the adaxial surface. The irrigated plants had lower stomatal frequency than the rainfed plants. Stomatal pore lenth was also unaffected by soil moisture .

Significant effect of soil moisture on epidermal cells of the adaxial surface was found. The irrigated plants had significantly lower number of epidermal cells on both surfaces of Sambal and on the abaxial surface of Daulat. The irrigated plants had significantly higher stomatal index on the abaxial surface. The irrigated Sambal plants had significantly higher stomatal index than that of the rainfed plants on both the adaxial and abaxial surfaces, while the irrigated Daulat had on the adaxial surface only. The stomatal index was higher in the abaxial surface than that of the adaxial surface.

Anatomical Characters of Leaves

Mean values of leaf thickness and cross-sectional areas of palisade and spongy paranchyma cells are presented in Table 1.9. Leaf thickness was significantly higher in the irrigated plants. The irrigated plants showed increased thickness of leaves and mean cross-sectional areas of palisade and spongy parenchyma cells. The mean cross-sectional area of spongy cells of Daulat was higher than that of Sambal. Within the irrigated plants, Daulat had higher leaf thickness and mean cross-sectional areas of palisade and spongy Table 1.8. Mean values of stomatal characters of two mustard cultivars as influenced by soil moisture .

	Stows (No. o	ital freque f stowata	ency 11 ⁻²)	Stowatal	pore leng	(th(pm)	Naube c	r of epide ells ss	rmal	Stomatal inder (%)		
	Sambal	Daolat	Nean	Saubal	Daulat	Nean	Saubal	Daelat	Kean	Sambal	Daulat	Rean
					Adaxia	al surface	e					
Irrigated	162	179	170	20.91	17.81	19.36	524	701	612	23	20	22
Rainfed	192	196	193	19.41	20.27	19.84	724	737	731	21	21	21
Nern	177	188		20.16	19.04		624	719		22	21	
LSD at 5X	(a)NS,(b)	NS,(c)NS,(d) RS	(a)NS,(b)	NS,(c)NS,(d)NS	(a)#S,(b)88,(c)88	,(d)184	(▲)∦S,(b)NS,(c)2.	0,(d)1.7
					Abari	al suface	!					
Irrigated	343	318	330	20.27	18.56	19.41	666	739	702	34	30	32
Rainled	356	319	338	20.69	21.33	21.01	816	916	866	30	26	28
Kean	350	319		20.48	19.95		714	828		32	28	
LSD at 5X	(a)N8,(b)NS,(c)NS,	(d)118	(a)RS,(b)AS,(c)AS,	, (d) 88	(a)135,	(b)185,(c)1	18, (d)115	(a)1.8,	(b)#S,(c)#	18,(d)3.9

(a) = Difference between treatments

(b) = Difference between cultivars

(c) = Difference within treatment between cultivars.

(d) = Difference within cultivar between treatments

parenchyma cells. In the irrigated plants, all these characters were higher than that of the rainfed plants in both the cultivars .

Table 1.9. Mean values of some anatomical characters of leaves of two mustard cultivars as influenced by soil moisture.

	Leaf thi	ckness(p n)	Cross-see palisade	ctional area parenchyma	of (µ ²)	Cross-sectional area of spongy parenchyma(pm ²)			
	Sambal	Daulat	Nean	Sambal	Daulat	Nean	Sambal	Daulat	Nean	
Irrigated	330	365	378	1000	862	931	511	394	453	
Rainfed	296	319	308	827	738	783	409	341	375	
Kean	313	342		914	800		460	368		
LSD at 5%	(a)31,(b)NS,(c)28,(d)19			(a)NS, (1	(a)NS,(b)NS,(c)NS,(d)NS			(a)NS,(b)NS,(c)NS,(d)NS		

(a) = Difference between treatments, (b) = Difference between cultivars

(c) = Difference within treatment between cultivars, (d) = Difference within cultivar between treatments

Characters Related to Flowering

Mean effects of soil moisture on plant height, number of leaves at first flowering, days taken to first flowering and duration of flowering are given in Table 1.10. Plant height at first flowering and duration of flowering were significantly higher in irrigated plants than that of the rainfed plants. All these the characters were significantly higher in Sambal, except duration of flowering. Within the irrigated plants, plant height and the number of leaves at first flowering were significantly higher in Sambal. Within the rainfed plants, all the above characters were significantly higher in Sambal, except duration of flowering. The irrigated plants were significantly taller in both the cultivars. The number of leaves was significantly higher in the irrigated plants than that of the rainfed plants of Sambal.

Table 1.10. Mean values	of some characters at	flowering stage of
two mustard	cultivars as influenced	1 by soil moisture.

	Plant height at first flowering (cm)		•)	Wumber of leaves at first flowering			Days taken to first flowering			Duration of flowering (days)		
	Sanbal	Daulat	Nean	Saubal	Daulat	Nean	Saubal	Daulat	Keaa	Seubal	Daulat	Kean
Irrigated	72.2Ż	69.67	71.94	13.67	11.11	12.39	38.67	38.00	38.34	22.89	20.11	21.50
Raialed	64.67	59.89	62.28	12.22	10.78	11.50	39.22	38.00	38.61	20.22	17.11	18.67
Kean	68.45	64.78		12.95	10.95		38.95	38.00		21.56	18.61	
LSD at 5X	(a)7.32, (c)3.99,	(b)2.82, (d)2.91		(a)NS,(b (c)1.78,)0.83, (d)0.80		(a)NS,(b (c)0.79,)0.55, (d)N8		(a)2.19,((c)N8,(d)	b) N 8 N 8	

Table 1.11. Mean values of pod abortion (%) on main axis and other branches of two mustard cultivars as influenced by soil moisture.

.

		Nain axis			Primary branch 1			Primary branch 2			Primary branch 3		
	Saabal	Daulat	Kean	Sambal	Daulat	Kean	Saubal	Danlat	Keaz	Sambal	Daulat	Kean	
Irrigated	28.95	35.95	32.45	47.44	51.11	49.28	51.00	58.33	54.67	54.67	59.89	57.28	
Rainfed	30.81	32.32	31.57	51.00	51.67	51.34	55.28	60.67	57.98	60.11	66.66	63.39	
Keaa	29.88	34.14		49.22	51.39		53.14	59.50		57.39	63.28		
LSD at SX	(a)N8,(b)) (c)3.24,(d	2.29, 1)2.65		(a)88,(b (c)88,(d) #S ,) #S		(a)88,(b) (c)88,(d))6.18,) 8 8		(a)18,(t (c)18,(d)		

(a) = Difference between treatments

(b) = Difference between cultivars

(c) = Difference within treatment between cultivars

(d) = Difference within cultivar between treatments

Flowering Pattern

Effect of soil moisture on cumulative number of flowers on the main axis and the primary branches 1-3 of two cultivars is shown in Figures 1.18a-d. The irrigated plants had always higher number of flowers than the rainfed plants. There was little variation regarding the initiation of flowering on different branches between treatments, except in primary branch 3, where flowers of the irrigated Sambal initiated at 44 DAS and the rainfed plants initiated at 48 DAS. The initiation of flowers on the main axis, primary branches 1, 2 and 3 were at 38, 42, 42 and 44 DAS, respectively.Flowering of the irrigated and the rainfed plants in all cases took place at the same time but in branch 3 it was accomplished later in the irrigated than the rainfed plants in both the cultivars. Thus, the duration of flowering of branch 3 of the irrigated plants was higher than the rainfed plants.

Pod Abortion

Mean values of pod abortion on the main axis and the primary branches 1-3 as influenced by soil moisture are given in Table 1.11.Pod abortion was less in the irrigated plants than the rainfed plants, though they were not significantly different. Significantly higher number of pods was aborted on the main axis and primary branch 2 of Daulat. The irrigated Daulat had lesser pod abortion than the rainfed plants on the main axis. The highest percentage of pod abortion was observed on the primary branch 3 and the lowest



Figure 1.18a. Influence of soil moisture on cumulative number of flowers opened on main axis.

95



Figure 1.18b. Influence of soil moisture on cumulative number of flowers opened on primary of branch 1.



Figure 1.18c. Influence of soil moisture on cumulative number of flowers opened on primary branch 2.





on the main axis in both the treatments and cultivars.

Pod Growth

Effect of soil moisture on pod growth is shown in Figure 1.19. Starting from a lower value, pod weight increased with increasing age. The irrigated plants had increased growth of pods than the rainfed plants at all the stages of growth, except at 10 and 30 days after flowering (DAF) in Sambal and at 10 DAF in Daulat. The irrigated Daulat had significantly higher pod growth at 20, 30, 35, 40 and 45 DAF, whereas the irrigated Sambal had higher pod growth only at 40 and 45 DAF. Daulat had higher pod growth than Sambal.

Seed Growth

Seed weight increased with increasing age at all the stages of growth, except at 45 DAF in rainfed Daulat (Figure 1.20). Soil moisture had no significant effect on seed growth of both the cultivars. The irrigated plants had higher seed weight at 20, 30, 40 and 45 DAF in Sambal and only at 45 DAF in Daulat. Sambal had higher seed weight than Daulat.

Yield and Yield Components, Consumptive Water Use and Water Use Efficiency

Mean values of plant height, length of main axis, number of branches/plant, number of pods on main axis, number of pods/plant, number of seeds/pod, pod weight/plant, straw weight, 1000-seed



Figure 1.19 . Effect of soil moisture on pod growth of two mustard cultivars at successive stages of growth.



Figure 1.20. Effect of soil moisture on seed growth of two mustard cultivars at different stages of growth.

weight, seed-husk weight ratio, harvest index, seed yield, oil content, consumptive water use and water use efficiency of two cultivars as influenced by soil moisture are given in Table 1.12. Soil moisture effect was significant for all the above characters, except the number of branches/plant, the number of pods on main axis seed-husk weight ratio, harvest index and oil content of seeds. Sambal had significantly higher plant height, length of main axis, number of pods on main axis straw weight, 1000-seed weight and water use efficiency than those of Daulat. But the number of seeds/pod and harvest index were significantly higher in Daulat than Sambal .

Plant height at maturity : The irrigated plants were taller than rainfed plants. Sambal was taller than Daulat.

Length of main axis : The main axis of the irrigated plants was longer than that of the rainfed plants. Sambal showed longer main axis than Daulat. Within the irrigated plants, longer main axis was also shown in Sambal. The longest main axis was present in the irrigated Sambal.

Number of branches/plant : The irrigated plants had higher number of branches/plant than the rainfed plants but the difference was not significant. Daulat showed higher number of branches/plant though it was not significantly different from Sambal.

Number of pods on main axis : The number of pods on main axis was higher in the irrigated than that of the rainfed plants but it was not significantly different. Sambal had significantly higher number of pods on main axis. The irrigated Daulat had significantly higher number of pods on main axis than that of the rainfed plants.

Number of pods/plant : The irrigated plants had higher number of pods/plant than the rainfed plants. Significantly higher number of pods/plant was found in the irrigated Daulat. In the rainfed conditions, Sambal had higher number of pods, though the differece was not significant .

Number of seeds/pod : The number of seeds/pod was significantly higher in the irrigated plants than that of the rainfed plants. Daulat had significantly higher number of seeds/pod.

Pod weight/plant : The irrigated plants had significantly higher pod weight than the rainfed plants. The irrigated Daulat had higher pod weight, while the rainfed Daulat had lower pod weight, though the differences were not significant .

Harvest index : Harvest index was higher in the irrigated plants, but it was not significantly different from the rainfed plants. Daulat had significantly higher harvest index than Sambal. Seed yield: Seed yield was significantly higher in the irrigated plants (2240 kg/ha) than that of the rainfed plants (1042 kg/ha). Average yield of Sambal was slightly higher than Daulat but the difference was not significant. In the irrigated condition Daulat yielded more seeds than Sambal but in the rainfed condition it yielded less though these differences were not significant.

Oil content of seeds : Oil content of seeds was not significantly influenced by soil moisture. In the seeds of the irrigated plants, oil content was lower than that of the seeds of the rainfed plants. Daulat yielded higher quantity of oil than Sambal .

Consumptive water use : Consumptive water use of the irrigated plants was significantly higher than the rainfed plants. Daulat had higher consumptive water use in the irrigated condition but lower in the rainfed condition than Sambal, though the differences were not significant.

Water use efficiency : The irrigated plants had significantly lower water use efficiecy (12.04 kg ha⁻¹ mm⁻¹) than the rainfed plants (15.30 kg ha⁻¹ mm⁻¹).Significantly higher water use efficiency was found in Daulat than Sambal.The highest water use efficiency was observed in the rainfed Daulat .

Table 1.12. Mean values of yield and yield components, consumptive water use and water use efficiency of two mustard cultivars as influenced by soil moisture.

.

	Plant heig	ght at matur	ity (cm)	Length of main axis (cm)			Number of branches/plant		
	Sambal	Daulat	Nean	Sambal	Daulat	Kean	Sambal	Daulat	Nean
Irrigated	180	139	160	81	61	71	5.08	5.50	5.29
Rainfed	170	116	143	68	48	58	4.75	4.92	4.83
Nean	175	128		75	55		4.92	5.21	
I.SD at 5%	(a)15.21, (c)32.19,	(b)22.76, (d)22.30		(a)7.14,((c)10.16,	b}7.18, (d)6.94		(a)NS,(b) (c)NS,(d)	NS, NS	

	Number of pods on main axis		Number of pods/plants			Number of seeds/pod			
	Sambal	Daulat	Kean	Sambal	Daulat	Kean	Sambal	Daulat	Kean
Irrigated	59 .	51	55	255	277	266	13.1	14.3	13.7
Rainfed	52	40	46	146	124	135	11.9	12.6	12.3
Kean	51	46		186	201		12.5	13.5	
LSD at 5%	(a)NS,(b)3.78 (c)5.35,(d)10.15			(a)9.94,(b)NS (c)NS,(d)38.18			(a}1.31,{b}0.83 (c)1.18,(d)0.80		

•.

Contd

.

Table	1.12.	(Continued).	

•

	Pod weight/plant (g)			Straw	weight/pla	nt (g)	1000-meed weight(g)		
	Sambal	Daulat	Nean	Sambal	Daulat	Nean	Sambal	Daulat	Nean
Irrigated	12.82	13.66	13.24	15.14	11.23	13.24	2.80	2.30	2.55
Rainfed	6.89	5.74	6.31	8.07	5.03	5.92	2.50	2.16	2.33
Nean	9.86	9.70		11.61	8.13		2.65	2.23	
LSD at 5%	(a)0.2 (c)NS,	5,(b)NS, (d)0.92		(a)1.88, (c)2.33,	(b)1.65, (d)1.59		(a)0.19, (c)0.40,	(b)0.29 (d)0.28	

•

	Seed	-husk weig	ht ratio	iiar	vest index	(X)	Seed yield (kg/ha)			
	Sambal	Daulat	Nean	Sambal	Daulat	Nean	Sambal	Daulat	Kean	
Irrigated	1.05	1.06	1.06	23.44	28.10	25.77	2163	2317	2240	
Rainfed	1.03	0.97	1.00	23.38	26.24	24.81	1151	933	1042	
Nean	1.04	1.02		23.41	27.17		1657	1625		
LSD at 5%	(a)NS,((c)NS,((a)NS,(b)NS, (c)NS,(d)0.04			(a)NS, (b)1.90, (c)2.68, (d)NS			(a)190,(b)NS (c)NS,(d)219		

Contd

.

	01	l content	(%)	Consump	Consumptive water use (mm)			Water use efficiency (kg ha ⁻¹ mm ⁻¹)		
	Sanbal	Daulat	Nean	Sambal	Daulat	Nean	Sambal	Daulat	Kean	
Irrigated	33.33	33.67	33.50	183	189	186	11.82	12.26	12.04	
Rainfed	34 .00 33.67	36.00	35.00	82 133	60 125	71	14.04	16.55 14.41	15.30	
LSD at 5%	(a)NS,((c)NS,(b)NS, d)NS		(a)55, (c)NS,	(b)NS, (d)69		(a)2.65,(b (c)NS,(d)3)2.12, 1.04		

.

Table 1.12. (Continued).

(a) = Difference between treatments

(b) = Difference between cultivars

•.

(c) = Difference within treatment between cultivars

(d) = Difference within cultivar between treatments

Correlation Coefficients Between Yield and Yield Components

Simple correlation coefficients between yield and yield components influenced by soil moisture are shown in Table 1.13.

Association between yield and yield components of the irrigated plants

The number of pods on main axis was positively correlated with the number of branches/plant. Pod weight/plant was positively associated with the number of pods/plant and the number of seeds/ pod. Thousand-seed weight was positively related to plant height and the number of pods on main axis but negatively related with the number of pods/plant and the number of seeds/pod. Harvest index was positively correlated with the number of pods/plant, number of seeds/pod and pod weight/plant but negatively correlated with plant height and 1000-seed weight. Seed yield was positively correlated with the number of pods/plant, number seeds/pod, pod weight/plant and harvest index. Oil content had no significant relation with any of the yield component.

Association between yield and yield components of the rainfed plants.

The number of pods on main axis was positively correlated with plant height, the number of branches/plant, number of pods/plant and pod weight/plant. Pod weight/plant was positively associated with plant height and the number of pods/plant. Thousand-seed weight was positively correlated with plant height. Harvest index had negative association with plant height and positive association with the number of seeds/pod.Seed yield showed positive correlation with plant height, number of pods on main axis, the number of pods/plant, pod weight/plant and harvest index.Oil content of seeds had positive relation with the number of seeds/pod and negative relation with plant height and the number of pods on main axis. Table 1.13. Simple correlation coefficients between yield and yield components (upper diagonal irrigated and lower diagonal rainfed values).

.

.

	Plant height at maturity	Number of branches/ plant	Rumber of pods on main axis	Number of pods/plant	Humber of Beeds/pod	Pod weight/ plant	1000-seed weight	Harvest índex	Seed yield	Oil contenl
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1		-0.01	0.51*	-0.02	-0.21	0.12	0.58**	-0.60**	0.13	-0.35
2	-0.06		-0.46*	0.29	0.05	0.37	-0.06	0.10	0.33	-0.24
3	0.64***	-0.45*		0.11	-0.01	0.14	0.44*	-0.19	0.18	0.10
4	0.34	0.08	0.54*		0.36	0.91***	-0.41*	0.48‡	0.90*1	* -0.04
5	-0.35	0.16	-0.30	-0.03		0.46‡	-0,43*	0.50*	0.48*	-0.02
6	0.42*	-0.03	0.47*	0.82***	0.24		-0.24	0.49‡	0.99**	* -0.11
7	0.50*	-0.22	0.10	-0.15	-0.06	0.23		-0.63**	-0.27	-0.08
8	-0,48‡	0.30	-0.36	0.26	0.63***	0.38	-0.09		0.55**	0.19
9	0.39*	0.06	0.39*	0.79***	0.30	0,98***	0.21	0.48‡		-0.12
10	-0.61**	0.21	-0.61**	-0.28	0.40‡	-0.17	-0.38	0.27	-0.17	

Like any other crop, growth and yield of mustard is under the control of many environmental factors; soil moisture is one of them. In a strict sense mustard is an irrigated crop (Andrews, 1972). Proper maintenance of soil moisture in the crop encourages its growth and development and ultimately reflects to the yield.

Although the use of exponential polynomial equation in growth analysis is advocated (Hurd, 1977; Hunt, 1978; Sivakumar and Shaw, 1978) due to the difficulty of selecting the degree of polynomial to be fitted, Nicholls and Calder (1973) suggested a statistically objective method. However, a serious problem arises when unequal degrees of polynomials describe the relationship between log, dry weight and log, leaf area with time in different sets of data under comparison. The main reason is that this results in strongly dissimilar patterns of derived growth parameters such as CGR, NAR, etc., for which no physiological explanation can be offered. Therefore, Hurd (1977) and Hunt (1978) suggested the use of simplest, quadratic relationship for all cultivars or treatments under comparison, and the same was followed in the present work. A close agreement between the harvest interval and curve-fitting method in the present work, also suggests that the selected polynomial adequately described the data except the dry weight.

The results of the present study showed that total dry matter production was significantly greater in the irrigated plants. Similar result was reported by Khan and Agarwal (1985), Mandal et al.(1986), Singh et al.(1987) and Haque et al. (1987) in mustard, Krogman and Hobbs (1975) and Clarke and Simpson (1978 a) in rape, Nageswara Rao et al.(1985) and Srinivasan et al.(1987) in groundnut and Farah (1981) in field beans. The increase of total dry matter was due to increased LAI in the early stages (Watson, 1947) in the irrigated plants. Total dry matter increased slowly at the early vegetative phases but increased rapidly with the advancement of the growth period (Figure 1.1). The rapid increase of total dry matter at the later stages was due to green stem and pod photosynthesis (Scott et al., 1973). The quadratic polynomials predicted dry weights to drop just prior to maturity, while the actual dry weights continued to increase. It may be attributed that cubic polynomials was most appropriate expression for fitting dry weight data.

Significantly higher LAI was found in the irrigated plants (Figure 1.2). This result corroborates the findings of Mandal *et al.* (1986) in mustard and Clarke and Simpson (1978 a) in rape. Similar result was also reported in several plants other than *Brassica*, like barley (Kirby, 1969), sorghum (Constable and Hearn, 1978; Sivakumar *et al.*, 1979 and Rabindrnath and Shiv Raj, 1983), sunflower (Connor and Jones, 1985) and peanut (Nageswara Rao *et al.*, 1988). The increase of LAI occurred due to increase of the leaf expansion in the irrigated plants. Soil moisture increased turgor pressure in the cells and turgor forces played a part in the process of leaf expansion (Hsiao and Acevedo, 1974). With the increase in the level of irrigation, uptake of nutrients was more hence more expansion of leaf took place (Mandal *et al.*, 1986). Soil moisture increased the

relative leaf water content(RLWC) (Figures 1.16a,b) which increased cell expansion and ultimately leaf area increased. Reduced LAI in the rainfed plants may be caused by the faster rate of leaf senescence and reduced number of leaves. Water stress is known to hasten leaf senescence (Asana, 1961) and reduces the number of leaves (Sivakumar and Shaw, 1978 and Turk et al., 1980). Mare and Palmer (1976) found that the total number of leaves produced by the primary stem of sunflower was reduced due to water stress .This may be due to the inhibition of the initiation and differentiation of leaf primordia due to water stress. Starting from a lower value, LAI reached a certain peak and then declined throughout (Figure 1.2). Senescence and abscission of the older leaves caused the depletion of LAI at the later stages. The pattern of LAI obtained from curvefitting method was similar with that of the original data. This is due to the best fit of the leaf area data in the quadratic polynomials.

The irrigated plants had significantly higher LAD in both the cultivars. Higher LAD of the irrigated plants was due to higher LAI for a longer period. Krogman and Hobbs (1975) found that irrigation of rape slowed the senescence of leaves and thereby increased the total green surface area of the plants. Constable and Hearn (1978) reported that soil water deficit during pod filling in soybeans caused early leaf death. The pattern of LAD calculated from curvefitting. method was closer to that calculated from the original data. The best fitness of leaf area data in the quadratic polynomials is the cause of this similar pattern. Significantly higher CGR was found in the irrigated plants at almost all the stages of growth in the present investigation (Figure 1.4a). Higher CGR in the irrigated plants was also observed by Clarke and Simpson (1978a) in rape and Srinivasan *et al.*(1987) in groundnut. Rabindranath and Shiv Raj (1983) reported that the CGR values reduced considerably with water stress. Higher CGR in the irrigated plants was due to higher production of dry matter owing to higher LAI. CGR calculated from the quadratic-fitted data increased up to a certain peak and decreased thereafter giving uncharacteristically negative values late in the season (Figure 1.4b). CGR is the most meaningful growth function, since it represents the net result of photosynthesis, respiration and canopy area interactions. As noted by Williams *et al.* (1965), CGR is also representative of the most common agronomic measurment, i.e., yield of dry matter per unit land area.

The irrigated plants had higher RGR at several stages of growth in both the cultivars but significantly higher only at (41-48) DAS in Daulat. This result corroborates the finding of El Nadi (1969) and Nerkar *et al.*(1981) in *Vicia faba*. Richards (1978) found that RGR was reduced in more severe drought treatment in rape. RGR in the irrigated plants increased because of the increase of LAR, as LAR is a component of RGR. In the present study, RGR declined with increasing age and plant dry weight.Similar result was reported for RGR in *Dactylis glomerata* (Eagles, 1969; 1971), in rape (Clarke and Simpson, 1979) and in *Pennisetum americanum* (Chanda *et al.*, 1987). It had been suggested that the decrese in RGR could be attributed to selfshading of lower leaves by upper leaves (Thorne, 1961). The decreasing trends in RGR with age was mainly due to the decline in LAR (Chanda *et al.*, 1987). The pattern of RGR was similar in both the curve-fitting and the classical methods.

At the early stages NAR values were slightly higher in the irrigated plants, but at the later stages it was significantly higher in the rainfed plants. Due to higher LAI, NAR was lower in the irrigated plants (Watson, 1952; Donald, 1961; Harper, 1963 and Rees, 1963). Watson (1958) has attributed the decline in NAR to increased mutual shading of leaves, resulting in reduced photosynthesis. Higher NAR due to higher soil moisture was found by El Nadi (1969) and Nerkar et al. (1981) in beans and Rabindranath and Shiv Raj (1983) in sorghum. Soil moisture effect in the present study (Figures 1.6a and 1.6b) showed that very little increase or depletion of NAR was found at the early stages but at the later stages it increased very sharply. A similar occurence was evident in Brassica napus(Allen and Morgan, 1972, 1975) and in soybean (Buttery, 1969; Koller et al., 1970). The increase of NAR at the later stages may be due to decreased leaf area and increased stem and pod photosynthesis. The upswing in NAR occurred after the cessation of flowering, and after maximum pod area would have occurred (Tayo and Morgan, 1975). Inanga and Kumara (1974) reported that maximum photosynthetic rate of rape occurred before the establishment of maximum pod area, after which the photosynthetic rate of all plant parts

declined. As a result, the observed upswing in NAR can be attributed to a sudden increase in photosynthetic contribution by pods. Koller *et al.*(1970) suggested that the observed upswing in NAR of soybean was the response of the photosynthetic apparatus to increased demand for assimilates by the rapidly growing seeds. Results of Thorne and Koller (1974) have shown that assimilate demand has a marked influence on source-leaf photosynthesis, and formation and export of carbohydrates in soybean. In *Brassica napus*, there is a rapid increase in seed weight late in the ripening phase (Norton and Harris, 1975); this effect would have provided the high sink demand which in turn increased photosynthetic activity. The pattern of NAR calculated from curve-fitting method showed a close correspondence with that calculated from the original data (Figures 1.6a and 1.6b).

The irrigated plants had higher LAR values at most of the stages of growth, but in Daulat it was significantly lower at 20 DAS. In Vicia faba, Nerkar et al. (1981) found that LAR increased with the increased level of soil moisture. Higher LAR in the irrigated plants may be due to increased SLA and LWR, since SLA and LWR are the two components of LAR. LAR values declined sharply with plant age in both the treatments and cultivars. Wallace and Munger (1965) reported that in dry beans LAR was highest during the early vegetative stage and later decreased rapidly with advancement of age. The pattern of curve-fitted LAR was similar with that of the original data.

Significantly higher RLGR was found in the irrigated plants of Sambal at (69-76) DAS and of Daulat at (27-34) and (48-55) DAS. Significantly lower RLGR was found at (55-62) DAS in Sambal and at (69-76) DAS in Daulat. Richards (1978) reported that RLGR was not affected by drought stress in rape. The maximum RLGR was found at (20-27) DAS in both the cultivars and treatments and then declined with increasing age. Similar decline of RLGR was found in sovbean (Buttery, 1969); in blackgram (Pandey et al., 1978) and in pearl millet (Chanda et al., 1987). Much of this decline can be attributed to self-shading at the early stages and abscission and senescence of leaves at the later stages. Saha (1983) observed a positive relation between RLGR and LAR in jute. So, due to decline of LAR, RLGR would have been declined. RLGR values obtained from the curve-fitting method drifted with increasing plant age and also with increasing plant weight (Figures 1.8b and 1.12).

The irrigated plants had significantly higher SLA at 34, 48, 55 and 62 DAS in Sambal and at 34 and 55 DAS in Daulat (Figure 1.9a). Higher SLA in the irrigated plants may be attributed due to less number of cells per unit leaf area and lower compactness between cells. SLA increased slightly at the later stages of growth in all the treatments and cultivars except that in the irrigated Daulat. The increase of SLA at the later stages may be due to translocation of assimilates from the leaves to the sink organs. Chanda *et al.* (1987) observed that SLA values decreased with increasing plant dry weight in pearl millet. In the present experiment, the LWR values of the irrigated plants were significantly greater at 48 and 55 DAS in Sambal and 55, 62, 69 and 76 DAS in Daulat (Figure 1.10a). The increase of LWR in the irrigated plants may be attributed due to increased leaf weight. LWR values declined very sharply with increasing plant age and plant dry weight. Strong drift of LWR with increasing plant dry weight was reported by Chanda *et al.* (1987) in pearl millet. The decrease of LWR was caused by increased plant dry weight and decreased LAI at the later stages.

Correlation analysis indicated that the pre-flowering CGR was positively correlated with pre- flowering LAI and negatively correlated with RGR and NAR at the pre-flowering stage of the rainfed plants (Table 1.4). This indicated that soil moisture had a direct effect on the association between CGR and LAI of the pre-flowering stage. CGR at the post-flowering stage of the rainfed plants showed positive correlation with the post-flowering LAI and negatively correlated with NAR (post-flowering) and LAR (pre-flowering).Pandey et al.(1981) observed a positive relation between CGR and LAI in cowpea. Post-flowering RGR (irrigated) was negatively correlated with LAT at the pre-flowering stage. This indicates that RGR decreased with increasing LAI at the pre-flowering stage of the irrigated plants. NAR (pre-flowering) was positively correlated with LAI at pre-flowering stage in the irrigated plants. NAR (postflowering) was positively correlated with LAR (pre-flowering) and negatively correlated with LAR at post-flowering in both the treatments. Post-flowering NAR of the rainfed plants was also

correlated negatively with the post-flowering LAI. This relationship postulates that with decreasing LAI in the rainfed plants NAR increased at the post-flowering stage. Thurling (1974) and Tan etn/.(1978) reported that NAR was negatively correlated with LAR and LAT. Seed yield was positively correlated with LAI at the preflowering stage and CGR and NAR at the post-flowering stage in irrigated plants. But in the rainfed plants, seed yield the was positively correlated with LAI and CGR at the post-flowering stage and negatively correlated with NAR at the post-flowering and LAR at the pre-flowering stages (Table 1.4). Seed yield was found significantly correlated with LAI in mustard (Reddy and Sinha, 1987 and Chaturvedi et al., 1988) and with RGR, NAR and LWR in B. campestris and with LAI and LWR in B. napus (Thurling, 1974). In other crops also, the growth attributes like LAI and LAD had positive correlation with yield (Thorne, 1960; Saini and Das, 1979).

Developmental factors affecting the accumulation of dry matter and subsequent partitioning of assimilates are of great importance in determining the final yield in crops (Watson, 1971; Wareing and Patric, 1975). In the present investigation, the percentage of DM accumulation for stem + petiole was lower at the early stages, at the middle stages it increased and then gradually decreased with time (Figures 1.14 and 1.15). Percentage of DM accumulation for stem + petiols was lower in the irrigated plants. DM accumulation in the leaves was found highest at the initial stage and gradually declined with increasing time and at 83 DAS it became zero. The irrigated plants accumulated higher percentage of DM for leaves. The irrigated plants had higher percentage of DM for buds + flowers. Pods accumulated DM first at 48 DAS and gradually increased with age. The rainfed plants had higher accumulation of DM for pods. Similar result were reported in rape (Krogman and Hobbs, 1975); in sovbean and sorghum (Constable and Hearn, 1978); in Vigna radiata (Nagarajah and Schulze, 1983) and in groundnut (Srinivasan et al., 1978). Allen et al. (1971) suggested that pods of rape produce the assimilates for their own development and for the seeds they contain and that the effect of leaves in seed growth is minor and indirect. However, Hozyo et al.(1972) and Freyman et al.(1973), who traced assimilates from the leaves to the seeds, showed that leaves do contribute to seed formation. Soil drought reduced partitioning into leaves but increased partitioning into stems (Nagarajah and Schulze, 1983). In sorghum and soybeans, the significant loss in stem dry weight during grain filling suggests that there could have been reallocation of dry matter from the stem to the developing grain (Constable and Hearn, 1978).

In the present study, the irrigated plants had higher RLWC than that of the rainfed plants. Similar result was reported by Raja and Bishnoi (1990) in rape and Joshi *et al.*(1988) in groundnut. Mehrotra *et al.* (1969) found that with increased soil moisture relative turgidity increased. Sivakumar *et al.*(1979) observed less negative leaf water potential in the irrigated plants than that of the rainfed sorghum. Average relative turgidity attained the minimum value at higher soil moisture tension and the maximum value at lower soil moisture tension (Nayak *et al.*,1983). The higher RLWC was associated with higher dry matter production rates of the irrigated plants because cell turgidity is important in relation to the opening and closing of stomata, expansion of leaves and flowers and movement of water and nutrients to various parts of the plant (Kramer,1969). Wein *et al.*(1979) reported that the water potential decreased with increasing soil moisture stress in *Vigna*. They suggested a increase in the plant resistance to water flow as soil moisture stress increased, which would help in maintaining suitable plant water status. But Hall and Schulze (1980) exhibited that the plant resistance to water flow definitely increased in *Vigna* with soil moisture stress. In the morning and afternoon, RLWC remaind higher but at noon it decreased (Figures 16a and 16b). The decrease of RLWC at the midday may be due to higher evapotranspiration owing to increased temperature and light intensity.

Chloroghyll a and b are the most important pigments active in the photosynthetic processes. In the present experiment, chlorophyll a,b and total chlorophyll slightly increased in the irrigated plants at the mid-pod filling stage, but the difference was not significant (Table 1.5). Hussein (1973) reported that water regime had no significant effect on the concentration of total chlorophyll in the leaves of cotton. There was an increase in the percentage of chlorophyll in the leaves of cotton with decreasing water stress at different periods of plant growth (El-Fouly *et al.*,1971; El-Saidi and El-Mossallami,1972). There was an increasing concentration of total chlorophyll, chlorophyll a and b in the leaves of cotton with a decreasing in the amount of available moisture depletion before irrigation at the early stages of growth and at the later stages of growth chlorophylls were highest at the 70% available soil moisture before irrigation (Moursi *et al.*,1976).

The rainfed plants accumulated significantly higher amount of proline than the irrigated plants at both the flowering and mid-pod filling stages. Proline content increased at the mid-pod filling stage than the flowering stage (Table 1.5). Increased proline accumulation in the stressed plants was found in sorghum (Blum and Ebercon, 1976), in wheat (Quarrie, 1980) and in foxtail millet (Narasimha Rao and Shiv Raj, 1985). But no relationship was observed by Richards (1978) in rape between the amount of proline accumulated and the water status of the leaves, nor between the proline content in turgid and wilted leaves of the same cultivar. In sorghum, leaves of water-stressed field-grown plants accumulated proline to a level of several times greater than that in non-stressed plants (Waldren et al., 1974). Stewart et al. (1966) and Blum and Ebercon (1976) suggested that the accumulated proline could also serve as a readily available energy and nitrogen source for use upon the relief of stress. Accumulated proline may be oxidized and serve as a source of energy, especially when carbohydrate content is low (Wang, 1968; Oakset al., 1970 and Stewart, 1972). However Hanson (1979) suggested that the free proline accumulated in response to inert and played no cellular role. Previous water stress was conjectures have projected proline: (a) as a detoxifying agent, rendering harmless NH3 released by proteolysis (Kemble et al.,

1954), but experimental evidence has discounted this role (Blum and Ebercon, 1976); (b) as a temporary storage compound for N, providing a major and readily available source of energy and N for post-stress recovery (Stewart *et al.*, 1966; Blum and Ebercon, 1976); and (c) as a hydrophilous agent for maintaining the colloidal properties of cell protoplasm against the extremes of dehydration (Protsenko *et al.*, 1968). More recent evidence suggests that proline may confer protection on some mitochondrial and solubilized enzymes against heat instability (Nash *et al.*, 1980). Beneficial as they may be, these proven and unproven cellular role of proline are usually to satisfactorily account for the agronomic expression of drought resistance.

Significantly higher sugar accumulation was found in the rainfed plants at both the flowering and mid-pod filling stages (Table 1.5).Sugar content increased at the mid-pod filling stage than the flowering stage.Stewart (1971) and Narasimha Rao and Shiv Raj (1985) reported that sugar accumulation increased as a result of water stress. The soluble sugars were significantly higher in both the unirrigated sunflower and sorghum plots than in the irrigated plots (Turner *et al.*, 1978). Easten and Ergle (1948) suggested that starch would be converted into sugars under stress. Sugars are also known to stabilize protoplasmic membranes (Larson, 1975).

Relative leaf water content (RLWC) at which stomata closed was not significantly affected by soil moisture treatment (Table
1.6). Similar result was reported by Nazrul-Islam and Yasmin (1982) in tomato. Bannister (1964) has shown that Calluna vulgaris from dry sites had a lower relative turgidity at stomatal closure than those from wet site. The water potential for stomatal closure was significantly correlated with the osmotic potential of the leaves at stomatal closure, i.e., as the osmotic potential of the leaves decreased the leaf water potential for stomatal closure decreased (Turner et al., 1978). The irrigated plants took significantly more time to close their stomata at the flowering stage (Table 1.6). Stomata of the plants of moist sites took more time to close than that of the dry site (Nazrul-Islam and Yasmin, 1982). In sunflower and sorghum, Turner et al. (1978) observed that time for stomatal closure was greater in the irrigated plants than that of the unirrigated plants. In the present investigation, Sambal closed its stomata well before Daulat. As the quickness of stomatal closure is a means of drought tolerance, Sambal is more resistant to drought.

At the flowering stage, total transpiration rate was signficantly higher in the irrigated plants. Stomatal and cuticular transpiration rates were not affected significantly by soil moisture treatment.Significant effect of soil moisture was found on total and stomatal transpiration by Nazrul-Islam and Yasmin (1982) in tomato and Nazrul-Islam and Alam (1986) in jute. Joshi *et al.*(1988) observed higher transpiration rate with stress-free plants than that of the stressed-plants of groundnut. Transpiration rates were similar in the irrigated and the unirrigated groundnut (Black *et al.*,1985).

Stomatal characters are known to be associated with CO2exchange rates (Dornhoff and Shibles, 1976). Soil moisture had no significant effect on stomatal frequency and pore length on both the surfaces of leaves. The epidermal cells decreased and stomatal index increased significantly on the abaxial surface of the irrigaleaves. Although soil moisture had no significant effect, ted stomatal frequency of both the surfaces decreased in the irrigated plants.Similar result was noted by Ciha and Brun (1975)in soybeans. Penfound (1931) reported fewer stomata/unit area when plants were grown under optimum soil moisture conditions compared with plants grown under water stress conditions. The decrease in stomatal frequency with water supply suggests that increased leaf expansion due to enlargement of cells caused this decrease. A greater stomatal frequency was observed on the abaxial surface than the adaxial surface. Similar results were also observed in Vicia faba (Nerkar et al., 1981), in soybean (Miller, 1938; Dornhoff, 1971; Teare and Kanesasu, 1972; Ciha and Brun, 1975).

Thickness of leaf was increased significantly in the irrigated plants (Table 1.9). Mean cross-sectional areas of palisade and spongy parenchyma were higher in the irrigated plants than the rainfed plants but the variations were not significant.Begum (1991) found that leaf thickness, mean cross-sectional area of palisade and spongy parenchyma increased with irrigation in mustard. The increase of leaf thickness may be due to increased cell size of the plants. Due to increased RLWC in the irrigated plants, cross-sectional area of mesophyll cells increased. Plant height at first flowering and duration of flowering were significantly increased and the number of leaves at first flowering and days taken to first flowering were unaffected by soil moisture (Table 1.10). Increased height of stem may be caused for increased elongation of the cells due to higher turgidity (Allen *et al.*, 1976; Vivekanandan and Gunasena, 1976). Longer duration of flowering for the irrigated plants may be due to delayed termination of flowering. This delaying may be caused as higher number of flower buds were initiated in these plants which took longer time to open.

Irrigated plants had always higher cumulative number of flowers on the main axis and the primary branches 1-3 than the rainfed plants (Figures 1.8a-d). In the primary branch 3, duration of flowering was shorter for the rainfed plants. The initiation and differentiation of reproductive primordia and further development of the differentiated cells are sensitive to moisture stress (Slatyer, 1973). Moisture shortage at final differentiation may result in defective floral organs, sterility, reduced number of flowers and failure of flowers to open (Slatyer, 1973).

The percentage of pod abortion was higher in the primary branches of the rainfed plants, but the difference was not significant (Table 1.11). The cause of pod abortion may be due to shedding of flowers. Some flower-shedding may be due to competition between a large number of flowers and metabolic products and root activity in relation to salt and water absorption may be retarded during the flowering phase (Lochwing, 1940).El Nadi (1969) reported that a wet regime during the flowering phase significantly reduced the intensity of flower shedding, but a dry treatment during this stage, the incidence of flower shedding was aggravated to a considarable extent. It is likely that the efficiency of water transport and mineral uptake is reduced during flowering and this is aggravated under conditions of water stress.

The irrigated plants showed higher pod growth rate. Kaufmann (1972) stated that reduction in fruit size was caused by water deficit during the period of fruit enlargement. Pod growth rate and the yield of seeds are not simple functions of leaf area index and that sources of materials other than the leaves are important in affecting the yield of pods and seeds (Milbourn and Hardwick, 1968).

In the present investigation, higher rate of seed growth was observed in the irrigated plants at 20, 30, 40 and 45 DAF in Sambal and at 45 DAF in Daulat, but the defference was non-significant (Figure 1.20).The increase in dry weight of grains was significantly affected under water stress in sorgham (Narasimha Rao and Shiv Raj,1988). Wardlaw (1967, 1968, 1971) concluded that reduction in grain yield under stress was not due to effects on photosynthetic rate but due to vein loading. However, Begg and Turner (1976) concluded that reduction in translocation caused by water stress was due to reduced photosynthesis or sink growth but not due to effect on conducting system. Plant height at maturity, length of main axis, number of pods/plant, number of seeds/pod, pod weight/plant, straw weight, 1000-seed weight and seed yield were significantly higher in the irrigated plants (Table 1.12). Similar results were reported by some other workers in mustard and rape. Signifiantly higher plant height, number of branches/plant, number of pods/plant and number of seeds/pod were found in mustard (Joarder *et al.*,1979; Bajpai *et al.*,1981; Singh *et al.*,1987 and Reddy and Sinha, 1987) and in rape (Maini *et al.*,1964; Krogman and Hobbs,1975; Clarke and Simpson ,1978 b and Bhan *et al.*,1980). But Bhati and Rathor (1982) and Chaniara and Damor (1982) found no significant effect of soil moisture on plant height, number of branches/plant and 1000-seed weight in mustard.

Seed yield was significanly higher in the irrigated plants. Similar result was observed by Mathur and Tomar (1972), Joarder et al.(1979), Bhan (1979), Chaniara and Damor (1982), Mandal et al. (1986), Singh et al. (1987), Reddy and Sinha (1978) and Reddy et al. (1988) in mustard and by Maini et al. (1964), Banerjee et al. (1967), Clarke and Simpson(1978b) and Bhan et al.(1980) in rape. No significant effect of soil moisture on yield was found in mustard by Bajpai et al.(1981) and Bhati and Rathor (1982) and in rape by Wankhede et al.(1970). Soil moisture had no significant effect on oil content of seeds (Table 1.12). Similar result was found in mustard(Chaniara and Damor, 1982 and Bhati and Rathor, 1982) and in rape (Wankhede et al., 1970). Oil content of seeds was higher in the irrigated plants in rape (Bhan et al., 1980) and in groundnut (Mehrotra et al., 1969). There was a slight increase in oil content up to one irrigation and then declined a little with increasing number of irrigations (Banerjee et al., 1967).

Consumptive water use increased and water use efficiency decreased with irrigation (Table 1.12). Similar relult was also reported by Singh and Singh (1978), Bhan (1981) and Reddy *et al.* (1988). The lower water use efficiency with higher soil moisture status was due to proportionately more increase in evapotranspiration than the increase in seed yield. The increased frequency of irrigation increased the amount of moisture use as well as water use efficiency in rape (Bhan *et al.*, 1980).

The correlation coefficients showed that 1000-seed weight was positively correlated with the number of pods on main branch and negatively correlated with the number of pods/plant and the number of seeds/pod in the irrigated plants (Table 1.13). This indicates that 1000-seed weight is increased with increasing number of pods on main branch but the relation is reverse in case of the number of pods/plant and the number of seeds/pod. Clarke and Simpson (1978b) found a negative correlation of 1000-seed weight with the number of pods/plant and the number of seeds/pod. Irrigation affected 1000seed weight, probably because of a combination of their effects on assimilate supply and distribution in the plants.

Seed yield in the present investigation was positively correlated with the number of pods/plant, pod weight/plant and harvest index in both the treatments and number of seeds/pod in the irrigated plants only (Table 1.13). Clarke and Simpson (1978b) reported that seed yield in the high irrigated plants of *Brassica napus* was positively associated with 1000-seed weight and negatively related with the branches/plant and the number of seeds/pod. Seed yield was positively correlated with the number of pods/plant, the number of seeds/pod, the number of primary branches/plant and pod weight/plant (Reddy and Sinha, 1987). Oil content of the rainfed plants was positively associated with the number of seeds/pod and negatively with plant height and the number of pods on main axis but no relation was observed in the irrigated plants.

The overall results of the present experiment indicate that TDM, LAI, LAD and some growth attributes like CGR, LAR, RLGR, SLA and LWR were increased significantly by soil moisture. Seed yield was correlated with LAI, CGR and NAR in the irrigated plants. Proline and sugar contents were significantly higher in the rainfed plants and increased with increasing age. The irrigated plants had higher leaf thickness, plant height at first flowering, duration of flowering and number of flowers. Yield and yield components like plant height at maturity, length of main axis, number of pods/plant, number of seeds/pod,pod weight/plant,straw weight, 1000-seed weight and seed yield were found significantly higher in the irrigated plants.Seed yield in the irrigated plants was positively correlated with the number of pods/plant, pod weight/plant, number of seeds/ pod and harvest index. Therefore, optimum soil moisture treatment be recommended and selection should be based on 1 may

characters for better yield. Proline and sugar accumulation and seed yield of Sambal were higher in the rainfed condition, whereas these characters of Daulat were higher in the irrigated condition. Thus, Daulat is more susceptible to water stress than Sambal. Therefore, to get higher yield, Sambal may be recommended for cultivation in the rainfed condition and Daulat in the irrigated condition.

EXPERIMENT 2

EFFECT OF SOIL MOISTURE AT SPECIFIC STAGES OF GROWTH ON DRY MATTER PRODUCTION AND YIELD OF MUSTARD (*BRASSICA JUNCEA* L.)

.

INTRODUCTION

Mustard is usually grown as a rainfed crop in Bangladesh. Soil moisture deficits frequently occur under rainfed condition; where the crop has to depend upon the stored soil moisture which is soon depleted through evapotranspiration causing depression in growth and yield of the crop. Plants fail to recover if permanent wilting occurs due to severe moisture stress. When rewatered after a prolonged and severe moisture stress plant will recover its growth. However, yield of crops may be depressed due to soil moisture stress of any degree. The rate of yield reduction will depend on the degree of stress.

Plants are not equally sensitive to soil moisture deficits at various stages of growth. There are some specific stages of plants during which they are more prone to soil moisture deficit. The vegetative stage of many plants are not sensitive to soil moisture stress. This may be due to the reason that plants resume their growth after releasing stress. Soil moisture stress at the flower initiation and flowering stage reduces the yield of legumes through the formation of defective floral organs, sterility, reduced number of flowers and failure of flowers to open (Slatyer, 1973). Moisture stress during the period of fruit development reduces the yield through the reduction of pod and seed size (Kaufmann, 1972). This is caused due to insufficient transportation of nutrient reserves into the pod and seed. Soil moisture stress at the stage of maturity is less harmful to the yield of crops as it is the stage of dehydration. Information is limited about the effects of water stress at different growth phases on crop growth and development of mustard. Due to an increase in the frequency of drought, declining water table, and increased costs of irrigation water, it is essential to evaluate water management of mustard. In spite of the reputed drought tolerance of mustard, information on the growth phases most sensitive to moisture stress is inadequate.

The aim of this study was to examine the effects of soil moisture stress imposed at specific growth phases on the growth, development and yield of two mustard cultivars.

.

MATERIALS AND METHODS

The plant material for the experiment was also Sambal and Daulat of mustard (*Brassica juncea* L.)

The experiment was arranged in a randomized complete block design with two replications. The size of each block was 2.1 m x 1 m. The distance between two blocks was 1 m. There were 8 rows in each block of 1 m long and 30 cm apart. The distance between plants within a row was approximately 10 cm. Seeds were sown on 2 Nvember, 1990. There were 5 irrigation treatments as follows :

- $T_0 = No irrigation$.
- $T_1 = No$ irrigation until the start of flowering, uniformly irrigation thereafter.
- T₂ = Irrigation until the start of flowering followed by no irrigation until mid-pod filling stage, uniformly irrigation thereafter.
 - T₃ = Irrigation until mid-pod filling stage followed by no irrigation until crop harvest.
 - $T_i = Uniformly irrigation throughout.$

The measured amount of water (10 mm on each occasion) was uniformly added with sprinklers. The treatments T_1 , T_2 , T_3 and T_4 of Sambal received 50, 70, 50 and 80 mm of irrigation water, respectively, whereas the corresponding treatments of Daulat received 40, 40, 50 and 70 mm of irrigation water. The similar cultural practices were done as in the experiment 1.

Dry Matter Determination

For dry matter determination, 5 harvests were taken at equal

intervals of 7 days. The first harvest was taken at 59 DAS. Four plants were harvested from each treatment of each replication of each cultivar. At each harvest, plants were cut off just below the cotyledonary node and tops were separated into leaves, stem and others (petiole + buds + flowers) and pods. Before weighing, all plant parts were dried separately in an oven at about 85 °C for 24 hours till they reached constant weight.

Yield and Yield Components

The final harvest was done at 100 DAS for Sambal and at 95 DAS for Daulat. Six plants per replication per treatment per cultivar were harvested and dried in the sun and the measurements of yield and yield contributing characters were taken. The following characters were recorded :

Plant height at maturity, length of main axis, number of branches/plant, number of pods on main axis, number of pods/plant, number of seeds/pod, pod weight/plant, straw weight, 1000-seed weight, seed-husk weight ratio, harvest index, seed yield/ha and oil content of seeds.

Statistical Analysis of Data

The experimenatal design was a randomized complete block design and the analysis of variance was done accordingly.

RESULTS

Total Dry Matter

Mean squares from the analysis of variance for total dry matter (TDM) as influenced by soil moisture stress at specific stages of growth are presented in Table 2.1. Soil moisture stress effect was significant at all the growth periods except 73 DAS. Cultivar x soil moisture interaction was non-significant.

The effect of soil moisture stress at specific stages of growth on dry matter is shown in Figure 2.1. In Sambal, decrease of TDM in the stressed plants over stress-free plants (T_4) was observed. The highest reduction of TDM production was found in continuous stressed (T_0) and the second and the third reduced position was found in the flowering to mid-pod filling stress (T_2) and pre-flowering stress (T_1) , respectively. The lowest reduction of TDM was found in T_3 , where water stress was imposed after midpod filling stage.

In Daulat, all the plants except T_3 yielded reduced TDM than the stress-free plants (T_4) . Maximum reduction of TDM production was found in the plants grown under continuous stress (T_0) . The socond position of reduced production was found in T_2 at 59, 66 and 87 DAS. Post mid-pod filling stressed plants (T_3) produced increased amount of TDM than that of the stress-free plants (T_4) , though the difference was not significant.

- .

Table 2.1. Mean squares from the analysis of variance of total dry matter (TDM) of two mustard cultivars as influenced by soil moisture stress at specific stages of growth.

.

.

Sources of	đf	Mean squares								
variation			Days	after sowing						
		59	66	73	80	87				
Replication	1	6444	1730	2761	16075	31				
Treatment (T)	4	15279*	20804**	20623	35783**	72485***				
Cultivar (C)	1	7881	7	551	238	2398				
тхс	4	2695	2537	7847	3580	8098				
Error	9	2709	2504	6445	4692	5298				



Figure 2.1. Effect of soil moisture stress at specific stages of growth of two mustard cultivars on total dry matter (TDM).

Distribution of Dry Matter

Influence of soil moisture stress at specific stages of growth of two cultivars on distribution of dry matter (DM) is shown in Figure 2.2. Percentage of DM of stem and others (petiole, buds and flowers), leaves and pods are shown in these figures. No significant effect of soil moisture stress on dry matter distribution was found. Percentage of DM accumulation in stem and others and leaves decreased with increasing age in both the cultivars. But in pods, DM accumulation increased with increasing age. Dry matter percentage of stem and others was highest in T_n and lowest in T_4 at the initial stages but reverse was found at the later stages. The highest percentage of DM accumulation in leaves of Sambal was found in T_1 . The lowest DM accumulation in leaves of Daulat was in T₀. The highest percentage of DM in pods was found in T_1 at 87, in T_2 at 59, in T_3 at 80 and T_4 at 73 and 87 DAS in Daulat.

Yield and Yield Components

Mean squares from the analysis of variance of plant height at maturity, length of main axis, number of branches/plant, number of pods on main axis, number of pods/plant, number of seeds/pod, pod weight/plant, straw weight, 1000-seed weight, seed-husk weight ratio, seed yield and oil content of seeds of Sambal and Daulat as influenced by soil moisture stress at specific stages of growth are given in Table 2.2. Soil moisture effect was significant for all the characters except 1000-seed weight, seed-husk weight ratio and



Figure 2.2. Influence of soil moisture stress at specific stages of growth of two mustard cultivars on distribution of dry matter (DM).

Table 2.2. Mean squares from the analysis of variance of yield and yield components of two mustard cultivars as influenced by soil moisture stress at specific stages of growth.

Sources of	dſ			Kean	Squares			
variation .		Plant beight at maturity (cm)	Length of main axis {cm}	Number of branches/ plant	Rumber of pods on main axis	Namber of pods/plant	Ru∎ber of seeds/pod	Pod weight/ plant(g)
Replication	1	9.80	24.20	0.01	3.20	5951	0.58	32.82**
freat∎ent (T)	4	208.83*	55.18*	1.71*	48.80**	25374***	3.15*	86.09***
Cultivar (C)	1	9331.20***	1036.80**	* 0.95**	352.80***	2808*	25.54***	2.95
f x C	4	58.33	34.18	0.22	8.05	1275	0.15	0.82
Brror	9	55.58	12.76	0.24	7.42	1009	0.57	4.32

Sources of	df	Nean squares								
Varlation		Straw weight /plant (g)	1000-seed weight (g)	Seed-husk weight ratio	Harvest index(I)	Seed yield (kg/ha)	Oil content of seeds(%)			
Repication	1	7.47**	0.07	0.0012	7.28	850000*	1.25			
Treat∎ent (T)	4	59,30***	0.08	0.0018	213.01	2502250*	15.68**			
Cultivar (C)	1	82.34***	0.08	0.0912‡	0.73	413000	22.05**			
T x C	4	6.89**	0.22	0.0023	1.65	14500	9.68*			
grior	9	0.56	0.11	0.0141	3.46	137749	2.25			

harvest index. Cultivar differences were significant for all the characters except pod weight/plant, 1000-seed weight, harvest index and seed yield. Cultivar x treatment interaction effect was significant for straw weight and oil content of seeds only.

Mean values of seed yield and its components of two mustard cultivars as influenced by soil moisture stress at specific stages are given in Table 2.3.

Plant height at maturity : Plant height at maturity was reduced in the stressed plants than that of the stress-free (T_{i}) plants. The continously stressed (T_{i}) plants had significantly shorter height than the T_{i} plants. Post mid-pod filling stressed plants (T_{i}) was taller than the stress-free plants but the difference was non-significant. Sambal was taller than Daulat. Plants grown under continuous (T_{i}) and pre-flowering stress (T_{i}) had significantly reduced plant height over the stress-free plants of Sambal.

Length of main axis : Length of main axis of the stressed plants was reduced over the stress-free plants, except in T_3 where it increased slighty. Significantly reduced main axis length was found in the continuous stressed (T_0) plants. Length of main axis was significantly higher in Sambal. Sambal had significantly reduced main axis length in T_0 and T_1 than that of the T_4 plants. Slight increase of main axis length was found in T_1 , T_2 and T_3 over T_4 of Daulat but the differences were not significant. Number of branches/plant : The stressed plants had reduced number of branches over the stress-free plants. Plants grown under T_0 , T_1 , T_2 and T_3 had 20, 13, 12 and 5% reduced number of branches, respectively over the stress-free plants. Significant reduction of the number of branches was found in T_0 and T_1 . Daulat had higher number of branches than Sambal. In Sambal, T_0 and T_2 had significantly reduced number of branches/plant over the stress-free plants.

Number of pods/plant : The stressed plants yielded reduced number of pods/plant than that of the stress-free plants, except T_3 where the number of pods/plant was increased slightly. The number of pods/plant was significantly reduced in T_0 (47%), T_1 (30%) and T_2 (33%) over T_4 . Daulat had higher number of pods/plant than Sambal, but the difference was not significant. Both Sambal and Daulat had significantly reduced number of pods/plant in T_0 , T_1 and T_2 over T_4 .

Pod weight/plant : Pod weight of the stressed plants diminished in respect to stress-free plants. Plants grown under T_0 (57%), T_1 (42%) and T_2 (36%) had significantly decreased pod yield than that of the stress-free plants (T_4). Daulat had higher pod weight/plant than Sambal, but the difference was non-significant.

Straw weight : Reduction of straw weight in the stressed plants was found over the stress-free plants. Straw weight reduced significantly in T_0 (57%), T_1 (38%), T_2 (36%) and T_3 (9%) than that

of the stress-free plants. Sambal had significnatly reduced straw weight in T_0 , T_1 , T_2 and T_3 and Daulat in T_0 , T_1 and T_2 .

1000-seed weight : No Significant effect of soil moisture stress on 1000-seed weight was found.

Seed-husk weight ratio : Seed-husk weight ratio was not significantly affected by soil moisture stress. The ratio was significantly higher in Daulat than Sambal.

Harvest index : Significantly higher harvest index was found in Daulat than that of Sambal. Harvest index increased slightly in the stressed plants of Sambal, though it was not significantly different.

Seed yield : Seed yield of the stressed plants was reduced than that of the stress-free plants. Significantly reduced seed yield was found in plants grown under T_0 (59%), T_1 (59%) and T_2 (37%) over that of T_4 plants. Seed yield was non-significantly higher in T_3 (8%) than T_4 . Seed yield in Daulat was higher than Sambal, but the difference was not significant. In Sambal, the reduction of seed yield in T_0 , T_1 , T_2 and T_3 over T_4 was 58, 37, 38 and 12%, respectively. The corresponding values of Daulat were 61, 37, 33 and 4%.

Oil content of seeds : Oil content of seeds of the stressed plants increased than the stress-free plants (T_i) except in

continuous stress treatment (T_0) . Significantly increased oil content was found in T_1 (9%) and T_2 (12%). Seeds of Daulat had significantly higher percentage of oil than that of Sambal. The increase of oil content of Sambal in T_0 , $T_{1,}$ T2 and T3 over T_4 was 3, 8, 5, and 6% respectively and of Daulat in T_1 and T_2 was 9 and 18%.

Correlation Coefficients Between Seed Yield and Its Components

Simple correlation coefficients between seed yield and its components as influenced by soil moisture stress are given in Table 2.4. Seed yield was positively correlated with the number of pods/plant, pod weight/plant and harvest index in T_0 , with the number of branches/plant, number of pods on main axis, number of pods/plant, pod weight/plant and harvest index in T_1 , with the number of pods/plant, pod weight/plant and harvest index in T_1 , with the number of pods/plant, pod weight/plant and harvest index in T_2 and T_3 and with the number of pods/plant, number of seeds/pod and harvest index in T_4 . There was no significant correlation between oil content and seed yield in any of the treatment.

Table 2.3.	Mean values of	yield and yield	components	of two mus	stard
	cultivars as	influenced by	soil moist	ture stres	s at
	specific stage	es of growth.			

	Plant heig	Plant height at maturity (cm)			main axis (c	ı)	Number of	Rumber of branches/plant		
	Sambal	Paulat	Nean	Sambal	Danlat	Nean	Saubal	Danlat	Nean	
ſŋ	157.50	118.50	138,00	79.50	56.50	63.50	4.75	5.42	5.06	
Т ₁	164.00	129.50	146.75	71.50	64.00	67.75	5.33	5.67	5.50	
۶,	172.00	128.50	150.25	74.50	62.50	68.50	4.92	6.25	5.59	
۲,	178.50	134.50	156,50	B1.00	66.00	73.50	6.00	6.48	6.04	
۲ ₁	181.50	126.50	154.00	82.05	59.00	70.75	6.08	£.59	6.34	
Negp	170.70	127.50		76.00	61.60		5.42	6.00		
LSP at 5X	(n)11.92.(b)7.54.(c)16	.86	(n)5.71.(h)3.61.(c)8.AB		(a)0,78,(b)0.50.(c)1.11		

	Number of	nods on sain	Axis	Number of	pods/plant.		Rumber of	Rumber of seeds/pod		
	Sambal	daulat	Nean	Saubal	Daulat	Nean	Sauba)	Daulat	Nean	
Γ Γ Γ	53.00	45.50	49.25	185	213	199	11.85	13.75	12.80	
l F	58.00	52.50	55.25	255	264	260	12.00	14.30	13.15	
۱ ۲ ₂	57.00	50.50	53,75	253	244	249	12.85	14.75	13.80	
T ₃	62.00	52.00	57.00	335	418	377	13.65	16.50	15.08	
T 4	64.50	52.00	58.25	369	378	374	12.95	15.30	14.13	
Nean	5R,90	50.50		279	303		12.66	14.92		
LSD at 5X	{a}1.21.(b)0.76.(c)1.71			(a)50.81,	{a}50.81,{b}BS,{c}71.86			{a}1.21,(b)0.76,(c)NS		

Table	2.3.	(Continued).
-------	------	--------------

	Pod weig	ht/plant (g)		Straw weig)	it/plant (g)		1000-see	1000-seed weight (g)		
	Sawbal	Daulat	Kean	Saubal	Daulat	Nean	Sambal	Daulai	Nean	
To	8.07	8.95	8.51	7.97	6.23	7.01	2.47	2.37	2.42	
۲ ₁	12.28	12.56	12.42	11.57	8.55	10.06	2.21	2.37	2.29	
۲ ₂	12.45	13.21	12.83	11.79	9.07	10.38	2.78	2.01	2.40	
۲ _э	17.06	19.24	18.15	17.30	12.94	14.90	2.42	2.87	2.65	
î4	20.09	19.84	19.97	20.64	12.19	16.42	2.74	2.38	2.56	
Near	13.99	14.78		13.82	9.80		2.52	1.40		
LSD at 5X	(a)3.32,(b)RS,(c)4.70			(a)1.20,(b)0.76,(c)1.69			(a)NS,(b)NS,(c)NS			

	Seed-	Seed-husk weight ratio			Harvest index (%)			Seed yield (kg/ba)		
	Saubal	Daulat	Nean	Saubal	Daulat	Nean	Sambal	Daulat	Nean	
To	0.94	1.09	1.02	24.14	30.48	27.31	1356	1353	1355	
T ₁	1.02	1.09	1.06	25.82	31.04	28.43	2025	2165	2095	
۴ ₂	0.95	1.14	1.05	24.93	31.36	28.15	2008	2314	2161	
۳ ₃	1.02	1.12	1.07	25.08	31.06	28.07	2842	3308	3075	
T ₄	0.95	1.12	1.04	23,90	32.56	28.23	3228	3454	3341	
Kean	0.98	1.11		24.77	31 .3 0		2269	2557		
LSD at 5%	(a)WS,(b)0.12,(c)HS			(a)NS,(b))	(a)NS,(b)1.88,(c)NS			(a)594, (b)RS, (c)840		

Table 2.3. (Continued).

	Oil content of	f seeds (%)	
	Sambal	Daulat	Mean
т ₀	32.00	30.50	31.25
т _l	33.50	36.00	34.75
т ₂	32.50	39.00	35.75
т	33.00	33.00	33.00
т ₄	31.00	33.00	32.00
Mean	32.40	34.30	
LSD at 5%	(a)2.40,(b)1.	50,(c)3.40	

- (a) = Difference between treatments
- (b) = Difference between cultivars
- (c) = Difference between interactions

Table 2.4. Correlation coefficients between seed yield and its components as influenced by soil moisture stress at specific stages of growth.

	Seed yield with											
	Plant height at maturity	Number of branches/ plant	Number of pods on main axis	Number of pods/plant	Number of seeds/pod	Pod weight/plant	1000-seed weight	Harvest index	Oil content			
۲o	0.07	0.42*	0.10	0.74***	0.38	0.97***	0.28	0.64***	-0.15			
٢ı	0.30	0.55**	0.47*	0.83***	0.34	0.96***	0.25	0.52**	-0.01			
۲ ₂	~0.08	0.21	0.30	0.80***	0.35	0.96***	-0.20	0.69***	0.18			
۲ ₃	-0.25	0.25	0.19	0.78***	0.25	0.47*	-0.26	0.52**	-0.34			
Ĩ4	-0.11	0.14	0.07	0.85***	0.54**	0.30	-0.18	0.48*	-0.11			

The growth and yield of crop plants are often reduced by soil moisture stress. The crops should be supplied with adequate water before the soil moisture has been depleted to such an extent as to cause the severe reduction in plant growth. When rewatered after a prolonged and severe moisture stress plant will recover its growth. However, yield of crops may be depressed due to soil moisture stress of any degree. The rate of yield reduction will depend on the degree of stress. Plants are not equally sensitive to soil moisture stress at various stages of growth.

In the present experiment, almost all the plants of Sambal and Daulat in the stressed condition yielded reduced TDM than the stress-free plants (T_i) . Maximum reduction of TDM production was found in the plants grown under continuous stress (T_{ij}) . The second highest reduction was found in the flowering to mid-pod filling stress (T_2) . Water shortage significantly affected total dry matter in beans (Farah, 1981). Increase in the water deficits resulted in progressive reductions in the total biomass accumulated (Nageswara Rao et al., 1985). A progressive increase in water stress will first affect the elongation of cells followed by cessation of elongation and marked reduction in cell multiplication and finally the reduction in dry weight of shoot through continued respiration (Jain and Misra, 1972). In groundnut, when the plants were stressed beyond early pod development phase, a loss in biomass production occurred (Ramesh Babu et al., 1984). Others have shown that continuous water deficit reduces dry matter production (Stansell et al.,

1976; Vivekanandan and Gunasena, 1976; Pallas et al., 1979).

Yield and yield components were reduced by soil moisture The height of the plants reduced due to water stress. The stress. continuously stressed plants (T_0) had significantly shorter height than that of the stress-free plants (T_i) . Plants grown under continuous stress (T_0) and pre-flowering stress (T_1) had significantly reduced plant height over the stress-free plants in Sambal. Similar result was found in Corchorus olitorius by Ayodele and Fawusi (1990). Plant height reduction is due to stunted growth of stem. Cell enlargement is so sensitive to water deficits that stem elongation can be inhibited in some species by small diurnal water deficits (Slatyer, 1973). Boote and Hammond (1981) reported that droughts during early pegging and pod formation period in groundnut reduced vegetative growth by reducing both the rate of node formation and by reducing elongation growth. Ramesh Babu et al. (1984) stated that moisture stress imposed at early flowering, at peak flowering and and at early pod development stages in groundnut caused a reduction in plant height. Stress at the late pod development stage did not reduce plant height because the plant elongation was completed by this time. However, continously stressed plants exhibited severe reduction in plant height during the entire life span.

Significantly higher number of branches/plant was found in continuous stress (T_0) over stress-free plants (T_1) . Similar result was found by Ayodele and Fawusi (1990) in *Corchorus capsularis*. The

number of branches/plant in groundnut was reported to be unaffected by moisture deficits (Lenka and Misra, 1973).

In the present investigation, the number of pods/plant was significantly reduced in T_0 , T_1 , and T_2 over T_4 . The cause of this reduction may be due to reduced plant height, length of main axis and number of branches/plant (Table 2.3). It may also be caused due to shedding of flowers by soil drought. Imposing moisture stress at flowering and pod formation stages led to a reduction of 64.20% and 46.92% in pod number compared with control plants (Ayodele and Fawusi, 1990). Reduced number of pods/plant was found due to stress in rapeseed by Richards and Thurling (1978a) and Bhan *et al.*(1980).

The plants grown under T_0 , T_1 and T_2 had significantly reduced pod weight/plant (57%, 42% and 36% respectively) than the plants grown under T_4 (Table 2.3). The main cause of this reduction was due to reduced number of pods/plant. Srinivasan *et al.*(1987) found higher pod yield in the plants grown under stress-free condition.

Straw weight was reduced significantly in $T_0(57\%), T_1(38\%), T_2(36\%)$ and $T_3(9\%)$ over the stress-free plants (T_4) . Similar result was found by Richards and Thurling (1978a) in rapeseed. Reduced straw yield was observed in the plants grown at low soil moisture than that grown at high soil moisture in mustard and chickpea (Singh, 1983).

No significant effect of soil moisture stress on 1000-seed weight was found (Table 2.3). Similar result was found by Bhan *et al.* (1980) in rapeseed. But Richards and Thurling (1978a) found significantly reduced 1000-seed weight in the stressed plants. Soil moisture stress had no clear cut effect on harvest index (Table (2.3). Richards and Thurling (1978a) and Singh (1983) observed similar effect in rapeseed.

Significantly reduced seed yield was found in plants grown under T_0 (59%), T_1 (37%) and T_2 (35%) over plants grown under stressfree condition (T_4). Similar result was reported by Richards and Thurling (1978a)in repessed. Seed yield was reduced in the stressed plants in rapeseed (Bhan *et al.*,1980) and in mustard and chickpea (Singh, 1983). The reason for reduced seed yield may be due to reduced number of pods/plant and weight of pods/plant. Haque *et al.* (1990) observed that in mustard seed yield increased significantly for two irrigations, one at the early vegetative stage and the other at the initial pod formation stage.

Oil content of seeds of the stressed plants grown under T_1 and T_2 was increased by 9 and 12%, respectively than that of the stress-free plants (Table 2.3). But Bhan *et al.*(1980) found reduced oil content in rapeseed due to soil moisture stress.

The overall results of the experiment indicate that TDM, seed yield and its components were highly reduced when moisture stress was imposed during the pre-flowering (T_i) and the flowering to pod filling (T_2) stages. Moisture stress at the post mid-pod filling stage had no significant effect on yield and its components. Thus, it may be concluded that the pre-flowering (T_1) and the flowering to pod filling (T_2) stages are more sensitive to soil moisture stress. Therefore, to obtain better yield it may be recommended that proper soil moisture should be maintained at those two stages for obtaining higher seed yield of mustard.



.

.

REFERENCES

- Allen, E.J. and D.G. Morgan. 1972. A quantitative analysis of the effect of nitrogen on the growth, development and yield of oilseed rape. J. agric. Sci., Camb. 78 : 315-324.
- Allen, E.J. and D.G. Morgan. 1975. A quantitative comparison of the growth, development and yield of different varieties of oilseed rape. J. agric. Sci., Camb. 85 : 159-174.
- Allen, E.J., D.G. Morgan, and W.J. Ridgman. 1971. A physiological analysis of the growth of oilseed rape. J. agric. Sci.,Camb. 77 : 339-341.
- Allen, L.H., Jr. K.J. Boote and L.C. Hammond. 1976. Peanut stomatal diffusion resistance affected by soil water and solar radiation. Proc. Soil Crop Sci. Soc. Fla. 35 : 42-46.
- Andrews, J.E. 1972. Research report. Can. Agri. Res. Sta. Lethbridge. Alta. P. 309.
- Anonymous. 1990. The Statistical Year Book of Bangladesh.Bangladesh Bureau of Statistics. P. 175.
- Arnon, D.I. 1949. Copper enzymes in isolated chloroplast polyphenol oxidase in *Beta vulgaris*. Plant Physiol. 24 : 1-15.
- Asana, R.D. 1961. Analysis of drought resistance in wheat. In Plant Water Relationship In Arid and Semi-Arid Conditions. Arid Zone Res. 16 : 183-190.
- Ayodele, V.I. and M.O.A. Fawusi. 1990. Studies on drought susceptibility of Corchorus olitorius L. II. Effects of moisture stress at different physiological stages on vegetative growth and seed yield of C. olitorius cv. 'Oniyaya'. Biotronics 19 : 33-37.
- Bajpai, M.R., G.D. Singh and B.R. Chhipa. 1981. Effect of sowing time and irrigation levels on growth and yield of mustard. Indian J. agric. Res. 15 : 197-200.

- Banerjee, H.T., M. Das and T.K. Bhattacharjee. 1967. A note on irrigational effect on yield and oil content of Toria (Brassica campestris) and its economics. Indian J. Agron. 12 : 323-324.
- Bannister, P.1964. Stomatal responses of heath plants to water deficits. J. Ecol. 52 : 151-158.
- Bates, L.S., R.P. Waldren and I.D. Teare. 1973. Rapid determination of free proline for water stress studies. Plant and Soil 39 : 205-207.
- Begg, J.E. and N.C. Turner. 1976. Crop water deficits. Adv. Agron. 28 : 161-217.
- Begum, F.A. 1991. Morphological and physiological responses of mustard (*Brassica juncea* L.) to water deficit. M.Sc. Thesis, Rajshahi University, Bangladesh.
- Bhan, S.1979. Effect of soil moisture and nitrogen on mustard under Gangetic alluvium of Uttar Pradesh. Indian J. Agron. 24 : 180-186.
- Bhan, S.1981. Water use, yield and water use efficiency of mustard in relation to variety, soil moisture regime and nitrogen under Gangetic alluvium of Uttar Pradesh. India J. Agron. 27 : 62-65.
- Bhan, S., M. Balaraju and Vidya Ram. 1980. Water use, yield and guality of rape seed as influenced by spacing, irrigation and time of harvest when raised in a multi-cropping system. Indian J. agric. Sci. 50 : 760-763.
- Bhati, T.K. and S.S. Rathor. 1982. Response of Indian mustard to irrigation and nitrogen fertilizer. Indian J. Agron. 27 : 451-453.
- Black, C.R., D.Y. Tang, C.K. Ong, A. Solon and L.P. Simmonds. 1985. Effects of soil moisture stress on the water relations and water use of groundnut stands. New Phytol. 100 : 313-328.

- Blum, A. and A. Ebercon. 1976. Genotypic responses in sorghum to drought stress. III. Free proline accumulation and drought resistance. Crop Sci. 16 : 428-431.
- Boote, K.J. and L.C. Hammond. 1981. Effect of drought on vegetative and reproductive development of peanut. Proc. Am. Peanut Res. Educ. Soc. 13 : p. 86.
- Buttery, B.R. 1969. Analysis of the growth of soybeans as affected by plant population and fertilizer. Can. J. Plant Sci. 49 : 675-684.
- Chanda, S.V., A.K. Joshi, P.P. Vaishnav and Y.D. Singh. 1987.Growth analysis using classical and curve-fitting methods in relation to productivity in pearl millet (*Pennisetum Americanum* L.). J. Agron. & Crop Sci. 159 : 312-319.
- Chaniara, N.J. and U.M. Damor. 1982. Effect of irrigation intervals and various levels of nitrogen and phosphorus on the yield of mustard variety Varuna. Indian J. Agron. 27 : 472-474.
- Chaturvedi, G.S., B.B. Singh, R. Prasad, Y.S. Chauhan and Padmakar. 1988. Physiological analysis of yield in Indian mustard (*Brassica juncea* L.) under irrigated conditions. Indian J. Plant Physiol. 31 : 38-44.
- Ciha, A.J. and W.A. Brun. 1975. Stomatal size and frequency in soybeans. Crop Sci. 15 : 309-313.
- Clarke, J.M. and G.M. Simpson. 1978a. Growth analysis of *Brassica* napus cv. Tower. Can. J. Plant Sci. 58 : 587-595.
- Clarke, J.M. and G.M. Simpson. 1978b. Influence of irrigation and seeding rate on yield and yield components of *Brassica napus* cv. Tower. Can. J. Plant Sci. 58 : 731-737.
- Clarke, J.M. and G.M. Simpson. 1978c. Leaf osmotic potential as an indicator of crop water deficit and irrigation in rape seed (*Brassica napus* L.). Agric. Water Manage. I : 351-356.
- Clarke, J.M. and G.M. Simpson. 1979. The application of a curvefitting technique to *Brassica napus* growth data.Feild Crops Res. 2 : 35-43.

- Connor, D.J. and T.R. Jones. 1985. Response of sunflower to strategies of irrigation. II. Morphological and physiological response to water stress. Field Crops Res. 12 : 91-103.
- Constable, G.A. and A.B. Hearn. 1978. Agronomic and physiological responses of soybean and sorghum to water deficits. I. Growth, development and yield. Aust. J. Plant Physiol. 5 : 159-167.
- Dastane, N.G. 1972. Practical Manual for Water Use Research in Agriculture. 2nd ed. Navabharat Prokashan, Poona, India.
- Denna, D.W. 1970. Transpiration and the waxy bloom in Brassica oleracea L. Aust. J. Biol. Sci. 23 : 27-31.
- Donald, C.M. 1961. Competition for light in crops and pastures. In Mechanisms in Biological Competition. Academic Press, Inc., New York. Soc. Exp. Biol. Symp. 15 : 282-313.
- Dornhoff, G.M. 1971. Net photosynthesis of soybean leaves as influenced by anatomy, respiration and variety. Ph.D. Thesis, Iowa State University.
- Dornhoff, G.M. and R. Shibles. 1976. Leaf morphology and anatomy in relation to CO₂ exchange rate of soybean leaves. Crop Sci. 16 : 377-381.
- Dunhum, R.J. and P.H. Nye. 1976. The influence of soil water content on the uptake of ions by roots. J. appl. Ecol. 13 : 967-984.
- Eagles, C.F. 1969. Time changes of relative growth rate in two natural populations of *Dactylis glomerata* L. Ann. Bot. 33 : 937-946.
- Eagles, C.F. 1971. Changes in net assimilation rate and leaf area ratio with time in *Dactylis glomerata* L. Ann. Bot. 35: 63-74.
- Easten, F.M. and D.R. Ergle. 1948. Carbohydrate accumulation in cotton plants at low moisture levels. Plant Physiol. 23: 169-187.
- El-Fouly, M.M., H.A. El-Hammawi and A.F.A. Fawzi. 1971. Different effects of chlormequat on chlorophyll, protein and yield in cotton grown under varying water regimes. Plant and Soil 35 : 183-186.
- El Nadi, A.H. 1969. Water relations of beans. I. Effects of water stress on growth and flowering. Expl. Agric. 5 : 195-207.
- El-Saidi, M.T. and A.A. El-Mossallami. 1972. The effect of different levels of water supply on transpiration coefficient and photosynthetic pigments of cotton plant. Z. Acker-und Pflanzenbau. 135: 135-142.
- Farah, S.M. 1981. An examination of the effects of water stress on leaf growth of crops of field beans (Vicia faba L.) I. Crop growth and yield. J. agric. Sci., Camb. 96 : 327-336.
- Freyman, S., W.A. Charnetski and R.K. Crookston.1973.Role of leaves in the formation of seed in rape. Can. J. Plant Sci. 53 : 693-694.
- Gardner, W.R. 1960.Dynamic aspects of water availability to plants. Soil Sci. 89 : 63-73.
- Hall, A.E. and E.D. Schulze. 1980. Drought effects on transpiration and leaf water status of cowpea in controlled environments. Aust. J. Plant Physiol. 7 : 141-147.
- Hanson, A.D., C.E. Nelson, A.R. Pederson and E.H. Everson. 1979. Capacity for proline accumulation during water stress in barley and its implications for breeding for drought resistance. Crop Sci. 19 : 489-493.
- Haque, M.R., M.U. Ahmed and M.A. Rahman. 1987. Irrigation scheduling for optimum yield of mustard. Bangladesh J. Agril. Sci. 14 : 31-34.
- Harper, P. 1963. Optimum leaf area index in the potato crop. Nature (Lond.) 197 : 917-918.
- Henry, J.L. and K.B. MacDonald. 1978. The effect of soil and fertilizer nitrogen and moisture stress on yield, oil and protein content of rape. Can. J. Soil Sci. 58 : 303-310.

- Hozyo, Y., S. Kato and H. Kobayashi. 1972. Photosynthetic activity of the pods of rape plants (*Brassica napus* L.) and the contribution of pods to the ripening of rape seeds (In Japanese). Proc. Crop Sci. Soc. Japan 41: 420-425. (English Abstr. in Field crop Abstr. 26: 708. 1973).
- Hsiao, T.C. and E. Acevedo. 1974.Plant responses to water deficits, water use efficiency and drought resistance. Agric. Mete. 14 : 59-84.
- Hughes, A.P. and P.R. Freeman. 1967. Growth analysis using frequent small harvests. J. appl. Ecol. 4 : 553-560.
- Hunt, R. 1978. Plant growth analysis. Studies in Biology No. 96 Edward Arnold, London, p. 67.
- Hurd, R.G. 1977. Vegetative plant growth analysis in controlled environments. Ann. Bot. 41 : 779-787.
- Hussein, M.M. 1973. Water relation of cotton plants and its effect on growth and shedding. Ph.D. Thesis. Fac. Agric., Ain Shams University.
- Hygen, G. 1951. Studies in plant transpiration. Physiol. Plant. 4 : 57-83.
- Inanga, S. and A. Kumara. 1974.Studies on dry matter production of rape plant (Brassica napus L.) I. Changes with growth in rates of photosynthesis and respiration of rape plant population (In Japanese, English Summary). Proc. Crop Sci. Jap. 43 : 267-277.
- Jain, T.C. and K.C.Jain. 1979. Response of Indian mustard to levels of irrigation and fertility in light-textured soils of Rajsthan. Indian J. agric. Sci. 49 : 470-472.
- Jain, T.C. and D.K. Misra. 1972. Moisture stress effect on leaf growth of groundnut. Indian J. Agron. Sci.17 : 283-286.
- Jarvis, P.G. and M.S. Jarvis. 1963. The water relations of tree seedlings. IV. Some aspect of tissue water relations and drought resistance. Physiol. Plant. 16 : 501-516.

- Joarder, O.I., N.K. Paul and S.K. Ghose.1979. Effects of irrigation and fertilizer on mustard (*Brassica juncea*). Expl. Agric. 15 : 299-302.
- Joshi, Y.C., P.C. Nautial, V. Ravindra and R. Snehi Dwivedi. 1988. Water relations in two cultivars of groundnut (Arachis hypogaea L.) under soil water deficit.Trop. Agric.(Trinidad) 65 : 182-184.
- Kaufmann, M.R. 1972. Water deficits and reproductive growth. In: Water Deficits and Plant Growth. T.T. Kozlowski (Ed). Academic Press, New York and London, vol.3, 91-124.
- Kaul, A.K. and M.L. Das. 1986. Oilseeds in Bangladesh. Bangladesh-Canada Agri. Sector team, Ministry of Agri. Govt. P.R. Bangladesh, Dhaka.
- Kemble, A.R. and H.T. MacPherson. 1954. Liberation of amino acids in perennial ryegrass during wilting. Biochem. J.58: 46-49.
- Khan, G.M. and S.K. Agarwal. 1985. Influence of sowing methods, moisture stress and nitrogen levels on growth, yield components and seed yield of mustard. Indian J. agric. Sci. 55 : 324-327.
- Kirby, E.J.M. 1969. The growth and development of some barley varieties in response to irrigation and nitrogen fertilizer. J. agric. Sci., Camb. 72 : 467-474.
- Koller, H.R., E.W. Nyquist and I.S. Chorush. 1970. Growth analysis of the soybean community. Crop Sci. 10 : 407-412.
- Kramer, P.J. 1969. Plant and soil water relationship : A modern synthesis. McGraw-Hill, New York.
- Krogman, K.K. and E.H. Hobbs.1975. Yield and morphological responses of rape (Brassica campestris L.cv. Span) to irrigation and fertilizer treatments. Can. J. Plant Sci. 55: 903-909.
- Larson, K.L. 1975. Drought injury and resistance of crop plants. In : Physiological Aspects of Dryland Farming. Gupta,U.S.(Ed.). Oxford and IBH Pub. Co., New Delhi. pp. 145-164.

- Lenka, D. and P.K. Misra. 1973. Response of groundnut (Arachis hypogaea L.) to irrigation. Indian J. Agron. 18 : 492-497.
- Lochwing, W.F. 1940. Flower and pod-shedding of soybean in relation to water and nutrient deficiency. Science 92 : p. 517.
- Mackinney, G. 1941. Absorption of light by chlorophyll solutions. J. Biol. Chem. 140 : 315-319.
- Maini, N.S., J.N. Sandhu and K.S. Johal. 1964. Effect of sowing date, spacing, irrigation and nitrogen levels on the grain yield and growth of Toria (*Brassica campestris*). Indian Oilseeds J. 8 : 128-132.
- Manalo, E.B. 1975. Agro-climatic survey of Bangladesh. P. 240.
- Mandal, B.K., P.K. Ray and S. Dasgupta. 1986. Water use by wheat, chickpea and mustard grown as sole crops and intercrops. Indian J. agric. Sci. 56 : 187-193.
- Mare, J. and A. Palmer. 1976. Effect of water stress on leaf growth of sunflower. Physiol. Plant 36 : 101-104.
- Mathur, O.P. and P.S. Tomar. 1972. Irrigation requirement of raya (Brassica juncea) crop in Western Rajsthan.Indian J. Agron. 17 : 306-308.
- Mehrotra, O.N., R.C. Garg and S.A. Ali. 1969. Influence of soil moisture on growth, yield and quality of groundnut. Indian J. Plant Sci. 12 : 58-63.
- Milbourn, G.M. and R.C. Hardwick. 1968. The growth of vining peas. I. The effect of time of sowing. J. agric. Sci.,Camb. 70 : 393-402.
- Miller, E.C. 1938. Plant Physiology. E.W. Sinott (ed.). McGraw-Hill Book Co., Inc. New York.
- Milthorpe, F.L. 1969. The significance and mechanism of stomatal movement. Aust. J. Sci. 32 : 31-35.

- Moursi, M.A., O.H. El-Bagoury and A.M. Shaheen.1976. Photosynthetic pigments in leaves of some field crops as affected by water deficit in soil. J. Agron. & Crop Sci. 142 : 206-213.
- Nagarajah, S. and E.D. Schulze.1983. Responses of Vigna unguiculata (L.) Walp. to atmospheric and soil drought. Aust. J. Plant Physiol. 10 : 385-394.
- Nageswara Rao, R.C., Sardar Singh, M.V.K. Sivakumar, K.L. Srivastava and J.H. Williams.1985.Effect of water deficit at different growth phases of peanut.1. Yield responses. Agron. J. 77 : 782-786.
- Nageswara Rao, R.C., J.H. Williams, M.V.K. Sivakumar and K.D.R. Wadia. 1988. Effect of water deficit at different growth phases of peanut. II. Response to drought during preflowering phase. Agron. J. 80 : 431-438.
- Narasimha Rao, C.L. and A. Shiv Raj. 1985. Sugar and proline accumulation in glossy and non-glossy cultivars of foxtail millet grown under water stress. Indian J. agric. Sci. 55 : 183-185.
- Narasimha Rao, C.L. and A. Shiv Raj. 1988. Effect of water stress on grain growth of glossy and non-glossy varieties of grain sorghum (Sorghum bicolor). Indian J. agric. Sci. 58 : 770-773.
- Nash, D., L.G. Paleg and J.T. Wiskich. 1980. The effect of proline and betaine on the heat stability of some mitochondrial enzymes. Proc. 21st Gen. Meet. Aust. Soc. Plant Physiol. Monash University, Melbourne, 13-16 May, Abstr. No. 63.
- Nathawat, K.S., P. Joshi. L.R. Bagrecha and P.D. Bhargava. 1969. Influence of soil drought on the relation between yield and yield components in *Brassica campestris* L. var. Sarson Prain. Indian J. agric. Sci. 39 : 397-399.
- Nayak, R.L., S.B. Roy, Gopeswar Sounda and S.R. Mandal. 1983. Relative turgidity of leaf tissue of wheat as affected by different soil moisture stress applied at various stages of growth. Bangladesh J. Agril. Sci. 10 : 215-216.

- Nazrul-Islam, A.K.M. and A. Yasmin. 1982. Tissue water relations in tomato grown in different soil moisture regimes. Geobios 10 : 49-53.
- Nazrul-Islam, A.K.M. and M. Alam. 1986. Comparative study of water relation in two cultivars of jute under waterlogged and non-waterlogged condition. Geobios 13 : 174-177.
- Nerkar, Y.S., D. Wilson and D.A. Lawes. 1981. Genetic variation in stomatal characteristics and behaviour,water use and growth of five *Vicia faba* L. genotypes under contrasting soil moisture regimes. Euphytica 30 : 335-345.
- Nicholls, A.O. and D.M. Calder. 1973. Comments on the use of regression analysis for the study of plant growth. New Phytol. 72 : 571-581.
- Norton, G. and J.F. Harris.1975.Compositional changes in developing rapeseed (*Brassica napus* L.). Planta 123 : 163-174.
- Oaks, A., D.J. Mitchell, R.A. Barnard and F.C. Johnson. 1970. The regulations of proline biosynthesis in maize roots. Can. J. Bot. 48 : 2249-2258.
- Palfi, G., M. Bito and J. Palfi. 1973. Free proline and water deficits. Sov. Plant Physiol. 20 : 189-193.
- Pallas, J.E. Jr., J.R. Stansell and T.J. Koske. 1979. Effects of drought on florunner peanuts. Agron. J. 71 : 853-858.
- Pandey, R.K., M.C. Saxena and V.B. Singh. 1978. Growth analysis of blackgram genotypes. Indian J. agric. Sci. 48 : 466-473.
- Pandey, R.P., P.K.R. Nair and J.P. Tiwari. 1981. Correlation of morpho-physiological and sink parameters in cowpea. Indian J. agric. Sci. 51 : 221-224.
- Penfound, W.T. 1931.Plant anatomy as conditioned by light intensity and soil moisture. Amer. J. Bot. 18 : 558-572.
- Protsenko, D.F., I.G. Shamat'ko and E.A. Rubanyuk. 1968. Drought hardiness of winter varieties as related to their amino acid composition. Sov. Plant Physiol. 15 : 567-573.

- Quarrie, S.A. 1980. Genotypic differences in leaf water potential, abscisic acid and proline concentrations in spring wheat during drought stress. Ann. Bot. 46 : 383-394.
- Rabindranath, P. and A. Shiv Raj.1983. Effect of moisture stress on growth, yield and yield components of field-grown sorghum varieties having glossy and non-glossy leaves. Indian J. agric. Sci. 53 : 428-430.
- Radford, P.J. 1967. Growth analysis formulae their use and abuse. Crop Sci. 7 : 171-175.
- Rahman, L. 1978. Studies on The Production Conditions of Mustard and Rapeseed at Farmers Level in 10 Police Stations of Comilla District Oilcrops Survey 2 : 16.
- Raja, V. and K.C. Bishnoi. 1990. Evapotranspiration, water use efficiency, moisture extraction pattern and plant water relations of rape (*Brassica campestris*) genotypes in relation to root development under varying irrigation schedules. Expl. Agric. 26 : 227-233.
- Ramesh Babu, V., P.S.S. Murty and D.N. Reddy. 1984. Effect of water stress at specific growth stages on growth and yield of groundnut. Ann. Arid Zone. 23 : 13-20.
- Reddy, B.N. and M.N. Sinha. 1987. Effect of irrigation and fertilization on yield and economics of Indian mustard. J. Oilseeds Res. 4 : 202-210.
- Reddy, B.N., M.N. Sinha and M.R. Hedge.1988. Studies on consumptive water use, water use efficiency and moisture extraction pattern by mustard as influenced by irrigation and fertilization. J. Oilseeds Res. 5 : 36-44.
- Rees, A.R. 1963.Relationship between crop growth rate and leaf area index in the oil palm. Nature (Lond.) 197 : 63-64.
- Richards, R.A. 1978. Variation between and within species of rapeseed (*Brassica campestris* and *B. napus*) in response to drought stress. III. Physiological and physicochemical characters. Aust. J. Agric. Res. 29 : 491-501.

- Richards, R.A. and N. Thurling. 1978a. Variation between and within species of rapeseed (*Brassica campestris* and *B. napus*) in response to drought stress. 1. Sensitivity at different stages of development. Aust. J. Agric. Res. 29 : 469-477.
- Richards, R.A. and N. Thurling. 1978b. Variation between and within species of rapeseed (*Brassica campestris* and *B. napus*) in response to drought stress.II. Growth and development under natural drought stresses.Aust. J. Agric. Res. 29: 479-490.
- Richards, R.A. and N. Thurling. 1978c. Genetic analysis of drought stress response in rapeseed (*Brassica campestris* and *B. napus*). III. Physiological characters. Euphytica 28: 755-759.
- Saha, D.K. 1983. Physiological studies on the growth, development and yield of two cultivated species of jute (Corchorus capsularis L. and C. olitorius L.). M.Sc. Thesis, Rajshahi University.
- Saini, A.D. and K. Das. 1979. Studies on growth and yield of three mungbean (Vigna radiata L.) cultivars. Indian J. Plant Physiol. 22 : 147-153.
- Sarkar, M.A.R., M.R. Ahsan, A.U. Sarker and S.M.A. Hossain. 1989. Effect of nitrogen and irrigation at different growth stages on mustard. Bangladesh J. Agril. Sci. 16 : 283-288.
- Sarker, A.A., M. Ahmed, M.S.U. Talukder and M.R. Khan. 1990. Yield and water use of mustard under different irrigation regimes. Bangladesh J. Agril. Sci. 17 : 293-298.
- Scott, R.K., E.A. Ogunremi, J.D. Ivins and N.J. Mendham. 1973. The effect of sowing date and season on growth and yield of oil seed rape (*Brassica napus*). J. agric. Sci., Camb. 18 : 277-285.
- Sharma, D.K. and A. Kumar. 1989. Effecct of irrigation on growth analysis, yield and water use in Indian mustard (*Brassica juncea* subsp. *juncea*). Indian J. agric. Sci. 59 : 162-165.
- Singh, B.P. 1983. Response of mustard and chickpea to moisture in soil profile and plant population on aridisols. Indian J. agric. Sci. 53 : 543-549.

- Singh, R.A. and H.R. Singh. 1978. Effect of nitrogen and phosphorus on yield, quality and moisture-use pattern of oilseed grown on rainfed lands. Indian J. agric. Sci. 48 : 583-588.
- Singh, R.A., S.C. Gupta and S. Moinbasha. 1987. Moisture needs of Indian mustard-chickpea intercropping. Indian J. agric. Sci. 57 : 553-558.
- Singh, V.B. and S.P. Tomar. 1971. Response of mustard to varying irrigation levels, spacing and fertilizer applications. Indian J. Agron. 16 : 465-467.
- Sivakumar, M.V.K. and R.H. Shaw. 1978. Methods of growth analysis in field-grown soyabeans (Glycine max (L.) Merill). Ann. Bot. 42 : 213-222.
- Sivakumar, M.V.K., N. Seetharama, Sardar Singh. and F.R. Bidinger. 1979. Water relations, growth and dry matter accumulation of sorghum under post-rainy season condition. Agron. J.71: 843-847.
- Slatyer, R.O. 1973. The effect of internal water status on plant growth, development and yield. *In* : Plant Response to Climatic Factors. Proc. Uppsala Symp. pp. 177-191.
- Srinivasan, P.S., R. Sathasivam, A. Arjunan, R. Sethupathi Ramalingam and M. Vaman Bhat. 1987. Effect of water stress on partitioning of dry matter and crop growth rate in relation to productivity in groundnut cultivars. J. Oilseeds Res.4 : 89-96.
- Stansell, J.R., J.L. Shepherd, J.E. Pallus, B.R. Bruce,N.A. Minton, D.K. Bell and L.W. Morgan. 1976. Peanut responses to soil water available in the south east. Peanut Sci. 3 : 44-48.
- Stewart, C.R. 1971. Effect of wilting on carbohydrate during incubation of excised bean leaves in the dark. Plant Physiol. 48 : 792-794.
- Stewart, C.R. 1972. The effect of wilting on proline metabolism in excised bean leaves in the dark.Plant Physiol. 51 : 508-511.

- Stewart, C.R., C.J. Morris and J.F. Thompson.1966. Changes in amino acid content of excised leaves during incubation. II. Role of sugar in the accumulation of proline in wilted leaves. Plant Physiol. 41 : 1585-1590.
- Tan, Geok-Yong, Wei-koon Tan and P.D. Walton.1978. Seedling growth of smooth bromegrass measured in five environments. Crop Sci. 18 : 601-604.
- Tayo, T.O. and D.G. Morgan. 1975. Quantitative analysis of the growth, development and distribution of flowers and pods in oil seed rape (*Brassica napus* L.). J. agric. Sci., Camb. 85: 103-110.
- Teare, I.D., and Kanemasu. 1972. Stomatal diffusion resistance and water potential of soybean and sorghum leaves. New Phytol. 71 : 805-810.
- Thorne, G.N. 1960. Variation with age in net assimilation rate and other growth attributes sugar beet, potato and barley in a controlled environment. Ann. Bot. 24 : 256-272.
- Thorne, G.N. 1961. Effects of age and environment on net assimilation rate of barley. Ann. Bot. 25 : 29-38.
- Thorne, J.H. and H.R. Koller. 1974. Influence of assimilate demand on photosynthesis, diffusion resistance, translocation and carbohydrate levels of soybean leaves. Plant Physiol. 54 : 201-207.
- Thurling, N. 1974. Morphophysiological determinants of yield in rapeseed (*Brassica campstris and Brassica napus*). I. Growth and morphological characters. Aust. J. Agric. Res. 25: 697-710.
- Turk, K.J., A.E. Hall and C.W. Asbell. 1980. Drought adaptation of cowpea. I. Influence of drought on seed yield: Agron. J. 72 : 413-420.
- Turner, N.C., J.E. Begg and M.L. Tonnet. 1978. Osmotic adjustment of sorghum and sunflower crops in response to water deficits and its influence on the water potential at which stomata close. Aust. J. Plant Physiol. 5 : 597-608.

- Vivekanandan, A.S. and H.P.M. Gunasena. 1976. Lysimetric studies on the effect of soil moisture tension on the growth and yield of maize (Zea mays L.) and groundnut (Arachis hypogaea L.). Beitr. Trop. Landwirtsch. Veterinaermed. 14 : 369-378.
- Waldren, R.P. and I.D. Teare. 1974. Free proline accumulation in drought stressed plants under laboratory conditions. Plant and Soil 40 : 689-692.
- Waldren, R.P., Teare and S.W. Ehrler. 1974. Changes in free proline concentration in sorghum and soybean plants under field conditions. Crop Sci. 14 : 447-450.
- Wallace, D.M. and Munger. 1965. Studies on the physiological basis for yield differences. I. Growth analysis of six dry bean varieties. Crop Sci. 5 : 343-348.
- Wang, D. 1968. Metabolism of ¹⁴C-labelled proline in higher plants. Contr. Boyce Thompson Inst. Plant Res. 24 : 117-122.
- Wankhede, N.P., I.P.S. Ahlawat and V.M. Sahni. 1970. Effect of varying levels of irrigation, plant population and fertilizer on yield and quality of Indian rape (Brasseca campestris L. var. Toria Duth. and Full.). Indian J. agric. Sci. 40: 873-877.
- Wardlaw, I.F. 1967. The effect of water stress on translocation in relation to photosynthesis and growth. I. Effect during grain development in wheat. Aust. J. biol. Sci. 20 : 25-39.
- Wardlaw, I.F. 1968. The control and pattern of movement of carbohydrates in plants. Botanical Review 34 : 79-105.
- Wardlaw, I.F. 1971. The early stages of grain development of wheat: Respose to water stress in a single variety. Aust. J. biol. Sci. 24 : 1047-1055.
- Wareing, P.F. and J. Patric. 1975. Source-sink relations and the partition of assimilates in the plant. *In*: Photosynthesis and productivity in different environments. J.P.Cooper (Ed.). Cambridge University Press, London. pp. 481-499.

- Watson, D.J. 1947. Comparative physiological studies on the growth of field crops. I. Variation in net assimilation rate and leaf area between species and varieties and within and between years. Ann. Bot. 11 : 41-76.
- Watson, D.J. 1952. The physiological basis of variation in yield. Adv. Agron. 4 : 101-145.
- Watson, D.J. 1958. The dependence of net assimilation rate on leaf area index. Ann. Bot. 22 : 37-54.
- Watson, D.J. 1971. Size, structure and activity of the productive system of crops. In : Potential Crop Production. P.F. Wareing and J.P. Cooper (Eds). Weinman Education Books, London.
- Weatherley, P.E. 1950. Studies in the water relations of cotton. I. The field measurement of water deficits in leaves.New Phytol. 49 : 81-87.
- Wein, H.C., S.T. Littleton and T.C. Ayanab. 1979. Drought stress of cowpea and soybean under tropical conditions. In: Stress Physiology in Crop Plants. C.H. Musell and R.C.Staples (Eds). Wiley Inc. New York. pp. 283-302.
- Williams, W.A., R.S. Loomis and C.R. Lepley.1965. Vegetative growth of corn as affected by population density. II. Components of growth, net assimilation rate and leaf area index. Crop Sci. 5 : 215-219.

Rajsham - - ty Library Poet e se section DUCULALIN. D-2644 Date 7:8:94 D-1744